



wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 4, 2024 – 11:20 am GMT

PDB ID : 5BXL
Title : Yeast 20S proteasome beta2-G170A mutant
Authors : Huber, E.M.; Groll, M.
Deposited on : 2015-06-09
Resolution : 2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3140 (2.80-2.80)
Clashscore	141614	3569 (2.80-2.80)
Ramachandran outliers	138981	3498 (2.80-2.80)
Sidechain outliers	138945	3500 (2.80-2.80)
RSRZ outliers	127900	3078 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	250	 2% 97%
1	O	250	 2% 96%
2	B	258	 3% 85% 9% • 5%
2	P	258	 4% 86% 8% • 5%
3	C	254	 4% 86% 7% • 6%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	Q	254	4% 87% 7% • 6%
4	D	260	% 85% 5% • 10%
4	R	260	% 85% 5% 10%
5	E	234	% 91% 6% ..
5	S	234	% 90% 8% .
6	F	288	2% 80% 5% 16%
6	T	288	% 79% 5% 16%
7	G	252	89% 6% .
7	U	252	2% 89% 6% .
8	H	232	3% 78% 16% • .
8	V	232	3% 77% 16% • • .
9	I	205	89% 10%
9	W	205	% 92% 8%
10	J	198	2% 89% 9% ..
10	X	198	2% 89% 9% ..
11	K	212	93% 5% .
11	Y	212	93% 5% .
12	L	222	89% 10% .
12	Z	222	90% 9% .
13	M	246	87% 5% • 7%
13	a	246	92% • 6%
14	N	196	% 94% 5% .
14	b	196	% 97% .

2 Entry composition

There are 17 unique types of molecules in this entry. The entry contains 49787 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Proteasome subunit alpha type-2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	250	Total	C	N	O	S	0	0	0
			1915	1219	315	377	4			
1	O	250	Total	C	N	O	S	0	0	0
			1915	1219	315	377	4			

- Molecule 2 is a protein called Proteasome subunit alpha type-3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	244	Total	C	N	O	S	0	0	0
			1904	1201	321	379	3			
2	P	244	Total	C	N	O	S	0	0	0
			1904	1201	321	379	3			

- Molecule 3 is a protein called Proteasome subunit alpha type-4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	C	240	Total	C	N	O	S	0	0	0
			1881	1176	329	372	4			
3	Q	240	Total	C	N	O	S	0	0	0
			1881	1176	329	372	4			

- Molecule 4 is a protein called Proteasome subunit alpha type-5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
4	D	235	Total	C	N	O	S	0	0	0
			1813	1136	304	366	7			
4	R	235	Total	C	N	O	S	0	0	0
			1813	1136	304	366	7			

- Molecule 5 is a protein called Proteasome subunit alpha type-6.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
5	E	231	Total 1773	C 1114	N 307	O 348	S 4	0	0	0
5	S	231	Total 1773	C 1114	N 307	O 348	S 4	0	0	0

- Molecule 6 is a protein called Probable proteasome subunit alpha type-7.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
6	F	243	Total 1892	C 1203	N 329	O 356	S 4	0	0	0
6	T	243	Total 1892	C 1203	N 329	O 356	S 4	0	0	0

- Molecule 7 is a protein called Proteasome subunit alpha type-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
7	G	241	Total 1907	C 1214	N 320	O 365	S 8	0	0	0
7	U	241	Total 1907	C 1214	N 320	O 365	S 8	0	0	0

- Molecule 8 is a protein called Proteasome subunit beta type-2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
8	H	222	Total 1685	C 1062	N 293	O 323	S 7	0	0	0
8	V	222	Total 1685	C 1062	N 293	O 323	S 7	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
H	170	ALA	GLY	conflict	UNP P25043
V	170	ALA	GLY	conflict	UNP P25043

- Molecule 9 is a protein called Proteasome subunit beta type-3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
9	I	204	Total 1581	C 1010	N 258	O 305	S 8	0	0	0
9	W	204	Total 1581	C 1010	N 258	O 305	S 8	0	0	0

- Molecule 10 is a protein called Proteasome subunit beta type-4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
10	J	195	Total	C	N	O	S	0	0	0
			1561	992	264	299	6			
10	X	195	Total	C	N	O	S	0	0	0
			1561	992	264	299	6			

- Molecule 11 is a protein called Proteasome subunit beta type-5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
11	K	212	Total	C	N	O	S	0	0	0
			1644	1045	280	312	7			
11	Y	212	Total	C	N	O	S	0	0	0
			1644	1045	280	312	7			

- Molecule 12 is a protein called Proteasome subunit beta type-6.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
12	L	222	Total	C	N	O	S	0	0	0
			1757	1115	303	335	4			
12	Z	222	Total	C	N	O	S	0	0	0
			1757	1115	303	335	4			

- Molecule 13 is a protein called Proteasome subunit beta type-7.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
13	M	229	Total	C	N	O	S	0	0	0
			1790	1133	306	344	7			
13	a	232	Total	C	N	O	S	0	0	0
			1815	1148	311	349	7			

- Molecule 14 is a protein called Proteasome subunit beta type-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
14	N	196	Total	C	N	O	S	0	0	0
			1512	955	250	300	7			
14	b	196	Total	C	N	O	S	0	0	0
			1512	955	250	300	7			

- Molecule 15 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
15	G	1	Total Mg 1 1	0	0
15	H	1	Total Mg 1 1	0	0
15	I	2	Total Mg 2 2	0	0
15	K	1	Total Mg 1 1	0	0
15	L	1	Total Mg 1 1	0	0
15	N	1	Total Mg 1 1	0	0
15	W	1	Total Mg 1 1	0	0
15	Z	1	Total Mg 1 1	0	0

- Molecule 16 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
16	G	1	Total Cl 1 1	0	0
16	U	1	Total Cl 1 1	0	0



- Molecule 17 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
17	A	15	Total O 15 15	0	0
17	B	15	Total O 15 15	0	0
17	C	16	Total O 16 16	0	0
17	D	15	Total O 15 15	0	0
17	E	13	Total O 13 13	0	0
17	F	15	Total O 15 15	0	0
17	G	21	Total O 21 21	0	0
17	H	23	Total O 23 23	0	0



Continued on next page...

Continued from previous page...



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
17	I	21	Total 21	O 21	0	0
17	J	20	Total 20	O 20	0	0
17	K	25	Total 25	O 25	0	0
17	L	29	Total 29	O 29	0	0
17	M	37	Total 37	O 37	0	0
17	N	20	Total 20	O 20	0	0
17	O	9	Total 9	O 9	0	0
17	P	12	Total 12	O 12	0	0
17	Q	15	Total 15	O 15	0	0
17	R	8	Total 8	O 8	0	0
17	S	7	Total 7	O 7	0	0
17	T	12	Total 12	O 12	0	0
17	U	21	Total 21	O 21	0	0
17	V	18	Total 18	O 18	0	0
17	W	19	Total 19	O 19	0	0
17	X	22	Total 22	O 22	0	0
17	Y	21	Total 21	O 21	0	0
17	Z	18	Total 18	O 18	0	0
17	a	29	Total 29	O 29	0	0
17	b	25	Total 25	O 25	0	0




- Molecule 10: Proteasome subunit beta type-4




- Molecule 10: Proteasome subunit beta type-4




- Molecule 11: Proteasome subunit beta type-5





- Molecule 11: Proteasome subunit beta type-5




- Molecule 12: Proteasome subunit beta type-6




- Molecule 12: Proteasome subunit beta type-6





- Molecule 13: Proteasome subunit beta type-7

Chain M:  87% 5% • 7%




- Molecule 13: Proteasome subunit beta type-7

Chain a:  92% • 6%




- Molecule 14: Proteasome subunit beta type-1

Chain N:  94% 5% •



- Molecule 14: Proteasome subunit beta type-1

Chain b:  97% •



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	134.80Å 300.94Å 144.49Å 90.00° 112.71° 90.00°	Depositor
Resolution (Å)	15.00 – 2.80 15.00 – 2.80	Depositor EDS
% Data completeness (in resolution range)	94.6 (15.00-2.80) 94.6 (15.00-2.80)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.68 (at 2.81Å)	Xtrriage
Refinement program	REFMAC 5.7.0032	Depositor
R, R_{free}	0.189 , 0.214 0.198 , 0.223	Depositor DCC
R_{free} test set	12204 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	54.5	Xtrriage
Anisotropy	0.127	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 45.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	49787	wwPDB-VP
Average B, all atoms (Å ²)	61.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.55% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CL, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.25	0/1952	0.48	0/2642
1	O	0.27	0/1952	0.49	0/2642
2	B	0.27	0/1934	0.51	0/2618
2	P	0.26	0/1934	0.51	0/2618
3	C	0.26	0/1910	0.51	0/2586
3	Q	0.26	0/1910	0.51	0/2586
4	D	0.25	0/1837	0.48	0/2475
4	R	0.25	0/1837	0.48	0/2475
5	E	0.25	0/1800	0.48	0/2433
5	S	0.25	0/1800	0.48	0/2433
6	F	0.26	0/1932	0.45	0/2609
6	T	0.26	0/1932	0.45	0/2609
7	G	0.26	0/1945	0.47	0/2634
7	U	0.28	0/1945	0.47	0/2634
8	H	0.35	0/1716	0.55	0/2328
8	V	0.36	0/1716	0.57	0/2328
9	I	0.30	0/1611	0.52	0/2174
9	W	0.27	0/1611	0.50	0/2174
10	J	0.26	0/1589	0.52	0/2142
10	X	0.26	0/1589	0.52	0/2142
11	K	0.26	0/1681	0.53	1/2274 (0.0%)
11	Y	0.26	0/1681	0.53	1/2274 (0.0%)
12	L	0.29	0/1795	0.54	0/2420
12	Z	0.30	0/1795	0.54	0/2420
13	M	0.27	0/1821	0.53	0/2470
13	a	0.26	0/1846	0.53	0/2503
14	N	0.25	0/1541	0.51	0/2087
14	b	0.25	0/1541	0.50	0/2087
All	All	0.27	0/50153	0.51	2/67817 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
11	K	4	LEU	CA-CB-CG	5.23	127.33	115.30
11	Y	4	LEU	CA-CB-CG	5.23	127.33	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1915	0	1929	2	0
1	O	1915	0	1929	10	0
2	B	1904	0	1904	11	0
2	P	1904	0	1904	11	0
3	C	1881	0	1895	6	0
3	Q	1881	0	1895	5	0
4	D	1813	0	1797	5	0
4	R	1813	0	1797	4	0
5	E	1773	0	1775	5	0
5	S	1773	0	1775	9	0
6	F	1892	0	1883	2	0
6	T	1892	0	1883	6	0
7	G	1907	0	1901	4	0
7	U	1907	0	1901	4	0
8	H	1685	0	1688	24	0
8	V	1685	0	1690	32	0
9	I	1581	0	1574	14	0
9	W	1581	0	1574	8	0
10	J	1561	0	1569	9	0
10	X	1561	0	1569	9	0
11	K	1644	0	1595	7	0
11	Y	1644	0	1595	6	0
12	L	1757	0	1711	13	0
12	Z	1757	0	1711	10	0
13	M	1790	0	1793	6	0
13	a	1815	0	1821	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
14	N	1512	0	1481	7	0
14	b	1512	0	1481	0	0
15	G	1	0	0	0	0
15	H	1	0	0	0	0
15	I	2	0	0	0	0
15	K	1	0	0	0	0
15	L	1	0	0	0	0
15	N	1	0	0	0	0
15	W	1	0	0	0	0
15	Z	1	0	0	0	0
16	G	1	0	0	0	0
16	U	1	0	0	0	0
17	A	15	0	0	0	0
17	B	15	0	0	1	0
17	C	16	0	0	0	0
17	D	15	0	0	0	0
17	E	13	0	0	0	0
17	F	15	0	0	0	0
17	G	21	0	0	0	0
17	H	23	0	0	0	0
17	I	21	0	0	0	0
17	J	20	0	0	0	0
17	K	25	0	0	0	0
17	L	29	0	0	0	0
17	M	37	0	0	0	0
17	N	20	0	0	1	0
17	O	9	0	0	0	0
17	P	12	0	0	0	0
17	Q	15	0	0	0	0
17	R	8	0	0	0	0
17	S	7	0	0	1	0
17	T	12	0	0	0	0
17	U	21	0	0	0	0
17	V	18	0	0	0	0
17	W	19	0	0	0	0
17	X	22	0	0	0	0
17	Y	21	0	0	0	0
17	Z	18	0	0	0	0
17	a	29	0	0	0	0
17	b	25	0	0	0	0
All	All	49787	0	49020	202	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 202 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
8:V:9:ASN:OD1	8:V:146:LEU:O	1.62	1.15
1:O:4:ARG:HH11	5:S:122:TYR:HD2	1.09	0.90
12:L:186:ASP:OD2	8:V:203:TYR:OH	1.88	0.90
1:O:4:ARG:NH1	5:S:122:TYR:HD2	1.68	0.89
8:H:9:ASN:H	8:H:9:ASN:ND2	1.71	0.87

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	248/250 (99%)	238 (96%)	9 (4%)	1 (0%)	34	66
1	O	248/250 (99%)	239 (96%)	9 (4%)	0	100	100
2	B	242/258 (94%)	235 (97%)	3 (1%)	4 (2%)	9	29
2	P	242/258 (94%)	235 (97%)	3 (1%)	4 (2%)	9	29
3	C	238/254 (94%)	232 (98%)	3 (1%)	3 (1%)	12	36
3	Q	238/254 (94%)	232 (98%)	3 (1%)	3 (1%)	12	36
4	D	231/260 (89%)	229 (99%)	2 (1%)	0	100	100
4	R	231/260 (89%)	229 (99%)	2 (1%)	0	100	100
5	E	229/234 (98%)	224 (98%)	5 (2%)	0	100	100
5	S	229/234 (98%)	223 (97%)	6 (3%)	0	100	100
6	F	241/288 (84%)	238 (99%)	3 (1%)	0	100	100
6	T	241/288 (84%)	238 (99%)	3 (1%)	0	100	100
7	G	239/252 (95%)	239 (100%)	0	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
7	U	239/252 (95%)	237 (99%)	2 (1%)	0	100	100
8	H	220/232 (95%)	215 (98%)	3 (1%)	2 (1%)	17	46
8	V	220/232 (95%)	214 (97%)	3 (1%)	3 (1%)	11	34
9	I	202/205 (98%)	194 (96%)	8 (4%)	0	100	100
9	W	202/205 (98%)	196 (97%)	6 (3%)	0	100	100
10	J	193/198 (98%)	188 (97%)	5 (3%)	0	100	100
10	X	193/198 (98%)	187 (97%)	6 (3%)	0	100	100
11	K	210/212 (99%)	206 (98%)	4 (2%)	0	100	100
11	Y	210/212 (99%)	206 (98%)	4 (2%)	0	100	100
12	L	220/222 (99%)	216 (98%)	4 (2%)	0	100	100
12	Z	220/222 (99%)	216 (98%)	4 (2%)	0	100	100
13	M	227/246 (92%)	218 (96%)	9 (4%)	0	100	100
13	a	230/246 (94%)	221 (96%)	9 (4%)	0	100	100
14	N	194/196 (99%)	190 (98%)	4 (2%)	0	100	100
14	b	194/196 (99%)	190 (98%)	4 (2%)	0	100	100
All	All	6271/6614 (95%)	6125 (98%)	126 (2%)	20 (0%)	41	72

5 of 20 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	51	VAL
2	B	220	ASN
2	B	222	GLY
3	C	202	GLN
8	H	195	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	209/209 (100%)	205 (98%)	4 (2%)	57	85

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	O	209/209 (100%)	204 (98%)	5 (2%)	49	81
2	B	203/216 (94%)	195 (96%)	8 (4%)	32	66
2	P	203/216 (94%)	195 (96%)	8 (4%)	32	66
3	C	212/226 (94%)	201 (95%)	11 (5%)	23	55
3	Q	212/226 (94%)	201 (95%)	11 (5%)	23	55
4	D	194/215 (90%)	183 (94%)	11 (6%)	20	50
4	R	194/215 (90%)	183 (94%)	11 (6%)	20	50
5	E	190/193 (98%)	180 (95%)	10 (5%)	22	54
5	S	190/193 (98%)	180 (95%)	10 (5%)	22	54
6	F	201/239 (84%)	191 (95%)	10 (5%)	24	56
6	T	201/239 (84%)	191 (95%)	10 (5%)	24	56
7	G	206/210 (98%)	196 (95%)	10 (5%)	25	57
7	U	206/210 (98%)	196 (95%)	10 (5%)	25	57
8	H	181/190 (95%)	169 (93%)	12 (7%)	16	44
8	V	181/190 (95%)	167 (92%)	14 (8%)	13	35
9	I	172/173 (99%)	167 (97%)	5 (3%)	42	76
9	W	172/173 (99%)	169 (98%)	3 (2%)	60	87
10	J	173/175 (99%)	167 (96%)	6 (4%)	36	70
10	X	173/175 (99%)	167 (96%)	6 (4%)	36	70
11	K	169/169 (100%)	161 (95%)	8 (5%)	26	59
11	Y	169/169 (100%)	161 (95%)	8 (5%)	26	59
12	L	185/185 (100%)	179 (97%)	6 (3%)	39	73
12	Z	185/185 (100%)	179 (97%)	6 (3%)	39	73
13	M	195/208 (94%)	189 (97%)	6 (3%)	40	74
13	a	198/208 (95%)	192 (97%)	6 (3%)	41	75
14	N	162/162 (100%)	157 (97%)	5 (3%)	40	74
14	b	162/162 (100%)	157 (97%)	5 (3%)	40	74
All	All	5307/5540 (96%)	5082 (96%)	225 (4%)	30	63

5 of 225 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	O	157	PHE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
14	b	36	ARG
4	R	214	ILE
13	a	187	ARG
10	X	110	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 120 such sidechains are listed below:

Mol	Chain	Res	Type
13	M	213	GLN
12	Z	55	ASN
4	R	15	GLN
12	Z	49	ASN
13	a	213	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 11 ligands modelled in this entry, 11 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	250/250 (100%)	-0.44	5 (2%) 65 56	39, 52, 87, 126	0
1	O	250/250 (100%)	-0.41	4 (1%) 72 66	25, 60, 104, 132	0
2	B	244/258 (94%)	-0.38	7 (2%) 51 41	38, 58, 99, 150	0
2	P	244/258 (94%)	-0.37	11 (4%) 33 23	44, 60, 101, 147	0
3	C	240/254 (94%)	-0.30	11 (4%) 32 22	39, 62, 121, 148	0
3	Q	240/254 (94%)	-0.07	11 (4%) 32 22	48, 74, 154, 182	0
4	D	235/260 (90%)	-0.47	2 (0%) 84 80	42, 62, 93, 134	0
4	R	235/260 (90%)	-0.41	3 (1%) 77 72	47, 67, 106, 141	0
5	E	231/234 (98%)	-0.37	2 (0%) 84 80	47, 66, 103, 146	0
5	S	231/234 (98%)	-0.25	3 (1%) 77 72	51, 75, 113, 154	0
6	F	243/288 (84%)	-0.49	5 (2%) 63 54	43, 60, 110, 137	0
6	T	243/288 (84%)	-0.38	3 (1%) 79 73	45, 70, 122, 155	0
7	G	241/252 (95%)	-0.57	1 (0%) 92 91	34, 54, 88, 137	0
7	U	241/252 (95%)	-0.47	5 (2%) 63 54	30, 58, 87, 133	0
8	H	222/232 (95%)	-0.51	6 (2%) 54 44	32, 50, 83, 150	0
8	V	222/232 (95%)	-0.50	7 (3%) 47 37	24, 53, 81, 139	0
9	I	204/205 (99%)	-0.77	1 (0%) 91 88	37, 49, 73, 96	0
9	W	204/205 (99%)	-0.73	2 (0%) 82 77	37, 48, 78, 102	0
10	J	195/198 (98%)	-0.62	3 (1%) 73 68	36, 51, 76, 130	0
10	X	195/198 (98%)	-0.56	3 (1%) 73 68	39, 53, 77, 137	0
11	K	212/212 (100%)	-0.66	1 (0%) 91 88	32, 48, 70, 90	0
11	Y	212/212 (100%)	-0.65	1 (0%) 91 88	41, 52, 75, 96	0
12	L	222/222 (100%)	-0.62	1 (0%) 91 88	38, 53, 82, 117	0
12	Z	222/222 (100%)	-0.64	1 (0%) 91 88	39, 53, 84, 119	0

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
13	M	229/246 (93%)	-0.68	1 (0%) 92 91	35, 53, 76, 92	0
13	a	232/246 (94%)	-0.66	1 (0%) 92 91	36, 53, 74, 92	0
14	N	196/196 (100%)	-0.69	2 (1%) 82 77	37, 48, 74, 102	0
14	b	196/196 (100%)	-0.67	2 (1%) 82 77	38, 49, 77, 110	0
All	All	6331/6614 (95%)	-0.50	105 (1%) 70 63	24, 56, 100, 182	0

The worst 5 of 105 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	Q	50	LEU	6.5
2	P	220	ASN	6.0
3	Q	49	THR	5.8
5	S	202	ASP	5.8
2	B	219	ALA	5.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
15	MG	H	301	1/1	0.91	0.28	30,30,30,30	0
15	MG	Z	301	1/1	0.92	0.24	59,59,59,59	0
15	MG	G	301	1/1	0.94	0.07	51,51,51,51	0
15	MG	W	301	1/1	0.95	0.27	62,62,62,62	0
15	MG	K	301	1/1	0.97	0.05	46,46,46,46	0
15	MG	L	301	1/1	0.98	0.06	61,61,61,61	0
15	MG	N	201	1/1	0.98	0.07	48,48,48,48	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
15	MG	I	302	1/1	0.98	0.05	52,52,52,52	0
15	MG	I	301	1/1	0.98	0.24	50,50,50,50	0
16	CL	U	301	1/1	0.98	0.14	41,41,41,41	0
16	CL	G	302	1/1	1.00	0.12	38,38,38,38	0

6.5 Other polymers [i](#)

There are no such residues in this entry.