



wwPDB X-ray Structure Validation Summary Report ⓘ

Dec 20, 2023 – 12:17 pm GMT

PDB ID : 8C0D
Title : UFL1/DDRGK1 bound to UFC1
Authors : Magnussen, H.M.; Kulathu, Y.
Deposited on : 2022-12-16
Resolution : 2.56 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.56 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1279 (2.58-2.54)
Clashscore	141614	1327 (2.58-2.54)
Ramachandran outliers	138981	1312 (2.58-2.54)
Sidechain outliers	138945	1312 (2.58-2.54)
RSRZ outliers	127900	1269 (2.58-2.54)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	194	 2% (poor fit), 70% (0-1 outliers), 13% (2 outliers), 15% (3+ outliers)
1	D	194	 5% (poor fit), 70% (0-1 outliers), 12% (2 outliers), 17% (3+ outliers)
2	B	110	 % (poor fit), 70% (0-1 outliers), 22% (2 outliers), 7% (3+ outliers)
2	E	110	 5% (poor fit), 63% (0-1 outliers), 20% (2 outliers), 15% (3+ outliers)
3	C	168	 2% (poor fit), 80% (0-1 outliers), 14% (2 outliers), % (3+ outliers)

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	F	168	 2% 77% 19% ...

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 12907 atoms, of which 6389 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called E3 UFM1-protein ligase 1.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	165	2405	763	1189	217	232	4	52	0	0
1	D	161	2394	759	1184	214	233	4	45	0	0

There are 30 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-14	MET	-	initiating methionine	UNP O94874
A	-13	GLY	-	expression tag	UNP O94874
A	-12	HIS	-	expression tag	UNP O94874
A	-11	HIS	-	expression tag	UNP O94874
A	-10	HIS	-	expression tag	UNP O94874
A	-9	HIS	-	expression tag	UNP O94874
A	-8	HIS	-	expression tag	UNP O94874
A	-7	HIS	-	expression tag	UNP O94874
A	-6	GLU	-	expression tag	UNP O94874
A	-5	ASN	-	expression tag	UNP O94874
A	-4	LEU	-	expression tag	UNP O94874
A	-3	TYR	-	expression tag	UNP O94874
A	-2	PHE	-	expression tag	UNP O94874
A	-1	GLN	-	expression tag	UNP O94874
A	0	GLY	-	expression tag	UNP O94874
D	-14	MET	-	initiating methionine	UNP O94874
D	-13	GLY	-	expression tag	UNP O94874
D	-12	HIS	-	expression tag	UNP O94874
D	-11	HIS	-	expression tag	UNP O94874
D	-10	HIS	-	expression tag	UNP O94874
D	-9	HIS	-	expression tag	UNP O94874
D	-8	HIS	-	expression tag	UNP O94874
D	-7	HIS	-	expression tag	UNP O94874
D	-6	GLU	-	expression tag	UNP O94874
D	-5	ASN	-	expression tag	UNP O94874

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	-4	LEU	-	expression tag	UNP O94874
D	-3	TYR	-	expression tag	UNP O94874
D	-2	PHE	-	expression tag	UNP O94874
D	-1	GLN	-	expression tag	UNP O94874
D	0	GLY	-	expression tag	UNP O94874

- Molecule 2 is a protein called DDRGK domain-containing protein 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
2	B	102	Total	C	H	N	O	S	22	0	0
			1562	491	782	132	156	1			
2	E	94	Total	C	H	N	O		17	0	0
			1449	458	729	124	138				

- Molecule 3 is a protein called Ubiquitin-fold modifier-conjugating enzyme 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
3	C	164	Total	C	H	N	O	S	41	0	0
			2532	834	1245	217	232	4			
3	F	164	Total	C	H	N	O	S	38	0	0
			2558	839	1260	220	235	4			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	116	LYS	CYS	engineered mutation	UNP Q9Y3C8
C	168	GLY	-	expression tag	UNP Q9Y3C8
F	116	LYS	CYS	engineered mutation	UNP Q9Y3C8
F	168	GLY	-	expression tag	UNP Q9Y3C8

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	O	0	0
			1	1		
4	C	2	Total	O	0	0
			2	2		
4	D	1	Total	O	0	0
			1	1		
4	E	1	Total	O	0	0
			1	1		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	F	2	Total	O	0	0
			2	2		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: E3 UFM1-protein ligase 1




- Molecule 1: E3 UFM1-protein ligase 1




- Molecule 2: DDRGK domain-containing protein 1



- Molecule 2: DDRGK domain-containing protein 1





- Molecule 3: Ubiquitin-fold modifier-conjugating enzyme 1

Chain C: 



- Molecule 3: Ubiquitin-fold modifier-conjugating enzyme 1

Chain F: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	79.21Å 124.56Å 131.29Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	65.73 – 2.56 65.65 – 2.56	Depositor EDS
% Data completeness (in resolution range)	61.0 (65.73-2.56) 61.0 (65.65-2.56)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.64 (at 2.58Å)	Xtrriage
Refinement program	REFMAC 5.8.0403	Depositor
R, R_{free}	0.232 , 0.286 0.233 , 0.285	Depositor DCC
R_{free} test set	1222 reflections (4.73%)	wwPDB-VP
Wilson B-factor (Å ²)	46.6	Xtrriage
Anisotropy	0.215	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 46.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	12907	wwPDB-VP
Average B, all atoms (Å ²)	51.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 35.38 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 5.8841e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.53	0/1228	0.94	3/1669 (0.2%)
1	D	0.52	0/1223	0.90	1/1660 (0.1%)
2	B	0.68	3/788 (0.4%)	1.00	1/1068 (0.1%)
2	E	0.66	1/728 (0.1%)	1.08	3/988 (0.3%)
3	C	0.58	0/1324	1.02	7/1806 (0.4%)
3	F	0.62	1/1335 (0.1%)	0.99	5/1818 (0.3%)
All	All	0.59	5/6626 (0.1%)	0.98	20/9009 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	4
1	D	0	3
2	B	0	2
2	E	0	1
3	C	0	1
3	F	0	1
All	All	0	12

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	274	GLU	CD-OE2	-6.54	1.18	1.25
3	F	11	SER	CA-CB	-5.95	1.44	1.52
2	E	274	GLU	CD-OE1	-5.92	1.19	1.25
2	B	274	GLU	CD-OE1	-5.53	1.19	1.25
2	B	218	GLU	CD-OE2	5.36	1.31	1.25

The worst 5 of 20 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	9	ARG	NE-CZ-NH2	-9.36	115.62	120.30
1	A	9	ARG	NE-CZ-NH1	8.55	124.57	120.30
3	C	64	ARG	CG-CD-NE	7.44	127.42	111.80
3	C	127	ARG	NE-CZ-NH1	-6.49	117.06	120.30
3	F	64	ARG	CG-CD-NE	6.39	125.21	111.80

There are no chirality outliers.

5 of 12 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	26	ARG	Sidechain
1	A	73	ARG	Sidechain
1	A	9	ARG	Sidechain
1	A	96	ARG	Sidechain
2	B	265	ARG	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1216	1189	1123	16	0
1	D	1210	1184	1135	15	0
2	B	780	782	769	14	0
2	E	720	729	719	12	0
3	C	1287	1245	1208	13	0
3	F	1298	1260	1233	22	0
4	A	1	0	0	0	0
4	C	2	0	0	0	0
4	D	1	0	0	0	0
4	E	1	0	0	0	0
4	F	2	0	0	0	0
All	All	6518	6389	6187	78	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 78 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:70:LEU:HD13	1:A:77:VAL:HG12	1.50	0.90
3:C:166:ASN:CB	3:F:164:LYS:HA	2.02	0.90
1:D:70:LEU:HD12	1:D:77:VAL:HG13	1.66	0.78
3:C:122:LYS:HB2	3:C:123:PRO:HD3	1.79	0.64
2:B:239:GLY:HA3	3:F:166:ASN:HA	1.80	0.62

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	161/194 (83%)	146 (91%)	13 (8%)	2 (1%)	13	17
1	D	157/194 (81%)	145 (92%)	12 (8%)	0	100	100
2	B	100/110 (91%)	98 (98%)	2 (2%)	0	100	100
2	E	92/110 (84%)	90 (98%)	1 (1%)	1 (1%)	14	19
3	C	162/168 (96%)	151 (93%)	8 (5%)	3 (2%)	8	9
3	F	162/168 (96%)	155 (96%)	6 (4%)	1 (1%)	25	33
All	All	834/944 (88%)	785 (94%)	42 (5%)	7 (1%)	19	27

5 of 7 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	133	GLU
3	C	164	LYS
1	A	161	LEU
3	C	6	THR
2	E	214	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	112/171 (66%)	108 (96%)	4 (4%)	35	47
1	D	117/171 (68%)	114 (97%)	3 (3%)	46	59
2	B	82/92 (89%)	80 (98%)	2 (2%)	49	63
2	E	74/92 (80%)	69 (93%)	5 (7%)	16	20
3	C	126/146 (86%)	122 (97%)	4 (3%)	39	51
3	F	130/146 (89%)	128 (98%)	2 (2%)	65	77
All	All	641/818 (78%)	621 (97%)	20 (3%)	40	52

5 of 20 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	E	276	LEU
2	E	294	LEU
3	F	167	GLN
3	F	17	LYS
3	C	17	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
3	C	161	HIS
1	D	44	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	165/194 (85%)	0.26	3 (1%) 68 75	31, 55, 111, 143	0
1	D	161/194 (82%)	0.39	9 (5%) 24 31	33, 52, 118, 139	0
2	B	102/110 (92%)	0.27	1 (0%) 82 87	32, 45, 68, 75	0
2	E	94/110 (85%)	0.48	6 (6%) 19 24	31, 51, 73, 92	0
3	C	164/168 (97%)	0.32	4 (2%) 59 67	25, 44, 87, 106	0
3	F	164/168 (97%)	0.25	3 (1%) 68 75	28, 43, 77, 96	0
All	All	850/944 (90%)	0.32	26 (3%) 49 58	25, 48, 99, 143	0

The worst 5 of 26 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	E	240	LEU	3.5
3	C	107	ALA	3.0
1	D	103	SER	2.9
3	C	166	ASN	2.9
1	A	131	LEU	2.9

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.