



wwPDB X-ray Structure Validation Summary Report ⓘ

Nov 7, 2023 – 03:39 AM EST

PDB ID : 3C1P
Title : Crystal Structure of an alternating D-Alanyl, L-Homoalanyl PNA
Authors : Cuesta-Seijo, J.A.; Sheldrick, G.M.; Zhang, J.; Diederichsen, U.
Deposited on : 2008-01-23
Resolution : 1.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



| Metric | Whole archive (#Entries) | Similar resolution (#Entries, resolution range(Å)) |
|------------|-----------------------------|---|
| Clashscore | 141614 | 1117 (1.06-0.94) |

ENTRY-COMPOSITION INFOmissingINFO

2 Residue-property plots

There is no protein, DNA or RNA chain in this entry to show sequence plots.

3 Data and refinement statistics

| Property | Value | Source |
|---|---|------------------|
| Space group | P 1 21 1 | Depositor |
| Cell constants a, b, c, α , β , γ | 26.31Å 30.73Å 33.55Å 90.00° 99.53° 90.00° | Depositor |
| Resolution (Å) | 40.00 – 1.00 33.09 – 1.00 | Depositor EDS |
| % Data completeness (in resolution range) | 99.1 (40.00-1.00) 99.5 (33.09-1.00) | Depositor EDS |
| R_{merge} | 0.06 | Depositor |
| R_{sym} | 0.06 | Depositor |
| $\langle I/\sigma(I) \rangle$ ¹ | 1.20 (at 1.00Å) | Xtrriage |
| Refinement program | SHELXL-97 | Depositor |
| R, R_{free} | 0.150 , 0.202 0.160 , (Not available) | Depositor DCC |
| R_{free} test set | No test flags present. | wwPDB-VP |
| Wilson B-factor (Å ²) | 12.1 | Xtrriage |
| Anisotropy | 0.472 | Xtrriage |
| Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²) | 0.27 , 55.4 | EDS |
| L-test for twinning ² | $\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$ | Xtrriage |
| Estimated twinning fraction | No twinning to report. | Xtrriage |
| F_o, F_c correlation | 0.98 | EDS |
| Total number of atoms | 814 | wwPDB-VP |
| Average B, all atoms (Å ²) | 22.0 | wwPDB-VP |

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 18.93% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

4 Model quality [i](#)

4.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: AGD, DLY, LHC, HGL, CUD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Chain | Bond lengths | | Bond angles | |
|-----|-------|--------------|---------|-------------|---------|
| | | RMSZ | # Z >5 | RMSZ | # Z >5 |
| 1 | D | 0.33 | 0/2 | 0.00 | - |

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

4.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

| Mol | Chain | Non-H | H(model) | H(added) | Clashes | Symm-Clashes |
|-----|-------|-------|----------|----------|---------|--------------|
| 1 | A | 102 | 70 | 65 | 0 | 0 |
| 1 | B | 100 | 66 | 64 | 0 | 0 |
| 1 | C | 100 | 66 | 65 | 0 | 0 |
| 1 | D | 109 | 61 | 56 | 0 | 0 |
| 2 | A | 44 | 0 | 0 | 0 | 0 |
| 2 | B | 29 | 0 | 0 | 0 | 0 |
| 2 | C | 34 | 0 | 0 | 0 | 0 |
| 2 | D | 33 | 0 | 0 | 0 | 0 |
| All | All | 551 | 263 | 250 | 0 | 0 |

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 0.

There are no clashes within the asymmetric unit.

There are no symmetry-related clashes.

4.3 Torsion angles [i](#)

4.3.1 Protein backbone [i](#)

There are no protein molecules in this entry.

4.3.2 Protein sidechains [i](#)

There are no protein molecules in this entry.

4.3.3 RNA [i](#)

There are no RNA molecules in this entry.

4.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

4.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

4.6 Ligand geometry [i](#)

There are no ligands in this entry.

4.7 Other polymers [i](#)

Of 37 such residues modelled in this entry, 3 are modelled with single atom - leaving 34 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Type | Chain | Res | Link | Bond lengths | | | Bond angles | | |
|-----|------|-------|-------|------|--------------|------|----------|-------------|------|----------|
| | | | | | Counts | RMSZ | # Z > 2 | Counts | RMSZ | # Z > 2 |
| 1 | AGD | A | 13 | 1 | 12,17,18 | 1.77 | 2 (16%) | 9,24,26 | 1.21 | 1 (11%) |
| 1 | CUD | C | 37 | 1 | 12,13,14 | 0.83 | 0 | 11,17,19 | 1.60 | 3 (27%) |
| 1 | LHC | B | 26 | 1 | 13,14,15 | 0.91 | 0 | 12,18,20 | 1.24 | 1 (8%) |
| 1 | AGD | C | 33 | 1 | 12,17,18 | 1.59 | 3 (25%) | 9,24,26 | 1.22 | 1 (11%) |
| 1 | AGD | C | 35 | 1 | 12,17,18 | 1.76 | 3 (25%) | 9,24,26 | 1.44 | 1 (11%) |
| 1 | CUD | D | 47[A] | 1 | 12,13,14 | 1.01 | 1 (8%) | 11,17,19 | 1.10 | 1 (9%) |
| 1 | HGL | A | 12 | 1 | 13,18,19 | 1.54 | 3 (23%) | 10,25,27 | 1.16 | 1 (10%) |
| 1 | AGD | B | 25 | 1 | 12,17,18 | 1.61 | 3 (25%) | 9,24,26 | 1.70 | 2 (22%) |
| 1 | LHC | A | 14 | 1 | 13,14,15 | 0.78 | 0 | 12,18,20 | 0.60 | 0 |
| 1 | HGL | C | 32 | 1 | 13,18,19 | 1.63 | 3 (23%) | 10,25,27 | 1.05 | 0 |
| 1 | LHC | C | 34 | 1 | 13,14,15 | 1.25 | 1 (7%) | 12,18,20 | 1.19 | 2 (16%) |
| 1 | LHC | D | 44 | 1 | 13,14,15 | 0.74 | 0 | 12,18,20 | 0.86 | 1 (8%) |
| 1 | LHC | B | 24 | 1 | 13,14,15 | 1.05 | 1 (7%) | 12,18,20 | 0.47 | 0 |
| 1 | CUD | A | 17[B] | - | 12,12,14 | 3.69 | 3 (25%) | 11,16,19 | 0.87 | 0 |
| 1 | LHC | C | 36 | 1 | 13,14,15 | 0.93 | 0 | 12,18,20 | 0.81 | 0 |
| 1 | AGD | B | 23 | 1 | 12,17,18 | 1.89 | 3 (25%) | 9,24,26 | 1.03 | 1 (11%) |
| 1 | LHC | D | 46 | 1 | 13,14,15 | 1.17 | 1 (7%) | 12,18,20 | 1.16 | 1 (8%) |
| 1 | HGL | D | 42[B] | 1 | 13,18,19 | 1.63 | 3 (23%) | 10,25,27 | 0.82 | 1 (10%) |
| 1 | AGD | D | 43 | 1 | 12,17,18 | 1.58 | 4 (33%) | 9,24,26 | 1.15 | 1 (11%) |
| 1 | CUD | D | 47[B] | 1 | 12,13,14 | 0.90 | 0 | 11,17,19 | 1.10 | 1 (9%) |
| 1 | CUD | A | 17[A] | 1 | 12,13,14 | 1.18 | 1 (8%) | 11,17,19 | 0.87 | 0 |
| 1 | LHC | A | 16 | 1 | 13,14,15 | 1.15 | 1 (7%) | 12,18,20 | 1.71 | 3 (25%) |
| 1 | AGD | D | 45 | 1 | 12,17,18 | 1.91 | 3 (25%) | 9,24,26 | 0.73 | 0 |
| 1 | HGL | B | 22 | 1 | 13,18,19 | 2.21 | 5 (38%) | 10,25,27 | 1.55 | 1 (10%) |
| 1 | HGL | D | 42[A] | 1 | 13,18,19 | 1.43 | 2 (15%) | 10,25,27 | 0.82 | 1 (10%) |
| 1 | AGD | A | 15 | 1 | 12,17,18 | 1.68 | 2 (16%) | 9,24,26 | 1.50 | 2 (22%) |
| 1 | CUD | B | 27 | 1 | 12,13,14 | 0.41 | 0 | 11,17,19 | 1.68 | 2 (18%) |

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

| Mol | Type | Chain | Res | Link | Chirals | Torsions | Rings |
|-----|------|-------|-----|------|---------|----------|---------|
| 1 | AGD | A | 13 | 1 | - | 0/1/6/8 | 0/2/2/2 |
| 1 | CUD | C | 37 | 1 | - | 0/3/6/8 | 0/1/1/1 |
| 1 | LHC | B | 26 | 1 | - | 0/6/7/9 | 0/1/1/1 |

Continued on next page...

Continued from previous page...

| Mol | Type | Chain | Res | Link | Chirals | Torsions | Rings |
|-----|------|-------|-------|------|---------|----------|---------|
| 1 | AGD | C | 33 | 1 | - | 0/1/6/8 | 0/2/2/2 |
| 1 | AGD | C | 35 | 1 | - | 0/1/6/8 | 0/2/2/2 |
| 1 | CUD | D | 47[A] | 1 | - | 0/3/6/8 | 0/1/1/1 |
| 1 | HGL | A | 12 | 1 | - | 0/6/7/9 | 0/2/2/2 |
| 1 | AGD | B | 25 | 1 | - | 0/1/6/8 | 0/2/2/2 |
| 1 | LHC | A | 14 | 1 | - | 0/6/7/9 | 0/1/1/1 |
| 1 | HGL | C | 32 | 1 | - | 0/6/7/9 | 0/2/2/2 |
| 1 | LHC | C | 34 | 1 | - | 0/6/7/9 | 0/1/1/1 |
| 1 | LHC | D | 44 | 1 | - | 0/6/7/9 | 0/1/1/1 |
| 1 | LHC | B | 24 | 1 | - | 0/6/7/9 | 0/1/1/1 |
| 1 | CUD | A | 17[B] | - | - | 1/3/4/8 | 0/1/1/1 |
| 1 | LHC | C | 36 | 1 | - | 0/6/7/9 | 0/1/1/1 |
| 1 | AGD | B | 23 | 1 | - | 0/1/6/8 | 0/2/2/2 |
| 1 | LHC | D | 46 | 1 | - | 0/6/7/9 | 0/1/1/1 |
| 1 | HGL | D | 42[B] | 1 | - | 0/6/7/9 | 0/2/2/2 |
| 1 | AGD | D | 43 | 1 | - | 0/1/6/8 | 0/2/2/2 |
| 1 | CUD | D | 47[B] | 1 | - | 0/3/6/8 | 0/1/1/1 |
| 1 | CUD | A | 17[A] | 1 | - | 0/3/6/8 | 0/1/1/1 |
| 1 | LHC | A | 16 | 1 | - | 0/6/7/9 | 0/1/1/1 |
| 1 | AGD | D | 45 | 1 | - | 0/1/6/8 | 0/2/2/2 |
| 1 | HGL | B | 22 | 1 | - | 0/6/7/9 | 0/2/2/2 |
| 1 | HGL | D | 42[A] | 1 | - | 0/6/7/9 | 0/2/2/2 |
| 1 | AGD | A | 15 | 1 | - | 0/1/6/8 | 0/2/2/2 |
| 1 | CUD | B | 27 | 1 | - | 1/3/6/8 | 0/1/1/1 |

The worst 5 of 48 bond length outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed(Å) | Ideal(Å) |
|-----|-------|-------|------|-------|-------|-------------|----------|
| 1 | A | 17[B] | CUD | O-C | 12.02 | 1.67 | 1.19 |
| 1 | A | 13 | AGD | C5-C6 | -4.18 | 1.38 | 1.47 |
| 1 | D | 45 | AGD | C8-N7 | -4.06 | 1.28 | 1.35 |
| 1 | B | 22 | HGL | C6-N1 | 3.87 | 1.43 | 1.37 |
| 1 | B | 22 | HGL | C8-N7 | -3.87 | 1.28 | 1.35 |

The worst 5 of 28 bond angle outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed(°) | Ideal(°) |
|-----|-------|-----|------|----------|-------|-------------|----------|
| 1 | B | 27 | CUD | C6-N1-C2 | 3.82 | 122.80 | 120.23 |
| 1 | A | 16 | LHC | C6-N1-C2 | 3.78 | 122.77 | 120.23 |
| 1 | C | 37 | CUD | N1-C2-N3 | 3.48 | 121.97 | 118.45 |
| 1 | B | 25 | AGD | O6-C6-N1 | 3.47 | 124.74 | 120.65 |
| 1 | B | 26 | LHC | C6-N1-C2 | -3.02 | 118.19 | 120.23 |

There are no chirality outliers.

All (2) torsion outliers are listed below:

| Mol | Chain | Res | Type | Atoms |
|-----|-------|-------|------|-------------|
| 1 | B | 27 | CUD | CA-CB-N1-C2 |
| 1 | A | 17[B] | CUD | C-CA-CB-N1 |

There are no ring outliers.

No monomer is involved in short contacts.

4.8 Polymer linkage issues

There are no chain breaks in this entry.

5 Fit of model and data [i](#)

5.1 Protein, DNA and RNA chains [i](#)

There are no RSRZ outliers to report within protein, DNA, RNA chains in this entry.

5.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.4 Ligands [i](#)

There are no ligands in this entry.

5.5 Other polymers [i](#)

There are no such residues in this entry.