



# Full wwPDB X-ray Structure Validation Report ⓘ

Nov 19, 2023 – 10:33 PM JST

PDB ID : 7C1E  
Title : Crystal structure of Kluyveromyces polyspora ADH (KpADH) mutant (Y127W)  
Authors : Wu, Y.F.; Zhou, J.Y.; Liu, Y.F.; Xu, G.C.; Ni, Y.  
Deposited on : 2020-05-03  
Resolution : 1.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2340 (1.76-1.76)
Clashscore	141614	2466 (1.76-1.76)
Ramachandran outliers	138981	2437 (1.76-1.76)
Sidechain outliers	138945	2437 (1.76-1.76)
RSRZ outliers	127900	2298 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	342	
1	B	342	

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 11872 atoms, of which 5493 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Epimerase domain-containing protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	342	5463	1760	2716	457	522	8	0	0	0
1	B	342	5474	1760	2727	457	522	8	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	127	TRP	TYR	engineered mutation	UNP A7TM80
B	127	TRP	TYR	engineered mutation	UNP A7TM80

- Molecule 2 is NADP NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NAP) (formula:  $C_{21}H_{28}N_7O_{17}P_3$ ) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
			Total	C	H	N	O	P		
2	A	1	73	21	25	7	17	3	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
			Total	C	H	N	O			P
2	B	1	73	21	25	7	17	3	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	368	Total	O	0	0
			368	368		
3	B	421	Total	O	0	0
			421	421		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Epimerase domain-containing protein



- Molecule 1: Epimerase domain-containing protein



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 61	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	102.69Å 102.69Å 134.41Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	25.67 – 1.75 25.67 – 1.75	Depositor EDS
% Data completeness (in resolution range)	90.6 (25.67-1.75) 90.6 (25.67-1.75)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.99 (at 1.75Å)	Xtrriage
Refinement program	PHENIX 1.17.1_3660	Depositor
R, $R_{free}$	0.153 , 0.183 0.161 , 0.187	Depositor DCC
$R_{free}$ test set	3628 reflections (4.97%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	17.0	Xtrriage
Anisotropy	0.098	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.42 , 47.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.032 for h,-h-k,-l	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	11872	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	22.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 71.36 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.6652e-06. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.50	0/2809	0.63	1/3804 (0.0%)
1	B	0.53	2/2809 (0.1%)	0.64	1/3804 (0.0%)
All	All	0.51	2/5618 (0.0%)	0.64	2/7608 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	112	LYS	CD-CE	-7.72	1.31	1.51
1	B	112	LYS	CE-NZ	7.59	1.68	1.49

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	35	ASP	N-CA-CB	6.50	122.30	110.60
1	B	112	LYS	CD-CE-NZ	-5.13	99.90	111.70

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2747	2716	2725	10	0
1	B	2747	2727	2725	18	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	48	25	25	2	0
2	B	48	25	25	4	0
3	A	368	0	0	2	0
3	B	421	0	0	5	0
All	All	6379	5493	5500	28	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (28) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:112:LYS:CE	1:B:112:LYS:NZ	1.68	1.55
1:B:112:LYS:NZ	1:B:112:LYS:CD	2.30	0.92
1:A:287:LYS:HE3	3:A:516:HOH:O	1.88	0.73
1:B:183:SER:HA	3:B:635:HOH:O	1.94	0.67
1:B:215:THR:HG21	2:B:401:NAP:N7N	2.10	0.66
1:A:210:LYS:HE3	3:A:658:HOH:O	1.95	0.66
1:B:215:THR:HG21	2:B:401:NAP:H72N	1.61	0.65
1:A:114:ALA:HB1	1:A:117:THR:HB	1.80	0.64
1:B:127:TRP:CD1	1:B:196:SER:HA	2.32	0.64
1:B:39:LYS:HD3	3:B:542:HOH:O	2.01	0.61
1:A:127:TRP:CD1	1:A:196:SER:HA	2.36	0.60
1:B:119:GLU:HG2	1:B:187:LYS:HE3	1.84	0.58
1:B:233:LYS:HD3	1:B:296:ASP:O	2.06	0.55
1:B:39:LYS:CD	3:B:542:HOH:O	2.60	0.49
1:B:189:LYS:HE2	3:B:679:HOH:O	2.13	0.49
1:A:215:THR:HG21	2:A:401:NAP:O7N	2.14	0.47
1:B:112:LYS:NZ	1:B:112:LYS:HD2	2.25	0.47
1:B:198:VAL:H	2:B:401:NAP:H72N	1.62	0.47
1:A:233:LYS:HD3	1:A:296:ASP:O	2.16	0.46
1:B:43:GLN:HE21	1:B:206:GLU:CD	2.19	0.46
1:B:255:LYS:NZ	3:B:513:HOH:O	2.49	0.45
1:B:125:ALA:O	2:B:401:NAP:H6N	2.18	0.42
1:B:189:LYS:HB3	1:B:259:ILE:HG23	2.02	0.41
1:B:14:LYS:NZ	1:B:207:ASP:OD2	2.37	0.41
1:A:67:PHE:HB3	1:A:113:TYR:HB2	2.02	0.41
1:A:125:ALA:O	2:A:401:NAP:H6N	2.21	0.40
1:A:43:GLN:HE21	1:A:206:GLU:CD	2.25	0.40
1:A:39:LYS:HA	1:A:39:LYS:HD3	1.92	0.40



There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	340/342 (99%)	330 (97%)	10 (3%)	0	100	100
1	B	340/342 (99%)	331 (97%)	9 (3%)	0	100	100
All	All	680/684 (99%)	661 (97%)	19 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	305/305 (100%)	298 (98%)	7 (2%)	50	28
1	B	305/305 (100%)	299 (98%)	6 (2%)	55	34
All	All	610/610 (100%)	597 (98%)	13 (2%)	53	31

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	35	ASP
1	A	42	LYS
1	A	127	TRP
1	A	131	MET
1	A	180	LYS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	187	LYS
1	A	239	PHE
1	B	42	LYS
1	B	127	TRP
1	B	131	MET
1	B	154	GLU
1	B	184	ASP
1	B	239	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	NAP	A	401	-	45,52,52	4.14	17 (37%)	56,80,80	2.06	5 (8%)
2	NAP	B	401	-	45,52,52	4.09	18 (40%)	56,80,80	2.30	12 (21%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAP	A	401	-	-	7/31/67/67	0/5/5/5
2	NAP	B	401	-	-	3/31/67/67	0/5/5/5

All (35) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401	NAP	C2D-C1D	-16.03	1.29	1.53
2	B	401	NAP	C2D-C1D	-15.96	1.29	1.53
2	B	401	NAP	C3D-C4D	-9.72	1.28	1.53
2	A	401	NAP	O4D-C1D	9.59	1.54	1.41
2	A	401	NAP	C3D-C4D	-9.35	1.29	1.53
2	B	401	NAP	C3B-C4B	-8.93	1.30	1.53
2	B	401	NAP	O4D-C1D	8.89	1.53	1.41
2	A	401	NAP	C3B-C4B	-8.65	1.30	1.53
2	A	401	NAP	O4B-C4B	7.81	1.62	1.45
2	B	401	NAP	O4B-C4B	7.17	1.61	1.45
2	B	401	NAP	C2D-C3D	5.26	1.67	1.53
2	A	401	NAP	O4B-C1B	-5.17	1.33	1.41
2	A	401	NAP	C7N-N7N	5.08	1.42	1.33
2	A	401	NAP	C2D-C3D	4.94	1.66	1.53
2	B	401	NAP	O4B-C1B	-4.66	1.34	1.41
2	B	401	NAP	C2A-N3A	4.48	1.39	1.32
2	B	401	NAP	O4D-C4D	4.42	1.54	1.45
2	A	401	NAP	O4D-C4D	4.29	1.54	1.45
2	B	401	NAP	C7N-N7N	4.21	1.41	1.33
2	A	401	NAP	C2A-N3A	3.99	1.38	1.32
2	A	401	NAP	C6A-N6A	3.49	1.46	1.34
2	B	401	NAP	C6A-N6A	3.32	1.46	1.34
2	B	401	NAP	O3B-C3B	3.30	1.50	1.43
2	B	401	NAP	P2B-O2B	3.00	1.65	1.59
2	B	401	NAP	C2A-N1A	2.80	1.39	1.33
2	A	401	NAP	O3B-C3B	2.67	1.49	1.43
2	A	401	NAP	C5A-C4A	-2.45	1.34	1.40
2	A	401	NAP	O3D-C3D	2.44	1.48	1.43
2	B	401	NAP	C5A-C4A	-2.42	1.34	1.40
2	A	401	NAP	P2B-O2B	2.35	1.63	1.59
2	B	401	NAP	O2B-C2B	-2.33	1.35	1.44
2	A	401	NAP	C2A-N1A	2.30	1.38	1.33

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	401	NAP	O3D-C3D	2.30	1.48	1.43
2	A	401	NAP	O7N-C7N	-2.29	1.19	1.24
2	B	401	NAP	O2D-C2D	2.01	1.47	1.43

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	NAP	C1B-N9A-C4A	-8.32	112.02	126.64
2	B	401	NAP	C5A-C6A-N6A	8.18	132.78	120.35
2	B	401	NAP	C1B-N9A-C4A	-7.69	113.13	126.64
2	A	401	NAP	C5A-C6A-N6A	7.21	131.31	120.35
2	B	401	NAP	N6A-C6A-N1A	-6.13	105.85	118.57
2	B	401	NAP	O7N-C7N-C3N	5.50	126.22	119.63
2	A	401	NAP	N6A-C6A-N1A	-5.43	107.29	118.57
2	A	401	NAP	N3A-C2A-N1A	-5.06	120.76	128.68
2	B	401	NAP	N3A-C2A-N1A	-4.84	121.11	128.68
2	B	401	NAP	C2B-C3B-C4B	2.98	108.46	101.99
2	B	401	NAP	O7N-C7N-N7N	-2.70	118.74	122.58
2	A	401	NAP	O7N-C7N-N7N	-2.64	118.83	122.58
2	B	401	NAP	C3N-C7N-N7N	-2.30	114.99	117.75
2	B	401	NAP	C5D-C4D-C3D	-2.28	106.62	115.18
2	B	401	NAP	O2D-C2D-C3D	-2.26	104.51	111.82
2	B	401	NAP	O3D-C3D-C2D	-2.20	104.71	111.82
2	B	401	NAP	O2D-C2D-C1D	2.10	118.60	110.85

There are no chirality outliers.

All (10) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	401	NAP	C5B-O5B-PA-O2A
2	A	401	NAP	O4D-C1D-N1N-C2N
2	A	401	NAP	O4B-C4B-C5B-O5B
2	A	401	NAP	PA-O3-PN-O2N
2	B	401	NAP	PA-O3-PN-O2N
2	A	401	NAP	C3B-C4B-C5B-O5B
2	A	401	NAP	C5B-O5B-PA-O3
2	B	401	NAP	PA-O3-PN-O1N
2	A	401	NAP	C5B-O5B-PA-O1A
2	B	401	NAP	O4B-C4B-C5B-O5B


There are no ring outliers.

2 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	401	NAP	2	0
2	B	401	NAP	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	342/342 (100%)	-0.03	11 (3%) 47 54	10, 19, 38, 60	0
1	B	342/342 (100%)	-0.20	3 (0%) 84 89	9, 16, 32, 54	0
All	All	684/684 (100%)	-0.12	14 (2%) 65 72	9, 17, 35, 60	0

All (14) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1	MET	4.4
1	A	184	ASP	3.8
1	B	184	ASP	3.4
1	B	185	ALA	2.7
1	A	72	LYS	2.5
1	A	228	ASP	2.5
1	A	112	LYS	2.4
1	A	89	LYS	2.3
1	A	230	LYS	2.3
1	A	183	SER	2.2
1	A	227	PHE	2.1
1	A	185	ALA	2.1
1	A	180	LYS	2.0
1	B	180	LYS	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	NAP	A	401	48/48	0.96	0.09	8,14,22,26	0
2	NAP	B	401	48/48	0.96	0.09	8,12,17,18	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







## 6.5 Other polymers [i](#)

There are no such residues in this entry.