



wwPDB X-ray Structure Validation Summary Report ⓘ

May 16, 2020 – 01:42 pm BST

PDB ID : 4C3I
Title : Structure of 14-subunit RNA polymerase I at 3.0 Å resolution, crystal form C2-100
Authors : Fernandez-Tornero, C.; Moreno-Morcillo, M.; Rashid, U.J.; Taylor, N.M.I.; Ruiz, F.M.; Gruene, T.; Legrand, P.; Steuerwald, U.; Muller, C.W.
Deposited on : 2013-08-24
Resolution : 3.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.











Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2092 (3.00-3.00)
Clashscore	141614	2416 (3.00-3.00)
Ramachandran outliers	138981	2333 (3.00-3.00)
Sidechain outliers	138945	2336 (3.00-3.00)
RSRZ outliers	127900	1990 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1664	
2	B	1203	
3	C	335	
4	D	137	
5	E	215	
6	F	155	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
7	G	326	
8	H	146	
9	I	125	
10	J	70	
11	K	142	
12	L	70	
13	M	415	
14	N	233	

2 Entry composition [i](#)

There are 18 unique types of molecules in this entry. The entry contains 34252 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA-DIRECTED RNA POLYMERASE I SUBUNIT RPA190.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	1484	11695	7388	2031	2213	63	0	0	0

- Molecule 2 is a protein called DNA-DIRECTED RNA POLYMERASE I SUBUNIT RPA135.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	1176	9322	5898	1629	1745	50	0	0	0

- Molecule 3 is a protein called DNA-DIRECTED RNA POLYMERASES I AND III SUBUNIT RPAC1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	C	304	2418	1536	414	460	8	0	0	0

- Molecule 4 is a protein called DNA-DIRECTED RNA POLYMERASE I SUBUNIT RPA14.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
4	D	59	466	292	80	94	0	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	12	SER	THR	conflict	UNP P50106

- Molecule 5 is a protein called DNA-DIRECTED RNA POLYMERASES I, II, AND III SUBUNIT RPABC 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
5	E	215	1759	1116	310	321	12	0	0	0

- Molecule 6 is a protein called DNA-DIRECTED RNA POLYMERASES I, II, AND III SUBUNIT RPABC 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
6	F	100	823	522	144	154	3	0	0	0

- Molecule 7 is a protein called DNA-DIRECTED RNA POLYMERASE I SUBUNIT RPA43.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
7	G	259	2052	1301	348	398	5	0	0	0

- Molecule 8 is a protein called DNA-DIRECTED RNA POLYMERASES I, II, AND III SUBUNIT RPABC 3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
8	H	134	1072	676	181	211	4	0	0	0

- Molecule 9 is a protein called DNA-DIRECTED RNA POLYMERASE I SUBUNIT RPA12.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
9	I	124	942	584	160	189	9	0	0	0

- Molecule 10 is a protein called DNA-DIRECTED RNA POLYMERASES I, II, AND III SUBUNIT RPABC 5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
10	J	69	569	362	101	100	6	0	0	0

- Molecule 11 is a protein called DNA-DIRECTED RNA POLYMERASES I AND III SUBUNIT RPAC2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
11	K	103	810	506	132	167	5	0	0	0

- Molecule 12 is a protein called DNA-DIRECTED RNA POLYMERASES I, II, AND III SUBUNIT RPABC 4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
12	L	45	359	221	71	63	4	0	0	0


- Molecule 13 is a protein called DNA-DIRECTED RNA POLYMERASE I SUBUNIT RPA49.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
13	M	105	831	528	137	166	0	0	0

- Molecule 14 is a protein called DNA-DIRECTED RNA POLYMERASE I SUBUNIT RPA34.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
14	N	139	1103	706	179	214	4	0	0	0

- Molecule 15 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
15	A	1	5	4	1	0	0
15	A	1	5	4	1	0	0
15	K	1	5	4	1	0	0


- Molecule 16 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
16	B	1	Total	Zn	0	0
			1	1		
16	A	2	Total	Zn	0	0
			2	2		
16	L	1	Total	Zn	0	0
			1	1		
16	J	1	Total	Zn	0	0
			1	1		
16	I	2	Total	Zn	0	0
			2	2		

- Molecule 17 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
17	B	1	Total	Mg	0	0
			1	1		

- Molecule 18 is (4S)-2-METHYL-2,4-PENTANEDIOL (three-letter code: MPD) (formula: C₆H₁₄O₂).





Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
18	G	1	Total	C	O	0	0
			8	6	2		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: DNA-DIRECTED RNA POLYMERASE I SUBUNIT RPA190






- Molecule 2: DNA-DIRECTED RNA POLYMERASE I SUBUNIT RPA135

Chain B: 80% 17%





- Molecule 3: DNA-DIRECTED RNA POLYMERASES I AND III SUBUNIT RPA1


Chain C: 73% 17% 9%




- Molecule 4: DNA-DIRECTED RNA POLYMERASE I SUBUNIT RPA14




● Molecule 5: DNA-DIRECTED RNA POLYMERASES I, II, AND III SUBUNIT RPABC 1





● Molecule 6: DNA-DIRECTED RNA POLYMERASES I, II, AND III SUBUNIT RPABC 2




● Molecule 7: DNA-DIRECTED RNA POLYMERASE I SUBUNIT RPA43




● Molecule 8: DNA-DIRECTED RNA POLYMERASES I, II, AND III SUBUNIT RPABC 3



● Molecule 9: DNA-DIRECTED RNA POLYMERASE I SUBUNIT RPA12



- Molecule 14: DNA-DIRECTED RNA POLYMERASE I SUBUNIT RPA34



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	400.53Å 140.22Å 122.89Å 90.00° 100.14° 90.00°	Depositor
Resolution (Å)	84.00 – 3.00 84.00 – 3.00	Depositor EDS
% Data completeness (in resolution range)	99.2 (84.00-3.00) 99.1 (84.00-3.00)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.95 (at 3.01Å)	Xtrriage
Refinement program	BUSTER 2.11.5	Depositor
R, R_{free}	0.199 , 0.231 0.219 , 0.256	Depositor DCC
R_{free} test set	6617 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	93.9	Xtrriage
Anisotropy	0.224	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.27 , 66.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	34252	wwPDB-VP
Average B, all atoms (Å ²)	120.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.30% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, MPD, ZN, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.39	0/11907	0.58	0/16088
2	B	0.39	0/9527	0.59	0/12879
3	C	0.39	0/2469	0.61	0/3347
4	D	0.39	0/472	0.53	0/639
5	E	0.40	0/1795	0.55	0/2416
6	F	0.39	0/838	0.54	0/1129
7	G	0.39	0/2094	0.58	0/2843
8	H	0.39	0/1090	0.57	0/1476
9	I	0.40	0/955	0.55	0/1288
10	J	0.40	0/578	0.62	0/775
11	K	0.39	0/821	0.60	0/1108
12	L	0.38	0/361	0.60	0/478
13	M	0.38	0/846	0.52	0/1136
14	N	0.37	0/1124	0.52	0/1512
All	All	0.39	0/34877	0.58	0/47114

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	11695	0	11764	116	0
2	B	9322	0	9187	93	0
3	C	2418	0	2401	24	0
4	D	466	0	466	3	0
5	E	1759	0	1788	7	0
6	F	823	0	841	7	0
7	G	2052	0	2016	17	0
8	H	1072	0	1042	7	0
9	I	942	0	935	7	0
10	J	569	0	585	12	0
11	K	810	0	801	11	0
12	L	359	0	381	1	0
13	M	831	0	820	10	0
14	N	1103	0	1106	6	0
15	A	10	0	0	0	0
15	K	5	0	0	0	0
16	A	2	0	0	0	0
16	B	1	0	0	0	0
16	I	2	0	0	0	0
16	J	1	0	0	0	0
16	L	1	0	0	0	0
17	B	1	0	0	0	0
18	G	8	0	14	2	0
All	All	34252	0	34147	270	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 270 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:712:ILE:H	11:K:106:GLN:HE22	1.22	0.88
2:B:99:VAL:HG21	2:B:417:ILE:HD11	1.55	0.87
2:B:16:PHE:HD2	2:B:978:ALA:HB2	1.51	0.75
1:A:86:TYR:H	1:A:431:GLN:HE22	1.36	0.72
1:A:824:THR:HG23	2:B:1023:ARG:HB2	1.71	0.71

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	1466/1664 (88%)	1393 (95%)	71 (5%)	2 (0%)	51	85
2	B	1166/1203 (97%)	1094 (94%)	70 (6%)	2 (0%)	47	82
3	C	300/335 (90%)	284 (95%)	16 (5%)	0	100	100
4	D	55/137 (40%)	52 (94%)	3 (6%)	0	100	100
5	E	213/215 (99%)	202 (95%)	11 (5%)	0	100	100
6	F	98/155 (63%)	96 (98%)	2 (2%)	0	100	100
7	G	251/326 (77%)	234 (93%)	16 (6%)	1 (0%)	34	72
8	H	130/146 (89%)	119 (92%)	10 (8%)	1 (1%)	19	57
9	I	122/125 (98%)	109 (89%)	11 (9%)	2 (2%)	9	40
10	J	67/70 (96%)	62 (92%)	5 (8%)	0	100	100
11	K	101/142 (71%)	96 (95%)	5 (5%)	0	100	100
12	L	43/70 (61%)	39 (91%)	4 (9%)	0	100	100
13	M	101/415 (24%)	93 (92%)	8 (8%)	0	100	100
14	N	131/233 (56%)	123 (94%)	8 (6%)	0	100	100
All	All	4244/5236 (81%)	3996 (94%)	240 (6%)	8 (0%)	47	82

5 of 8 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	237	GLY
2	B	532	HIS
2	B	1154	ASP
1	A	1338	ARG
8	H	84	ALA

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	1306/1465 (89%)	1234 (94%)	72 (6%)	21	57
2	B	1024/1053 (97%)	926 (90%)	98 (10%)	8	32
3	C	269/296 (91%)	244 (91%)	25 (9%)	9	33
4	D	56/116 (48%)	50 (89%)	6 (11%)	6	26
5	E	197/197 (100%)	192 (98%)	5 (2%)	47	79
6	F	90/137 (66%)	86 (96%)	4 (4%)	28	65
7	G	234/291 (80%)	219 (94%)	15 (6%)	17	51
8	H	116/128 (91%)	111 (96%)	5 (4%)	29	66
9	I	109/110 (99%)	98 (90%)	11 (10%)	7	29
10	J	64/65 (98%)	57 (89%)	7 (11%)	6	25
11	K	93/130 (72%)	84 (90%)	9 (10%)	8	31
12	L	40/57 (70%)	37 (92%)	3 (8%)	13	43
13	M	94/371 (25%)	89 (95%)	5 (5%)	22	58
14	N	128/220 (58%)	118 (92%)	10 (8%)	12	42
All	All	3820/4636 (82%)	3545 (93%)	275 (7%)	14	45

5 of 275 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	617	THR
2	B	965	GLU
11	K	99	ASN
2	B	692	THR
2	B	840	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 61 such sidechains are listed below:

Mol	Chain	Res	Type
2	B	351	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	724	GLN
9	I	97	HIS
2	B	361	HIS
2	B	480	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 8 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
15	SO4	A	2664	-	4,4,4	0.15	0	6,6,6	0.04	0
15	SO4	K	1143	-	4,4,4	0.13	0	6,6,6	0.08	0
18	MPD	G	1317	-	7,7,7	0.57	0	9,10,10	0.61	0
15	SO4	A	2665	-	4,4,4	0.14	0	6,6,6	0.08	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
18	MPD	G	1317	-	-	3/5/5/5	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
18	G	1317	MPD	O2-C2-C3-C4
18	G	1317	MPD	C2-C3-C4-C5
18	G	1317	MPD	C2-C3-C4-O4

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
18	G	1317	MPD	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	1484/1664 (89%)	-0.26	6 (0%) 92 79	62, 98, 156, 225	0
2	B	1176/1203 (97%)	-0.26	4 (0%) 94 84	60, 107, 142, 176	0
3	C	304/335 (90%)	-0.26	5 (1%) 72 44	93, 122, 160, 197	0
4	D	59/137 (43%)	-0.38	0 100 100	82, 124, 147, 158	0
5	E	215/215 (100%)	-0.26	3 (1%) 75 49	78, 136, 183, 202	0
6	F	100/155 (64%)	-0.48	0 100 100	69, 86, 125, 142	0
7	G	259/326 (79%)	-0.22	1 (0%) 92 79	71, 121, 196, 231	0
8	H	134/146 (91%)	-0.18	3 (2%) 62 33	94, 123, 152, 165	0
9	I	124/125 (99%)	0.78	20 (16%) 1 0	116, 187, 215, 228	0
10	J	69/70 (98%)	-0.12	0 100 100	94, 112, 141, 161	0
11	K	103/142 (72%)	-0.53	0 100 100	82, 106, 136, 171	0
12	L	45/70 (64%)	-0.46	0 100 100	107, 128, 149, 153	0
13	M	105/415 (25%)	-0.01	1 (0%) 82 59	137, 229, 262, 267	0
14	N	139/233 (59%)	0.16	12 (8%) 10 3	111, 249, 267, 276	0
All	All	4316/5236 (82%)	-0.22	55 (1%) 77 51	60, 111, 202, 276	0

The worst 5 of 55 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
9	I	98	THR	8.2
9	I	99	LEU	7.0
9	I	118	GLY	6.9
9	I	105	ASP	5.9
9	I	87	PRO	5.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
15	SO4	A	2665	5/5	0.86	0.20	166,167,168,168	0
18	MPD	G	1317	8/8	0.86	0.37	99,101,103,104	0
15	SO4	A	2664	5/5	0.87	0.12	177,177,178,180	0
16	ZN	I	1127	1/1	0.88	0.11	153,153,153,153	0
17	MG	B	2204	1/1	0.92	0.90	89,89,89,89	0
15	SO4	K	1143	5/5	0.96	0.13	147,147,149,149	0
16	ZN	A	2667	1/1	0.99	0.14	95,95,95,95	0
16	ZN	A	2666	1/1	0.99	0.14	108,108,108,108	0
16	ZN	I	1126	1/1	0.99	0.12	138,138,138,138	0
16	ZN	L	1071	1/1	0.99	0.11	125,125,125,125	0
16	ZN	J	1070	1/1	1.00	0.23	104,104,104,104	0
16	ZN	B	2205	1/1	1.00	0.17	79,79,79,79	0

6.5 Other polymers [i](#)

There are no such residues in this entry.