

## wwPDB X-ray Structure Validation Summary Report (i)

#### Nov 19, 2023 – 10:53 PM JST

PDB ID	:	7C5Q
Title	:	Crystal Structure of H177A mutant Glyceraldehyde-3-phosphate dehydroge-
		nase1 from Escherichia coli complexed with BPG at $2.13$ Angstrom resolution
Authors	:	Zhang, L.; Liu, M.R.; Bao, L.Y.; Yao, Y.C.; Bostrom, I.K.; Wang, Y.D.; Chen,
		A.Q.; Li, J.X.; Gu, S.H.; Ji, C.N.
Deposited on	:	2020-05-20
Resolution	:	2.13  Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:


		4 001 407
MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
$\mathrm{EDS}$	:	2.36
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36

## 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:  $X\text{-}RAY \, DIFFRACTION$ 

The reported resolution of this entry is 2.13 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Percentile relative to X-ray structures of similar resolution

Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$		
$R_{free}$	130704	2523 (2.16-2.12)		
Clashscore	141614	2653 (2.16-2.12)		
Ramachandran outliers	138981	2618 (2.16-2.12)		
Sidechain outliers	138945	2617 (2.16-2.12)		
RSRZ outliers	127900	2485 (2.16-2.12)		

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	О	352	90%	5%	5%
1	Р	352	90%	5%	5%
1	Q	352	91%	·	5%
1	R	352	% 91%	•	5%



The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry Clash		Electron density
3	G3H	Q	402	-	-	Х	-
3	G3H	R	402	-	-	Х	-



 $\mathbf{2}$ 

# Entry composition (i)

There are 8 unique types of molecules in this entry. The entry contains 21814 atoms, of which 10407 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues		Atoms					ZeroOcc	AltConf	Trace
1	0	334	Total	С	Η	Ν	0	S	0	14	0
1	0	004	5171	1624	2596	432	512	7	0	14	0
1	р	333	Total	С	Η	Ν	0	S	0	7	0
1	1	000	5113	1606	2571	427	502	7	0		
1	0	333	Total	С	Η	Ν	0	S	0	2	0
	I Q		5069	1593	2550	423	496	$\overline{7}$	0		0
1	1 R	333	Total	С	Н	Ν	0	S	0	0	0
			5053	1588	2543	422	493	7		0	U

• Molecule 1 is a protein called Glyceraldehyde-3-phosphate dehydrogenase.

There are 80 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
0	-18	HIS	-	expression tag	UNP A0A140NCK4
0	-17	HIS	-	expression tag	UNP A0A140NCK4
0	-16	HIS	-	expression tag	UNP A0A140NCK4
0	-15	HIS	-	expression tag	UNP A0A140NCK4
0	-14	HIS	-	expression tag	UNP A0A140NCK4
0	-13	HIS	-	expression tag	UNP A0A140NCK4
0	-12	SER	-	expression tag	UNP A0A140NCK4
0	-11	SER	-	expression tag	UNP A0A140NCK4
0	-10	GLY	-	expression tag	UNP A0A140NCK4
0	-9	LEU	-	expression tag	UNP A0A140NCK4
0	-8	VAL	-	expression tag	UNP A0A140NCK4
0	-7	PRO	-	expression tag	UNP A0A140NCK4
0	-6	ARG	-	expression tag	UNP A0A140NCK4
0	-5	GLY	-	expression tag	UNP A0A140NCK4
0	-4	SER	-	expression tag	UNP A0A140NCK4
0	-3	HIS	-	expression tag	UNP A0A140NCK4
0	-2	MET	-	expression tag	UNP A0A140NCK4
0	-1	ALA	-	expression tag	UNP A0A140NCK4
0	0	SER	-	expression tag	UNP A0A140NCK4
0	177	ALA	HIS	engineered mutation	UNP A0A140NCK4
Р	-18	HIS	_	expression tag	UNP A0A140NCK4




7	C5	Q
•	$\sim \circ$	~€

Continued from previous page										
Chain	Residue	Modelled	Actual	Comment	Reference					
Р	-17	HIS	-	expression tag	UNP A0A140NCK4					
Р	-16	HIS	-	expression tag	UNP A0A140NCK4					
Р	-15	HIS	-	expression tag	UNP A0A140NCK4					
Р	-14	HIS	-	expression tag	UNP A0A140NCK4					
Р	-13	HIS	-	expression tag	UNP A0A140NCK4					
Р	-12	SER	-	expression tag	UNP A0A140NCK4					
Р	-11	SER	-	expression tag	UNP A0A140NCK4					
Р	-10	GLY	-	expression tag	UNP A0A140NCK4					
Р	-9	LEU	-	expression tag	UNP A0A140NCK4					
Р	-8	VAL	-	expression tag	UNP A0A140NCK4					
Р	-7	PRO	-	expression tag	UNP A0A140NCK4					
Р	-6	ARG	-	expression tag	UNP A0A140NCK4					
Р	-5	GLY	-	expression tag	UNP A0A140NCK4					
Р	-4	SER	-	expression tag	UNP A0A140NCK4					
Р	-3	HIS	-	expression tag	UNP A0A140NCK4					
Р	-2	MET	-	expression tag	UNP A0A140NCK4					
Р	-1	ALA	-	expression tag	UNP A0A140NCK4					
Р	0	SER	-	expression tag	UNP A0A140NCK4					
Р	177	ALA	HIS	engineered mutation	UNP A0A140NCK4					
$\mathbf{Q}$	-18	HIS	-	expression tag	UNP A0A140NCK4					
Q	-17	HIS	-	expression tag	UNP A0A140NCK4					
Q	-16	HIS	-	expression tag	UNP A0A140NCK4					
Q	-15	HIS	-	expression tag	UNP A0A140NCK4					
Q	-14	HIS	-	expression tag	UNP A0A140NCK4					
Q	-13	HIS	-	expression tag	UNP A0A140NCK4					
Q	-12	SER	-	expression tag	UNP A0A140NCK4					
$\mathbf{Q}$	-11	SER	-	expression tag	UNP A0A140NCK4					
$\mathbf{Q}$	-10	GLY	-	expression tag	UNP A0A140NCK4					
Q	-9	LEU	-	expression tag	UNP A0A140NCK4					
$\mathbf{Q}$	-8	VAL	-	expression tag	UNP A0A140NCK4					
Q	-7	PRO	-	expression tag	UNP A0A140NCK4					
Q	-6	ARG	-	expression tag	UNP A0A140NCK4					
Q	-5	GLY	-	expression tag	UNP A0A140NCK4					
Q	-4	SER	-	expression tag	UNP A0A140NCK4					
Q	-3	HIS	-	expression tag	UNP A0A140NCK4					
Q	-2	MET	-	expression tag	UNP A0A140NCK4					
Q	-1	ALA	-	expression tag	UNP A0A140NCK4					
Q	0	SER	-	expression tag	UNP A0A140NCK4					
Q	177	ALA	HIS	engineered mutation	UNP A0A140NCK4					
R	-18	HIS	-	expression tag	UNP A0A140NCK4					
R	-17	HIS	-	expression tag	UNP A0A140NCK4					
R	-16	HIS	-	expression tag	UNP A0A140NCK4					



Chain	Residue	Modelled	Actual	Comment	Reference
R	-15	HIS	-	expression tag	UNP A0A140NCK4
R	-14	HIS	-	expression tag	UNP A0A140NCK4
R	-13	HIS	-	expression tag	UNP A0A140NCK4
R	-12	SER	-	expression tag	UNP A0A140NCK4
R	-11	SER	-	expression tag	UNP A0A140NCK4
R	-10	GLY	-	expression tag	UNP A0A140NCK4
R	-9	LEU	-	expression tag	UNP A0A140NCK4
R	-8	VAL	-	expression tag	UNP A0A140NCK4
R	-7	PRO	-	expression tag	UNP A0A140NCK4
R	-6	ARG	-	expression tag	UNP A0A140NCK4
R	-5	GLY	-	expression tag	UNP A0A140NCK4
R	-4	SER	-	expression tag	UNP A0A140NCK4
R	-3	HIS	-	expression tag	UNP A0A140NCK4
R	-2	MET	-	expression tag	UNP A0A140NCK4
R	-1	ALA	-	expression tag	UNP A0A140NCK4
R	0	SER	-	expression tag	UNP A0A140NCK4
R	177	ALA	HIS	engineered mutation	UNP A0A140NCK4

• Molecule 2 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (three-letter code: NAD) (formula:  $C_{21}H_{27}N_7O_{14}P_2$ ) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
0	0	1	Total	С	Η	Ν	Ο	Р	0	0
	2 0	1	70	21	26	7	14	2	0	0
0	D	1	Total	С	Η	Ν	Ο	Р	0	0
	2 P	1	70	21	26	7	14	2	U	U



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf			
2	0	1	Total	С	Η	Ν	Ο	Р	0	0
	Q	1	70	21	26	7	14	2	0	0
0	D	1	Total	С	Η	Ν	Ο	Р	0	0
	n		70	21	26	7	14	2	0	0


• Molecule 3 is GLYCERALDEHYDE-3-PHOSPHATE (three-letter code: G3H) (formula:  $C_3H_7O_6P$ ) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	О	1	Total         C         H         O         P           15         3         5         6         1	0	0
3	Q	1	Total         C         H         O         P           15         3         5         6         1	0	0
3	R	1	Total         C         H         O         P           15         3         5         6         1	0	0

• Molecule 4 is PHOSPHATE ION (three-letter code: PO4) (formula:  $O_4P$ ) (labeled as "Ligand of Interest" by depositor).






Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	О	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{P} \\ 5 & 4 & 1 \end{array}$	0	0
4	О	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{P} \\ 5 & 4 & 1 \end{array}$	0	0
4	О	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{P} \\ 5 & 4 & 1 \end{array}$	0	0
4	Р	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{P} \\ 5 & 4 & 1 \end{array}$	0	0
4	Р	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{P} \\ 5 & 4 & 1 \end{array}$	0	0
4	Q	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{P} \\ 5 & 4 & 1 \end{array}$	0	0
4	Q	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{P} \\ 5 & 4 & 1 \end{array}$	0	0
4	R	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{P} \\ 5 & 4 & 1 \end{array}$	0	0
4	R	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{P} \\ 5 & 4 & 1 \end{array}$	0	0

• Molecule 5 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula:  $C_2H_6O_2$ ) (labeled as "Ligand of Interest" by depositor).






Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	О	1	$\begin{array}{cccc} \text{Total} & \text{C} & \text{H} & \text{O} \\ 10 & 2 & 6 & 2 \end{array}$	0	0
5	Q	1	Total         C         H         O           20         4         12         4	0	1
5	R	1	Total         C         H         O           10         2         6         2	0	0

• Molecule 6 is CHLORIDE ION (three-letter code: CL) (formula: Cl) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	О	2	Total Cl 2 2	0	0
6	Р	1	Total Cl 1 1	0	0
6	Q	1	Total Cl 1 1	0	0
6	R	1	Total Cl 1 1	0	0

• Molecule 7 is phosphono (2S)-2-oxidanyl-3-phosphonooxy-propanoate (three-letter code: DG4) (formula:  $C_3H_8O_{10}P_2$ ) (labeled as "Ligand of Interest" by depositor).





Mol	Chain	Residues	Atoms			ZeroOcc	AltConf		
7	D	1	Total	С	Η	Ο	Р	0	0
1	Ľ	1	19	3	4	10	2	0	U

• Molecule 8 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	О	276	Total O 276 276	0	0
8	Р	270	Total         O           270         270	0	0
8	Q	253	Total O 254 254	0	1
8	R	173	Total O 174 174	0	1



## 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Glyceraldehyde-3-phosphate dehydrogenase





## 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants	89.67Å 89.67Å 341.80Å	Depositor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $90.00^{\circ}$ $90.00^{\circ}$	Depositor
Resolution (Å)	48.08 - 2.13	Depositor
Resolution (A)	48.08 - 2.13	EDS
% Data completeness	99.9 (48.08-2.13)	Depositor
(in resolution range)	99.9 (48.08-2.13)	EDS
R <sub>merge</sub>	0.15	Depositor
R <sub>sym</sub>	(Not available)	Depositor
$< I/\sigma(I) > 1$	$3.46 (at 2.12 \text{\AA})$	Xtriage
Refinement program	PHENIX (1.10_2155: ???)	Depositor
D D.	0.138 , $0.190$	Depositor
$R, R_{free}$	0.139 , $0.192$	DCC
$R_{free}$ test set	3931 reflections $(4.96%)$	wwPDB-VP
Wilson B-factor $(Å^2)$	21.9	Xtriage
Anisotropy	0.036	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.38 , $49.8$	EDS
L-test for twinning <sup>2</sup>	$ \langle L  \rangle = 0.49, \langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	21814	wwPDB-VP
Average B, all atoms $(Å^2)$	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 4.43% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality (i)

### 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: NAD, PO4, EDO, G3H, CL, DG4  $\,$ 

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Chain		Bond	lengths	Bond angles		
	Unam	RMSZ	# Z  > 5	RMSZ	# Z  > 5	
1	0	0.50	0/2665	0.68	0/3617	
1	Р	0.50	0/2600	0.67	0/3529	
1	Q	0.50	0/2568	0.65	0/3483	
1	R	0.46	0/2550	0.64	0/3460	
All	All	0.49	0/10383	0.66	0/14089	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	0	2575	2596	2531	8	0
1	Р	2542	2571	2552	10	0
1	Q	2519	2550	2538	10	0
1	R	2510	2543	2545	10	0
2	0	44	26	25	1	0
2	Р	44	26	24	0	0
2	Q	44	26	25	2	0
2	R	44	26	23	2	0
3	0	10	5	5	2	0



	-	<i>i</i> previous				
Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	Q	10	5	5	5	0
3	R	10	5	5	4	0
4	0	15	0	0	0	0
4	Р	10	0	0	1	0
4	Q	10	0	0	0	0
4	R	10	0	0	1	0
5	0	4	6	6	0	0
5	Q	8	12	12	0	0
5	R	4	6	6	1	0
6	0	2	0	0	0	0
6	Р	1	0	0	0	0
6	Q	1	0	0	0	0
6	R	1	0	0	0	0
7	Р	15	4	0	3	0
8	0	276	0	0	1	0
8	Р	270	0	0	2	0
8	Q	254	0	0	0	0
8	R	174	0	0	0	0
All	All	11407	10407	10302	44	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 44 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:R:182:THR:HG21	3:R:402:G3H:H32	1.61	0.81
1:R:182:THR:CG2	3:R:402:G3H:H32	2.26	0.65
1:P:28:ASP:OD1	1:P:70:LYS:NZ	2.27	0.60
2:Q:401:NAD:PN	3:Q:402:G3H:H2	2.45	0.57
1:R:19:ARG:HH22	5:R:405:EDO:H12	1.70	0.57

There are no symmetry-related clashes.

### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	Ο	346/352~(98%)	331~(96%)	15~(4%)	0	100	100
1	Р	338/352~(96%)	325~(96%)	13~(4%)	0	100	100
1	Q	333/352~(95%)	323~(97%)	10 (3%)	0	100	100
1	R	331/352~(94%)	316~(96%)	15~(4%)	0	100	100
All	All	1348/1408~(96%)	1295 (96%)	53 (4%)	0	100	100

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

There are no Ramachandran outliers to report.

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles		
1	Ο	286/288~(99%)	283~(99%)	3~(1%)	76	79	
1	Р	278/288~(96%)	276~(99%)	2(1%)	84	87	
1	Q	274/288~(95%)	273 (100%)	1 (0%)	91	93	
1	R	272/288~(94%)	271 (100%)	1 (0%)	91	93	
All	All	1110/1152~(96%)	1103 (99%)	7 (1%)	86	89	

5 of 7 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	Р	229	HIS
1	Р	303	ASP
1	R	229	HIS
1	Q	229	HIS
1	0	330	PHE

Sometimes side chains can be flipped to improve hydrogen bonding and reduce clashes. There are no such side chains identified.



#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

#### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

#### 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

#### 5.6 Ligand geometry (i)

Of 26 ligands modelled in this entry, 5 are monoatomic - leaving 21 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	B	ond leng	gths	B	Sond ang	gles
MOI	Type	Ullalli	nes	LIIIK	Counts	RMSZ	# Z >2	Counts	RMSZ	# Z >2
2	NAD	Ο	401	-	42,48,48	4.68	17 (40%)	50,73,73	1.74	8 (16%)
4	PO4	0	408	-	4,4,4	0.76	0	$6,\!6,\!6$	0.54	0
4	PO4	R	403	-	4,4,4	0.60	0	$6,\!6,\!6$	0.77	0
5	EDO	Q	405[A]	-	3, 3, 3	0.57	0	$2,\!2,\!2$	0.19	0
4	PO4	0	403	-	$4,\!4,\!4$	0.92	0	$6,\!6,\!6$	0.56	0
7	DG4	Р	402	-	12,14,14	1.90	3 (25%)	$16,\!21,\!21$	1.97	4 (25%)
5	EDO	Q	405[B]	-	3,3,3	0.47	0	2,2,2	0.42	0
4	PO4	0	404	-	4,4,4	0.76	0	$6,\!6,\!6$	0.53	0
4	PO4	R	404	-	4,4,4	0.83	0	$6,\!6,\!6$	0.59	0
3	G3H	R	402	-	$8,\!9,\!9$	1.74	3 (37%)	$10,\!12,\!12$	1.93	2 (20%)
4	PO4	Р	404	-	4,4,4	0.82	0	$6,\!6,\!6$	0.50	0
2	NAD	Р	401	-	42,48,48	4.74	17 (40%)	50,73,73	1.63	10 (20%)
5	EDO	0	405	-	3,3,3	0.52	0	2,2,2	0.21	0
4	PO4	Р	403	-	4,4,4	0.82	0	$6,\!6,\!6$	0.72	0
5	EDO	R	405	-	3,3,3	0.46	0	2,2,2	0.85	0
3	G3H	Q	402	-	$8,\!9,\!9$	0.99	0	$10,\!12,\!12$	1.74	2 (20%)



Mal	Mol Type Chain		Res	Link	Bond lengths			Bond angles		
		nes	LIIIK	Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z >2	
4	PO4	Q	404	-	4,4,4	0.79	0	6,6,6	0.52	0
4	PO4	Q	403	-	4,4,4	0.69	0	6,6,6	0.78	0
3	G3H	Ο	402	-	$8,\!9,\!9$	1.26	1 (12%)	10,12,12	1.45	3 (30%)
2	NAD	R	401	-	42,48,48	4.97	16 (38%)	50,73,73	1.59	8 (16%)
2	NAD	Q	401	-	42,48,48	4.58	16 (38%)	50,73,73	1.94	10 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAD	Р	401	-	-	5/26/62/62	0/5/5/5
5	EDO	0	405	-	-	1/1/1/1	-
5	EDO	R	405	-	-	0/1/1/1	-
2	NAD	Ο	401	-	-	5/26/62/62	0/5/5/5
3	G3H	Q	402	-	-	5/7/8/8	-
5	EDO	Q	405[B]	-	-	0/1/1/1	-
5	EDO	Q	405[A]	-	-	1/1/1/1	-
7	DG4	Р	402	-	-	8/13/15/15	-
3	G3H	R	402	-	-	3/7/8/8	-
3	G3H	Ο	402	-	-	4/7/8/8	-
2	NAD	R	401	-	-	5/26/62/62	0/5/5/5
2	NAD	Q	401	-	-	5/26/62/62	0/5/5/5

The worst 5 of 73 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	R	401	NAD	C2D-C1D	-15.11	1.30	1.53
2	R	401	NAD	O4D-C1D	14.20	1.60	1.41
2	R	401	NAD	C2B-C1B	-14.13	1.32	1.53
2	Р	401	NAD	C2D-C1D	-14.07	1.32	1.53
2	0	401	NAD	C2D-C1D	-14.06	1.32	1.53

The worst 5 of 47 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Ζ	$Observed(^{o})$	$Ideal(^{o})$
2	Q	401	NAD	O7N-C7N-C3N	7.45	128.55	119.63
2	0	401	NAD	C5A-C6A-N6A	5.74	129.07	120.35



Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
2	R	401	NAD	N3A-C2A-N1A	-5.37	120.28	128.68
2	Р	401	NAD	N3A-C2A-N1A	-5.24	120.49	128.68
2	0	401	NAD	N3A-C2A-N1A	-5.17	120.59	128.68

There are no chirality outliers.

5 of 42 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	0	401	NAD	O4D-C1D-N1N-C2N
2	0	401	NAD	O4D-C1D-N1N-C6N
2	0	401	NAD	C2D-C1D-N1N-C2N
2	0	401	NAD	C2D-C1D-N1N-C6N
2	Р	401	NAD	O4D-C1D-N1N-C2N


There are no ring outliers.

10 monomers are involved in 19 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	0	401	NAD	1	0
4	R	403	PO4	1	0
7	Р	402	DG4	3	0
3	R	402	G3H	4	0
4	Р	403	PO4	1	0
5	R	405	EDO	1	0
3	Q	402	G3H	5	0
3	0	402	G3H	2	0
2	R	401	NAD	2	0
2	Q	401	NAD	2	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and sufficient the outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.
































































































## 5.7 Other polymers (i)

There are no such residues in this entry.

### 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



## 6 Fit of model and data (i)

## 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	$\langle RSRZ \rangle$	#RSRZ>2	$\mathbf{OWAB}(\mathbf{A}^2)$	Q < 0.9
1	Ο	334/352~(94%)	-0.68	0 100 100	13, 19, 35, 51	0
1	Р	333/352~(94%)	-0.68	1 (0%) 94 95	13, 20, 39, 62	0
1	Q	333/352~(94%)	-0.60	0 100 100	13, 21, 40, 70	0
1	R	333/352~(94%)	-0.31	5 (1%) 73 78	13, 26, 50, 66	0
All	All	1333/1408~(94%)	-0.57	6 (0%) 91 93	13, 21, 42, 70	0

The worst 5 of 6 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	Р	1	MET	2.8
1	R	110	LEU	2.2
1	R	103	ALA	2.2
1	R	92	ILE	2.1
1	R	114	ALA	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

### 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




7	C5O	
•	000	


Mol	Type	Chain	Res	Atoms	RSCC	RSR	$B-factors(Å^2)$	Q<0.9
4	PO4	0	408	5/5	0.83	0.20	41,45,51,107	0
5	EDO	0	405	4/4	0.83	0.20	34,41,46,49	0
4	PO4	Р	404	5/5	0.84	0.25	66,72,78,96	0
5	EDO	R	405	4/4	0.84	0.15	30,43,53,64	0
5	EDO	Q	405[B]	4/4	0.86	0.17	23,29,37,37	10
5	EDO	Q	405[A]	4/4	0.86	0.17	$25,\!38,\!46,\!46$	10
7	DG4	Р	402	15/15	0.88	0.20	11,38,53,55	19
4	PO4	R	404	5/5	0.89	0.27	55,56,88,90	0
3	G3H	Q	402	10/10	0.89	0.27	19,29,52,63	15
3	G3H	R	402	10/10	0.92	0.22	$25,\!43,\!51,\!54$	15
4	PO4	Q	404	5/5	0.92	0.37	$61,\!65,\!92,\!109$	0
4	PO4	0	404	5/5	0.93	0.17	54,62,85,88	0
3	G3H	0	402	10/10	0.93	0.18	$21,\!35,\!47,\!52$	15
4	PO4	Р	403	5/5	0.94	0.11	$37,\!48,\!52,\!58$	0
4	PO4	R	403	5/5	0.94	0.17	$35,\!52,\!55,\!62$	0
4	PO4	Q	403	5/5	0.95	0.16	32,51,65,66	0
2	NAD	0	401	44/44	0.98	0.09	$11,\!17,\!22,\!25$	0
2	NAD	Р	401	44/44	0.98	0.08	13,19,26,31	0
2	NAD	Q	401	44/44	0.98	0.09	13,20,30,33	0
6	CL	0	406	1/1	0.98	0.06	$27,\!27,\!27,\!27$	0
2	NAD	R	401	44/44	0.98	0.08	18,23,31,38	0
6	CL	Р	405	1/1	0.99	0.06	22,22,22,22	0
6	CL	Q	406	1/1	0.99	0.05	26,26,26,26	0
6	CL	R	406	1/1	0.99	0.07	29,29,29,29	0
4	PO4	0	403	5/5	0.99	0.10	28,34,39,51	0
6	CL	0	407	1/1	1.00	0.07	20,20,20,20	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




























































































































## 6.5 Other polymers (i)

There are no such residues in this entry.

