



wwPDB X-ray Structure Validation Summary Report ⓘ

Nov 20, 2023 – 12:17 AM JST

PDB ID : 7C7X
Title : Structural insights into nucleosome reorganization by NAP1-RELATED PROTEIN 1 (NRP1)
Authors : Luo, Q.; Baihui, W.
Deposited on : 2020-05-27
Resolution : 3.00 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2092 (3.00-3.00)
Clashscore	141614	2416 (3.00-3.00)
Ramachandran outliers	138981	2333 (3.00-3.00)
Sidechain outliers	138945	2336 (3.00-3.00)
RSRZ outliers	127900	1990 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	93	 4% 62% 23% 13%
1	C	93	 4% 57% 28% 13%
2	B	98	 74% 14% 7%
2	D	98	 69% 17% 5% 8%
3	E	239	 5% 58% 13% 27%
3	F	239	 8% 49% 17% 31%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 5502 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Histone H2A.6.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
1	C	81	620	396	114	110	0	0	0
1	A	81	615	393	112	110	0	0	0

- Molecule 2 is a protein called Histone H2B.1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	D	90	707	455	120	130	2	0	0	0
2	B	91	715	459	122	132	2	0	0	0

- Molecule 3 is a protein called NAP1-related protein 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	E	174	1426	922	221	282	1	0	0	0
3	F	166	1371	890	211	269	1	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
E	18	SER	-	expression tag	UNP Q9CA59
F	18	SER	-	expression tag	UNP Q9CA59

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	D	1	Total C O 6 3 3	0	0
4	D	1	Total C O 6 3 3	0	0
4	D	1	Total C O 6 3 3	0	0
4	F	1	Total C O 6 3 3	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	C	4	Total O 4 4	0	0
5	D	3	Total O 3 3	0	0
5	E	3	Total O 3 3	0	0
5	F	5	Total O 5 5	0	0
5	A	6	Total O 6 6	0	0
5	B	3	Total O 3 3	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Histone H2A.6




- Molecule 1: Histone H2A.6




- Molecule 2: Histone H2B.1



- Molecule 2: Histone H2B.1



- Molecule 3: NAP1-related protein 1



ASP
GLY
ASP
ASP
ASP
GLY
ASP
GLU
GLU
GLY
GLU
GLU
ASP
ASP
ASP
ASP
GLU
GLU
GLU
GLU
GLU
GLY
GLU
GLU

● Molecule 3: NAP1-related protein 1



SER
ASN
LEU
E21
Q22
I23
D24
I31
E32
E36
D39
K47
A48
S49
L53
E54
V55
E56
Y59
V66
V73
F80
W81
F85
L91
G92
D93
L94
L95
T96
E97
E98
D99
Q100
K101
I102
F103
K104
Y105
L106
N107
S108
D113
D116
S119

T128
S129
M130
F131
F132
F133
E134
F144
L145
GLU
E147
G148
T149
T150
K151
A154
K158
W159
K160
E161
GLY
LYS
GLY
LEU
PRO
ASN
GLY
VAL
ASN
HIS
ASP
ASP
ASP
LYS
LYS
GLY
ASN
LYS
ARG
ALA
LEU
PRO
GLU
SER
PHE
F187
F190
T191
ASP
ALA
GLN
HIS
LYS
GLU
ASP
ALA

GLY
ASP
GLU
ILE
H204
D205
E206
S217
F223
N224
ASN
ASP
ALA
ASP
GLU
GLU
ASP
PHE
ASP
GLY
ASP
ASP
GLY
ASP
GLU
GLU
GLU
GLY
GLU
GLU
ASP
ASP
ASP
ASP
GLU
GLU
GLU
GLU
GLY
GLU

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	66.71Å 128.40Å 140.28Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.62 – 3.00 29.60 – 2.99	Depositor EDS
% Data completeness (in resolution range)	84.9 (29.62-3.00) 84.2 (29.60-2.99)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.10 (at 3.00Å)	Xtrriage
Refinement program	REFMAC 5.8.0253	Depositor
R, R_{free}	0.223 , 0.262 0.227 , 0.260	Depositor DCC
R_{free} test set	1044 reflections (4.94%)	wwPDB-VP
Wilson B-factor (Å ²)	54.9	Xtrriage
Anisotropy	0.201	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 68.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.39$, $\langle L^2 \rangle = 0.21$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	5502	wwPDB-VP
Average B, all atoms (Å ²)	80.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.97% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.73	1/622 (0.2%)	0.85	0/840
1	C	0.70	0/628	0.84	0/848
2	B	0.70	0/726	0.87	0/976
2	D	0.71	0/718	0.90	0/965
3	E	0.71	0/1457	0.86	1/1969 (0.1%)
3	F	0.67	0/1401	0.82	0/1895
All	All	0.70	1/5552 (0.0%)	0.85	1/7493 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	69	ALA	C-O	5.64	1.34	1.23

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	E	192	ASP	CB-CG-OD2	5.16	122.94	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	615	0	647	31	0
1	C	620	0	652	21	0
2	B	715	0	753	11	0
2	D	707	0	747	23	0
3	E	1426	0	1382	19	0
3	F	1371	0	1333	55	0
4	D	18	0	24	2	0
4	F	6	0	8	0	0
5	A	6	0	0	2	0
5	B	3	0	0	0	0
5	C	4	0	0	1	0
5	D	3	0	0	1	0
5	E	3	0	0	1	0
5	F	5	0	0	2	0
All	All	5502	0	5546	139	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

The worst 5 of 139 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:F:132:PHE:O	3:F:160:LYS:N	1.81	1.14
3:F:131:PRO:O	3:F:160:LYS:HB3	1.51	1.10
3:F:131:PRO:O	3:F:160:LYS:CB	1.99	1.09
1:A:81:ARG:HD2	1:A:102:ILE:HD11	1.33	1.09
3:F:95:LEU:HD11	3:F:103:PHE:CE2	1.93	1.02

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	79/93 (85%)	69 (87%)	10 (13%)	0	100	100
1	C	79/93 (85%)	65 (82%)	11 (14%)	3 (4%)	3	18
2	B	89/98 (91%)	83 (93%)	5 (6%)	1 (1%)	14	50
2	D	88/98 (90%)	81 (92%)	7 (8%)	0	100	100
3	E	166/239 (70%)	141 (85%)	23 (14%)	2 (1%)	13	48
3	F	158/239 (66%)	133 (84%)	25 (16%)	0	100	100
All	All	659/860 (77%)	572 (87%)	81 (12%)	6 (1%)	17	55

5 of 6 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	E	223	PHE
1	C	100	VAL
1	C	74	LYS
2	B	70	LYS
3	E	37	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	60/70 (86%)	56 (93%)	4 (7%)	16	49
1	C	61/70 (87%)	56 (92%)	5 (8%)	11	39
2	B	80/87 (92%)	71 (89%)	9 (11%)	6	24
2	D	79/87 (91%)	70 (89%)	9 (11%)	5	24
3	E	159/214 (74%)	144 (91%)	15 (9%)	8	32
3	F	153/214 (72%)	131 (86%)	22 (14%)	3	15
All	All	592/742 (80%)	528 (89%)	64 (11%)	6	26

5 of 64 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	60	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	102	SER
3	E	130	ASN
3	E	116	ASP
2	B	115	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 6 such sidechains are listed below:

Mol	Chain	Res	Type
3	E	130	ASN
3	F	100	GLN
3	F	130	ASN
3	E	22	GLN
2	D	73	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	GOL	F	301	-	5,5,5	0.57	0	5,5,5	0.47	0
4	GOL	D	201	-	5,5,5	0.09	0	5,5,5	0.37	0
4	GOL	D	202	-	5,5,5	0.58	0	5,5,5	0.36	0
4	GOL	D	203	-	5,5,5	0.65	0	5,5,5	0.45	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	F	301	-	-	4/4/4/4	-
4	GOL	D	201	-	-	2/4/4/4	-
4	GOL	D	202	-	-	0/4/4/4	-
4	GOL	D	203	-	-	1/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 7 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	D	201	GOL	O1-C1-C2-C3
4	F	301	GOL	C1-C2-C3-O3
4	F	301	GOL	O1-C1-C2-C3
4	D	201	GOL	O1-C1-C2-O2
4	F	301	GOL	O1-C1-C2-O2

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	D	201	GOL	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	81/93 (87%)	-0.27	1 (1%) 79 54	25, 65, 118, 144	0
1	C	81/93 (87%)	-0.22	4 (4%) 29 11	25, 63, 119, 154	0
2	B	91/98 (92%)	-0.53	0 100 100	25, 52, 94, 105	0
2	D	90/98 (91%)	-0.54	0 100 100	26, 52, 93, 115	0
3	E	174/239 (72%)	0.13	12 (6%) 16 5	39, 89, 156, 184	0
3	F	166/239 (69%)	0.24	18 (10%) 5 2	43, 93, 167, 190	0
All	All	683/860 (79%)	-0.11	35 (5%) 28 10	25, 71, 152, 190	0

The worst 5 of 35 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	F	147	GLU	6.2
3	E	161	GLU	4.8
3	E	162	GLY	4.2
1	C	73	ASN	4.1
3	F	150	THR	3.9

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	GOL	D	203	6/6	0.83	0.21	28,31,31,31	0
4	GOL	D	202	6/6	0.92	0.27	28,30,30,30	0
4	GOL	D	201	6/6	0.94	0.22	53,57,59,61	0
4	GOL	F	301	6/6	0.95	0.30	30,32,32,32	0

6.5 Other polymers [i](#)

There are no such residues in this entry.