

wwPDB X-ray Structure Validation Summary Report (i)

Aug 20, 2020 – 08:50 PM BST

PDB ID : 3C9T

Title : AaThiL complexed with AMPPCP and TMP

Authors: McCulloch, K.M.; Kinsland, C.; Begley, T.P.; Ealick, S.E.

Deposited on : 2008-02-18

Resolution : 2.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS: 2.13.1

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac : 5.8.0158

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.13.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \text{Whole archive} \\ (\#\text{Entries}) \end{array}$	$\begin{array}{c} {\rm Similar \; resolution} \\ (\#{\rm Entries, \; resolution \; range(\AA)}) \end{array}$
R_{free}	130704	3163 (2.60-2.60)
Clashscore	141614	3518 (2.60-2.60)
Ramachandran outliers	138981	3455 (2.60-2.60)
Sidechain outliers	138945	3455 (2.60-2.60)
RSRZ outliers	127900	3104 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain				
4		0.40	% •				
1	A	342	55%	32%	•	10%	
	_		% ■				
1	В	342	57%	30%	٠	10%	



2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 5011 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Thiamine monophosphate kinase.

\mathbf{Mol}	Chain	Residues		\mathbf{At}	oms			ZeroOcc	AltConf	Trace
1	Λ	307	Total	С	N	О	S	0	0	0
1	А	307	2432	1562	404	457	9	U	U	U
1	D	309	Total	С	N	О	S	0	0	0
1	Ъ	309	2449	1574	406	460	9	U	0	

There are 72 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-35	MET	-	EXPRESSION TAG	UNP O67883
A	-34	GLY	-	EXPRESSION TAG	UNP O67883
A	-33	SER	_	EXPRESSION TAG	UNP O67883
A	-32	HIS	-	EXPRESSION TAG	UNP O67883
A	-31	HIS	_	EXPRESSION TAG	UNP O67883
A	-30	HIS	-	EXPRESSION TAG	UNP O67883
A	-29	HIS	-	EXPRESSION TAG	UNP O67883
A	-28	HIS	-	EXPRESSION TAG	UNP O67883
A	-27	HIS	-	EXPRESSION TAG	UNP O67883
A	-26	ASP	-	EXPRESSION TAG	UNP O67883
A	-25	ILE	_	EXPRESSION TAG	UNP O67883
A	-24	THR	-	EXPRESSION TAG	UNP O67883
A	-23	SER	-	EXPRESSION TAG	UNP O67883
A	-22	LEU	-	EXPRESSION TAG	UNP O67883
A	-21	TYR	-	EXPRESSION TAG	UNP O67883
A	-20	LYS	-	EXPRESSION TAG	UNP O67883
A	-19	LYS	-	EXPRESSION TAG	UNP O67883
A	-18	ALA	-	EXPRESSION TAG	UNP O67883
A	-17	GLY	-	EXPRESSION TAG	UNP O67883
A	-16	SER	-	EXPRESSION TAG	UNP O67883
A	-15	ALA	_	EXPRESSION TAG	UNP O67883
A	-14	ALA	-	EXPRESSION TAG	UNP O67883
A	-13	ALA	-	EXPRESSION TAG	UNP O67883
A	-12	VAL	-	EXPRESSION TAG	UNP O67883
A	-11	LEU	-	EXPRESSION TAG	UNP O67883



Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	-10	GLU	-	EXPRESSION TAG	UNP 067883
A	-9	GLU	-	EXPRESSION TAG	UNP 067883
A	-8	ASN	-	EXPRESSION TAG	UNP 067883
A	-7	LEU	_	EXPRESSION TAG	UNP O67883
A	-6	TYR	-	EXPRESSION TAG	UNP O67883
A	-5	PHE	-	EXPRESSION TAG	UNP O67883
A	-4	GLN	-	EXPRESSION TAG	UNP O67883
A	-3	GLY	-	EXPRESSION TAG	UNP O67883
A	-2	SER	-	EXPRESSION TAG	UNP O67883
A	-1	PHE	-	EXPRESSION TAG	UNP O67883
A	0	THR	-	EXPRESSION TAG	UNP O67883
В	-35	MET	-	EXPRESSION TAG	UNP O67883
В	-34	GLY	-	EXPRESSION TAG	UNP O67883
В	-33	SER	-	EXPRESSION TAG	UNP O67883
В	-32	HIS	-	EXPRESSION TAG	UNP O67883
В	-31	HIS	-	EXPRESSION TAG	UNP O67883
В	-30	HIS	-	EXPRESSION TAG	UNP O67883
В	-29	HIS	-	EXPRESSION TAG	UNP O67883
В	-28	HIS	-	EXPRESSION TAG	UNP O67883
В	-27	HIS	-	EXPRESSION TAG	UNP O67883
В	-26	ASP	_	EXPRESSION TAG	UNP 067883
В	-25	ILE	-	EXPRESSION TAG	UNP O67883
В	-24	THR	-	EXPRESSION TAG	UNP O67883
В	-23	SER	-	EXPRESSION TAG	UNP O67883
В	-22	LEU	-	EXPRESSION TAG	UNP O67883
В	-21	TYR	-	EXPRESSION TAG	UNP O67883
В	-20	LYS	-	EXPRESSION TAG	UNP O67883
В	-19	LYS	-	EXPRESSION TAG	UNP 067883
В	-18	ALA	-	EXPRESSION TAG	UNP O67883
В	-17	GLY	-	EXPRESSION TAG	UNP O67883
В	-16	SER	-	EXPRESSION TAG	UNP O67883
В	-15	ALA	-	EXPRESSION TAG	UNP 067883
В	-14	ALA	-	EXPRESSION TAG	UNP O67883
В	-13	ALA		EXPRESSION TAG	UNP O67883
В	-12	VAL	-	EXPRESSION TAG	UNP O67883
В	-11	LEU	-	EXPRESSION TAG	UNP O67883
В	-10	GLU	-	EXPRESSION TAG	UNP O67883
В	-9	GLU	-	EXPRESSION TAG	UNP O67883
В	-8	ASN	-	EXPRESSION TAG	UNP O67883
В	-7	LEU	-	EXPRESSION TAG	UNP O67883
В	-6	TYR	-	EXPRESSION TAG	UNP O67883
В	-5	PHE	-	EXPRESSION TAG	UNP O67883




~ 1	e		
Continued	trom	nremous	naae
	110110	production	paycon

Chain	Residue	Modelled	Actual	Comment	Reference
В	-4	GLN	_	EXPRESSION TAG	UNP O67883
В	-3	GLY	-	EXPRESSION TAG	UNP O67883
В	-2	SER	_	EXPRESSION TAG	UNP O67883
В	-1	PHE	-	EXPRESSION TAG	UNP O67883
В	0	THR	-	EXPRESSION TAG	UNP O67883

• Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	В	5	Total Mg 5 5	0	0
2	A	5	Total Mg 5 5	0	0


 $\bullet \ \, \text{Molecule 3 is THIAMIN PHOSPHATE (three-letter code: TPS) (formula: $C_{12}H_{18}N_4O_4PS)$.}$



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf						
2	Λ	1	Total	С	N	О	Р	S	0	0					
)	3 A	1	22	12	4	4	1	1		U					
9	D	1	Total	С	N	О	Р	S	0	0					
3	В	D	В	Б	В	\mathbf{D}	22	12	4	4	1	1		0	

• Molecule 4 is PHOSPHOMETHYLPHOSPHONIC ACID ADENYLATE ESTER (three-letter code: ACP) (formula: $C_{11}H_{18}N_5O_{12}P_3$).





Mol	Chain	Residues	${f Atoms}$				ZeroOcc	AltConf		
4	Α	1	Total	С	N	О	Р	0	0	
4	4 A	1	31	11	5	12	3	U		
4	D	1	Total	С	N	О	Р	0	0	
4	D	1	31	11	5	12	3	U	U	

• Molecule 5 is water.


ľ	Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
	5	A	8	Total O 8 8	0	0
	5	В	6	Total O 6 6	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Thiamine monophosphate kinase





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	60.75Å 66.84Å 196.86Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	31.65 - 2.60	Depositor
resolution (A)	46.83 - 2.13	EDS
% Data completeness	86.8 (31.65-2.60)	Depositor
(in resolution range)	86.5 (46.83-2.13)	EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	0.00 (at 2.14Å)	Xtriage
Refinement program	CNS 1.1	Depositor
P. P.	0.213 , 0.270	Depositor
R, R_{free}	0.208 , 0.263	DCC
R_{free} test set	4406 reflections (10.15%)	wwPDB-VP
Wilson B-factor (Å ²)	50.1	Xtriage
Anisotropy	0.725	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.32, 37.5	EDS
L-test for twinning ²	$ < L >=0.49, < L^2>=0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	5011	wwPDB-VP
Average B, all atoms (Å ²)	56.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.93% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $< L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, ACP, TPS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles		
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	A	0.39	0/2481	0.64	1/3346 (0.0%)	
1	В	0.39	0/2499	0.62	0/3370	
All	All	0.39	0/4980	0.63	1/6716 (0.0%)	

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}(^{o})$
1	Α	288	GLY	N-CA-C	5.05	125.71	113.10

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2432	0	2448	115	0
1	В	2449	0	2462	104	0
2	A	5	0	0	0	0
2	В	5	0	0	0	0
3	A	22	0	16	0	0
3	В	22	0	16	0	0
4	A	31	0	14	0	0
4	В	31	0	14	2	0



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	A	8	0	0	0	0
5	В	6	0	0	0	0
All	All	5011	0	4970	202	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 20.

The worst 5 of 202 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$egin{aligned} ext{Interatomic} \ ext{distance} \ (ext{Å}) \end{aligned}$	Clash overlap (Å)
1:A:28:THR:HB	1:A:75:ASN:HD22	1.05	1.15
1:A:28:THR:HG21	1:A:39:LEU:HB3	1.43	1.00
1:A:142:ARG:HG3	1:A:222:ARG:HG2	1.43	0.96
1:B:22:LYS:HE2	1:B:22:LYS:H	1.30	0.93
1:A:219:LEU:HD22	1:A:227:ILE:HD11	1.48	0.93

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed Favoured Allowed		Outliers	Percentiles	
1	A	305/342~(89%)	284 (93%)	18 (6%)	3 (1%)	15 32
1	В	$307/342 \ (90\%)$	284 (92%)	17 (6%)	6 (2%)	7 14
All	All	$612/684 \ (90\%)$	568 (93%)	35 (6%)	9 (2%)	10 21

5 of 9 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	27	ASP
1	A	289	GLU



Continued from previous page...

Mol	Chain	Res	Type
1	В	24	ILE
1	В	297	LYS
1	В	299	GLU

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed Rotameric Outl		Outliers	Perce	\mathbf{ntiles}
1	A	263/291 (90%)	251 (95%)	12 (5%)	27	51
1	В	265/291 (91%)	253 (96%)	12 (4%)	27	52
All	All	528/582 (91%)	504 (96%)	24 (4%)	27	52

5 of 24 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	268	HIS
1	В	22	LYS
1	В	232	GLU
1	A	272	ARG
1	В	11	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 15 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	221	GLN
1	A	238	ASN
1	В	250	ASN
1	A	203	ASN
1	В	221	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.



5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 14 ligands modelled in this entry, 10 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Mol True Chair		ain Res Li		Link Bond lengths			Bond angles		
MIOI	Mol Type Chain	nes	Link	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2	
4	ACP	A	307	2	27,33,33	2.20	6 (22%)	32,52,52	2.47	15 (46%)
3	TPS	A	308	2	20,23,23	2.28	7 (35%)	25,33,33	1.76	3 (12%)
4	ACP	В	307	2	27,33,33	2.21	6 (22%)	32,52,52	2.48	14 (43%)
3	TPS	В	308	2	20,23,23	2.49	5 (25%)	25,33,33	1.89	2 (8%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	ACP	A	307	2	-	3/15/38/38	0/3/3/3
3	TPS	A	308	2	-	2/10/11/11	0/2/2/2
4	ACP	В	307	2	-	3/15/38/38	0/3/3/3
3	TPS	В	308	2	-	2/10/11/11	0/2/2/2

The worst 5 of 24 bond length outliers are listed below:

Mol	Chain	Res	Type	${f Atoms}$	\mathbf{Z}	${f Observed(\AA)}$	$\operatorname{Ideal}(ext{\AA})$
3	В	308	TPS	C6-C5	-7.77	1.47	1.50



$\alpha \cdots \tau$	r	•	
Continued	trom	nromanne	naae
\circ	110116	picolous	puyc

Mol	Chain	Res	Type	${f Atoms}$	\mathbf{Z}	$\operatorname{Observed}(\operatorname{\AA})$	$oxed{Ideal(\AA)}$
4	A	307	ACP	PB-O3A	7.57	1.66	1.58
4	В	307	ACP	PB-O3A	7.05	1.66	1.58
3	A	308	TPS	C6-C5	-6.34	1.48	1.50
3	В	308	TPS	C7A-N3	4.04	1.56	1.48

The worst 5 of 34 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^o)$	$Ideal(^{o})$
3	В	308	TPS	C6-C5-C4	7.19	133.21	127.43
4	В	307	ACP	O5'-C5'-C4'	7.04	133.21	108.99
3	A	308	TPS	C6-C5-C4	6.70	132.81	127.43
4	A	307	ACP	O5'-C5'-C4'	6.51	131.41	108.99
4	A	307	ACP	PA-O3A-PB	4.58	147.08	132.56

There are no chirality outliers.

5 of 10 torsion outliers are listed below:

Mol	Chain	${f Res}$	Type	Atoms
4	В	307	ACP	O4'-C4'-C5'-O5'
4	В	307	ACP	C3'-C4'-C5'-O5'
4	A	307	ACP	O4'-C4'-C5'-O5'
4	A	307	ACP	C3'-C4'-C5'-O5'
3	A	308	TPS	C4-C5-C6-C7


There are no ring outliers.


1 monomer is involved in 2 short contacts:


Mol	Chain	${f Res}$	Type	Clashes	Symm-Clashes
4	В	307	ACP	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.














5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	$egin{array}{c c} \mathbf{Analysed} & <& \mathbf{RSRZ}> \end{array}$		$\#\mathrm{RSRZ}{>}2$	$OWAB(A^2)$	Q<0.9
1	A	307/342~(89%)	-0.12	3 (0%) 82 80	35, 53, 77, 96	1 (0%)
1	В	309/342 (90%)	-0.09	5 (1%) 72 68	35, 55, 81, 97	1 (0%)
All	All	616/684 (90%)	-0.10	8 (1%) 77 73	35, 54, 80, 97	2 (0%)

The worst 5 of 8 RSRZ outliers are listed below:

Mol	Chain	${f Res}$	Type	RSRZ
1	В	300	PRO	3.3
1	В	299	GLU	3.3
1	A	302	GLY	2.4
1	В	298	VAL	2.3
1	В	172	GLU	2.2

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.


6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q < 0.9
2	MG	A	312	1/1	0.89	0.29	40,40,40,40	0
2	MG	В	313	1/1	0.94	0.28	27,27,27,27	0
2	MG	В	312	1/1	0.94	0.28	29,29,29,29	0
2	MG	В	314	1/1	0.94	0.26	35,35,35,35	0
4	ACP	A	307	31/31	0.95	0.18	44,54,64,65	0
4	ACP	В	307	31/31	0.95	0.18	39,49,64,65	0
2	MG	A	313	1/1	0.96	0.26	25,25,25,25	0
2	MG	В	310	1/1	0.96	0.26	30,30,30,30	0
2	MG	A	310	1/1	0.96	0.34	32,32,32,32	0
3	TPS	В	308	22/22	0.96	0.17	46,51,57,58	0
2	MG	A	309	1/1	0.96	0.29	25,25,25,25	0
2	MG	В	311	1/1	0.97	0.28	25,25,25,25	0
3	TPS	A	308	22/22	0.97	0.18	47,52,56,60	0
2	MG	A	314	1/1	0.99	0.23	25,25,25,25	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.











6.5 Other polymers (i)

There are no such residues in this entry.

