



# Full wwPDB X-ray Structure Validation Report ⓘ

Dec 18, 2023 – 03:04 am GMT

PDB ID : 4C9L  
Title : Structure of Cyanide and Camphor bound wild type CYP101D1  
Authors : Batabyal, D.; L Poulos, T.  
Deposited on : 2013-10-02  
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*



The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5950 (1.80-1.80)
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)
RSRZ outliers	127900	5850 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	421	
1	B	421	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	CYN	A	1420	-	-	X	-

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 14105 atoms, of which 6546 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CYTOCHROME P450.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	A	408	Total	C	H	N	O	S	0	0	0
			6471	2060	3231	577	590	13			
1	B	407	Total	C	H	N	O	S	0	0	0
			6454	2055	3223	575	588	13			

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula:  $C_{34}H_{32}FeN_4O_4$ ).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
			Total	C	Fe	H	N			O
2	A	1	Total	C	Fe	H	N	O	0	0
			73	34	1	30	4	4		
2	B	1	Total	C	Fe	H	N	O	0	0
			73	34	1	30	4	4		

- Molecule 3 is CAMPHOR (three-letter code: CAM) (formula:  $C_{10}H_{16}O$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	H	O		
3	A	1	27	10	16	1	0	0
3	B	1	27	10	16	1	0	0

- Molecule 4 is CYANIDE ION (three-letter code: CYN) (formula: CN).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	N		
4	A	1	2	1	1	0	0
4	B	1	2	1	1	0	0


- Molecule 5 is water.

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
5	A	515	Total 515	O 515	0	0
5	B	461	Total 461	O 461	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: CYTOCHROME P450



- Molecule 1: CYTOCHROME P450



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 64 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	151.53Å 151.53Å 196.04Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	46.30 – 1.80 46.30 – 1.80	Depositor EDS
% Data completeness (in resolution range)	99.6 (46.30-1.80) 97.3 (46.30-1.80)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.15 (at 1.81Å)	Xtrriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, $R_{free}$	0.152 , 0.182 0.154 , 0.183	Depositor DCC
$R_{free}$ test set	2000 reflections (1.64%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	18.4	Xtrriage
Anisotropy	0.335	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.42 , 49.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	14105	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	25.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.92% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, CYN, CAM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.50	0/3322	0.73	4/4514 (0.1%)
1	B	0.47	0/3313	0.68	2/4502 (0.0%)
All	All	0.48	0/6635	0.71	6/9016 (0.1%)

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	134	ARG	NE-CZ-NH2	-8.45	116.07	120.30
1	B	201	LEU	CA-CB-CG	5.83	128.72	115.30
1	A	55	LEU	CA-CB-CG	5.79	128.61	115.30
1	A	201	LEU	CA-CB-CG	5.38	127.68	115.30
1	A	220	ARG	NE-CZ-NH2	-5.34	117.63	120.30
1	B	338	ARG	NE-CZ-NH2	-5.28	117.66	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3240	3231	3215	17	0
1	B	3231	3223	3207	27	1
2	A	43	30	30	2	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	43	30	30	1	0
3	A	11	16	16	4	0
3	B	11	16	16	2	0
4	A	2	0	0	3	0
4	B	2	0	0	1	0
5	A	515	0	0	4	1
5	B	461	0	0	5	0
All	All	7559	6546	6514	50	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (50) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:225:THR:O	1:A:230:ARG:NH1	2.07	0.87
1:B:193:THR:HG22	1:B:196:GLU:H	1.54	0.73
1:A:111:THR:HG21	5:A:2238:HOH:O	1.94	0.66
1:B:226:ASP:OD1	1:B:229:THR:HG23	1.98	0.64
1:A:49:GLN:HB2	1:A:52:THR:HG22	1.79	0.63
1:A:230:ARG:NH2	5:A:2256:HOH:O	2.32	0.62
1:B:278:THR:HG21	5:B:2348:HOH:O	2.00	0.61
1:B:49:GLN:HB2	1:B:52:THR:HG22	1.84	0.59
1:B:314:GLU:HG3	1:B:319:MET:HE2	1.84	0.58
3:B:1419:CAM:H52	4:B:1420:CYN:C	2.34	0.58
1:A:352:ASP:OD1	1:A:354:THR:HG23	2.03	0.58
1:A:212:VAL:HG11	1:A:250:VAL:HB	1.86	0.57
1:A:341:ASP:OD2	1:B:338:ARG:NH2	2.39	0.56
3:A:1419:CAM:H52	4:A:1420:CYN:C	2.36	0.56
1:B:13:ARG:HG3	1:B:59:PRO:HB2	1.90	0.54
1:A:220:ARG:NH2	1:A:241:MET:O	2.37	0.53
1:B:219:ARG:HD3	1:B:229:THR:HG22	1.91	0.52
1:B:229:THR:HG21	5:B:2325:HOH:O	2.09	0.52
3:A:1419:CAM:H83	4:A:1420:CYN:N	2.25	0.52
2:A:1418:HEM:HBC2	2:A:1418:HEM:HHD	1.93	0.50
1:A:193:THR:HG22	1:A:195:GLU:H	1.77	0.49
1:B:103:THR:HG21	3:B:1419:CAM:H31	1.95	0.49
1:B:49:GLN:H	1:B:52:THR:CG2	2.26	0.49
1:B:12:HIS:HD2	5:B:2003:HOH:O	1.96	0.48
1:A:228:ILE:O	1:A:232:LEU:HB2	2.14	0.48
1:A:103:THR:HG21	3:A:1419:CAM:H31	1.94	0.47

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:212:VAL:HG12	1:A:216:ILE:CD1	2.45	0.47
1:B:314:GLU:CD	1:B:319:MET:HE1	2.35	0.46
5:A:2455:HOH:O	1:B:338:ARG:HD3	2.17	0.45
1:B:147:ARG:HG2	1:B:149:HIS:CE1	2.52	0.45
1:B:219:ARG:HB3	1:B:229:THR:HG22	1.98	0.45
1:A:111:THR:HG22	5:A:2233:HOH:O	2.17	0.45
1:B:57:TRP:CH2	1:B:59:PRO:HA	2.52	0.44
1:B:163:ARG:HD2	1:B:173:VAL:HG13	2.00	0.44
1:B:111:THR:HG22	5:B:2205:HOH:O	2.17	0.44
1:B:216:ILE:O	1:B:220:ARG:HG3	2.18	0.43
1:A:216:ILE:O	1:A:220:ARG:HG3	2.19	0.43
3:A:1419:CAM:H61	4:A:1420:CYN:N	2.35	0.42
1:A:308:TYR:CZ	1:A:323:GLY:HA2	2.55	0.42
1:B:228:ILE:O	1:B:232:LEU:HB2	2.19	0.42
2:B:1418:HEM:HMB1	2:B:1418:HEM:HBB2	2.01	0.42
1:B:275:HIS:HB3	1:B:278:THR:HG23	2.02	0.42
1:B:368:MET:HE2	1:B:372:ARG:CZ	2.49	0.42
1:B:163:ARG:HD2	1:B:173:VAL:CG1	2.49	0.42
1:B:314:GLU:CG	1:B:319:MET:HE2	2.50	0.41
2:A:1418:HEM:HMB1	2:A:1418:HEM:HBB2	2.01	0.41
1:B:308:TYR:CZ	1:B:323:GLY:HA2	2.55	0.41
1:B:390:ARG:NH1	5:B:2443:HOH:O	2.50	0.40
1:A:58:THR:O	1:A:63:GLY:HA2	2.22	0.40
1:A:368:MET:HE2	1:A:368:MET:HB2	1.87	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:351:ARG:NH2	5:A:2008:HOH:O[8_555]	2.17	0.03

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	406/421 (96%)	396 (98%)	10 (2%)	0	100	100
1	B	405/421 (96%)	395 (98%)	10 (2%)	0	100	100
All	All	811/842 (96%)	791 (98%)	20 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	346/356 (97%)	329 (95%)	17 (5%)	25	11
1	B	345/356 (97%)	330 (96%)	15 (4%)	29	14
All	All	691/712 (97%)	659 (95%)	32 (5%)	27	13

All (32) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	14	VAL
1	A	38	GLN
1	A	52	THR
1	A	55	LEU
1	A	86	ARG
1	A	89	TRP
1	A	111	THR
1	A	201	LEU
1	A	205	ASN
1	A	232	LEU
1	A	260	THR
1	A	277	GLU
1	A	287	LEU
1	A	345	THR
1	A	351	ARG
1	A	391	LEU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	393	ASP
1	B	52	THR
1	B	55	LEU
1	B	89	TRP
1	B	111	THR
1	B	193	THR
1	B	195	GLU
1	B	201	LEU
1	B	205	ASN
1	B	229	THR
1	B	232	LEU
1	B	239	LYS
1	B	260	THR
1	B	278	THR
1	B	345	THR
1	B	391	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	149	HIS
1	B	12	HIS
1	B	149	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
2	HEM	B	1418	1,4	41,50,50	1.67	5 (12%)	45,82,82	1.39	9 (20%)
3	CAM	A	1419	-	12,12,12	1.99	3 (25%)	20,21,21	1.13	2 (10%)
4	CYN	A	1420	2	0,1,1	-	-	-	-	-
4	CYN	B	1420	2	0,1,1	-	-	-	-	-
2	HEM	A	1418	1,4	41,50,50	1.66	4 (9%)	45,82,82	1.55	8 (17%)
3	CAM	B	1419	-	12,12,12	2.08	3 (25%)	20,21,21	1.40	4 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	CAM	A	1419	-	-	-	0/3/2/2
2	HEM	B	1418	1,4	-	2/12/54/54	-
2	HEM	A	1418	1,4	-	2/12/54/54	-
3	CAM	B	1419	-	-	-	0/3/2/2

All (15) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1418	HEM	C3D-C2D	6.25	1.50	1.36
2	A	1418	HEM	C3D-C2D	6.11	1.49	1.36
3	B	1419	CAM	O-C2	4.66	1.29	1.21
2	A	1418	HEM	C3C-C2C	-4.66	1.33	1.40
2	B	1418	HEM	C3C-C2C	-4.14	1.34	1.40
3	A	1419	CAM	O-C2	4.11	1.28	1.21
2	B	1418	HEM	C3C-CAC	3.09	1.54	1.47
2	A	1418	HEM	C3C-CAC	2.89	1.53	1.47
3	B	1419	CAM	C1-C2	-2.85	1.49	1.52
3	A	1419	CAM	C1-C2	-2.67	1.49	1.52
2	A	1418	HEM	CAB-C3B	2.42	1.54	1.47
2	B	1418	HEM	CAB-C3B	2.14	1.53	1.47

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1419	CAM	C6-C1	-2.06	1.49	1.55
3	B	1419	CAM	C3-C4	-2.05	1.48	1.53
2	B	1418	HEM	CMA-C3A	2.01	1.55	1.51

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1418	HEM	C4D-ND-C1D	4.54	109.76	105.07
3	B	1419	CAM	O-C2-C1	3.67	129.86	125.39
2	B	1418	HEM	C4C-CHD-C1D	3.38	127.02	122.56
2	A	1418	HEM	C4C-CHD-C1D	3.26	126.86	122.56
2	A	1418	HEM	C4B-CHC-C1C	3.13	126.69	122.56
2	A	1418	HEM	CHD-C1D-ND	2.98	127.67	124.43
2	B	1418	HEM	C4D-ND-C1D	2.90	108.07	105.07
2	B	1418	HEM	CMA-C3A-C4A	-2.74	124.25	128.46
2	A	1418	HEM	C1B-NB-C4B	2.71	107.88	105.07
3	A	1419	CAM	C5-C4-C3	-2.60	99.25	106.40
2	A	1418	HEM	CMD-C2D-C1D	2.44	128.75	125.04
2	B	1418	HEM	C3C-C4C-NC	-2.42	106.38	110.94
3	B	1419	CAM	C10-C1-C2	2.33	118.61	108.99
2	B	1418	HEM	C2C-C3C-C4C	2.32	108.52	106.90
3	B	1419	CAM	O-C2-C3	-2.26	121.77	126.47
2	B	1418	HEM	CMD-C2D-C1D	2.25	128.47	125.04
2	B	1418	HEM	CMC-C2C-C3C	2.22	128.83	124.68
3	A	1419	CAM	C6-C5-C4	2.10	106.43	103.15
2	B	1418	HEM	CHD-C1D-ND	2.09	126.71	124.43
3	B	1419	CAM	C6-C5-C4	2.08	106.40	103.15
2	A	1418	HEM	CMA-C3A-C4A	-2.08	125.27	128.46
2	A	1418	HEM	CMC-C2C-C3C	2.03	128.47	124.68
2	B	1418	HEM	C4A-C3A-C2A	2.01	108.39	107.00

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	1418	HEM	CAA-CBA-CGA-O1A
2	A	1418	HEM	CAA-CBA-CGA-O2A
2	B	1418	HEM	CAA-CBA-CGA-O2A
2	A	1418	HEM	CAA-CBA-CGA-O1A


There are no ring outliers.


6 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	1418	HEM	1	0
3	A	1419	CAM	4	0
4	A	1420	CYN	3	0
4	B	1420	CYN	1	0
2	A	1418	HEM	2	0
3	B	1419	CAM	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	408/421 (96%)	-0.62	7 (1%) 70 66	11, 18, 36, 56	0
1	B	407/421 (96%)	-0.55	5 (1%) 79 76	12, 21, 40, 65	0
All	All	815/842 (96%)	-0.59	12 (1%) 73 70	11, 20, 39, 65	0

All (12) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	239	LYS	4.1
1	B	417	GLN	3.9
1	B	238	GLY	3.2
1	A	239	LYS	3.0
1	A	237	ASP	2.6
1	B	237	ASP	2.6
1	A	221	GLY	2.3
1	A	223	SER	2.2
1	A	225	THR	2.1
1	A	238	GLY	2.1
1	A	230	ARG	2.0
1	B	284	ARG	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.


## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	CAM	A	1419	11/11	0.85	0.18	16,29,43,43	0
3	CAM	B	1419	11/11	0.92	0.09	19,24,35,35	0
4	CYN	B	1420	2/2	0.96	0.12	21,21,21,25	0
2	HEM	B	1418	43/43	0.99	0.11	11,15,24,29	0
4	CYN	A	1420	2/2	0.99	0.10	12,12,12,29	0
2	HEM	A	1418	43/43	0.99	0.08	9,13,24,29	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





## 6.5 Other polymers [i](#)

There are no such residues in this entry.