



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 11, 2021 – 02:51 AM EDT

PDB ID : 3CBW
Title : Crystal structure of the YdhT protein from *Bacillus subtilis*
Authors : Bonanno, J.B.; Rutter, M.; Bain, K.T.; Iizuka, M.; Romero, R.; Smith, D.; Wasserman, S.; Sauder, J.M.; Burley, S.K.; Almo, S.C.; New York SGX Research Center for Structural Genomics (NYSGXRC)
Deposited on : 2008-02-23
Resolution : 1.27 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : 1.13
EDS : 2.23.2
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.23.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.27 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1023 (1.28-1.24)
Clashscore	141614	1060 (1.28-1.24)
Ramachandran outliers	138981	1029 (1.28-1.24)
Sidechain outliers	138945	1028 (1.28-1.24)
RSRZ outliers	127900	1004 (1.28-1.24)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	353	 3% 89% 6% 5%
1	B	353	 5% 90% 7%

2 Entry composition i

There are 3 unique types of molecules in this entry. The entry contains 6012 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called YdhT protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	336	2702	1728	448	518	8	0	5	0
1	B	330	2679	1716	448	506	9	0	7	0

There are 28 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	18	MET	-	expression tag	UNP O05512
A	19	SER	-	expression tag	UNP O05512
A	20	LEU	-	expression tag	UNP O05512
A	153	LYS	ASN	engineered mutation	UNP O05512
A	158	SER	ALA	engineered mutation	UNP O05512
A	160	VAL	ALA	engineered mutation	UNP O05512
A	363	GLU	-	expression tag	UNP O05512
A	364	GLY	-	expression tag	UNP O05512
A	365	HIS	-	expression tag	UNP O05512
A	366	HIS	-	expression tag	UNP O05512
A	367	HIS	-	expression tag	UNP O05512
A	368	HIS	-	expression tag	UNP O05512
A	369	HIS	-	expression tag	UNP O05512
A	370	HIS	-	expression tag	UNP O05512
B	18	MET	-	expression tag	UNP O05512
B	19	SER	-	expression tag	UNP O05512
B	20	LEU	-	expression tag	UNP O05512
B	153	LYS	ASN	engineered mutation	UNP O05512
B	158	SER	ALA	engineered mutation	UNP O05512
B	160	VAL	ALA	engineered mutation	UNP O05512
B	363	GLU	-	expression tag	UNP O05512
B	364	GLY	-	expression tag	UNP O05512
B	365	HIS	-	expression tag	UNP O05512
B	366	HIS	-	expression tag	UNP O05512
B	367	HIS	-	expression tag	UNP O05512

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	368	HIS	-	expression tag	UNP O05512
B	369	HIS	-	expression tag	UNP O05512
B	370	HIS	-	expression tag	UNP O05512

- Molecule 2 is CITRIC ACID (three-letter code: CIT) (formula: C₆H₈O₇).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	1	Total C O 13 6 7	0	0
2	B	1	Total C O 13 6 7	0	0
2	B	1	Total C O 13 6 7	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	319	Total O 319 319	0	0
3	B	273	Total O 273 273	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: YdhT protein



- Molecule 1: YdhT protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	57.73Å 69.62Å 72.50Å 90.00° 91.09° 90.00°	Depositor
Resolution (Å)	20.00 – 1.27 16.02 – 1.27	Depositor EDS
% Data completeness (in resolution range)	96.3 (20.00-1.27) 96.2 (16.02-1.27)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	0.10	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.52 (at 1.27Å)	Xtrriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.149 , 0.172 0.153 , 0.176	Depositor DCC
R_{free} test set	7332 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	9.5	Xtrriage
Anisotropy	0.261	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.44 , 50.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.000 for -h,l,k 0.014 for -h,-l,-k 0.018 for h,-k,-l	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	6012	wwPDB-VP
Average B, all atoms (Å ²)	14.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 38.44 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 3.6861e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CIT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.43	0/2796	0.60	0/3809
1	B	0.42	0/2777	0.59	0/3774
All	All	0.42	0/5573	0.59	0/7583

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2702	0	2564	12	0
1	B	2679	0	2552	9	0
2	B	39	0	15	2	0
3	A	319	0	0	4	0
3	B	273	0	0	2	0
All	All	6012	0	5131	21	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (21) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:308:ILE:HB	1:B:339:LEU:HD12	1.63	0.79
1:A:41:LYS:HE2	3:A:649:HOH:O	1.82	0.78
1:B:304:TYR:CD1	1:B:339:LEU:HD22	2.25	0.71
1:B:304:TYR:HD1	1:B:339:LEU:HD22	1.58	0.69
1:B:321:PHE:CZ	1:B:339:LEU:HD21	2.40	0.57
1:A:223:LYS:HE2	3:A:425:HOH:O	2.10	0.52
1:A:72:SER:HB2	3:A:637:HOH:O	2.09	0.51
1:A:80:ARG:NH1	3:A:513:HOH:O	2.43	0.51
1:B:325:ASN:HB2	3:B:487:HOH:O	2.11	0.50
1:A:225:TYR:HA	1:A:239:TRP:CH2	2.49	0.48
1:B:368:HIS:CD2	2:B:2:CIT:O1	2.69	0.46
1:A:117:MET:CG	1:A:179:LEU:HD23	2.45	0.46
1:B:325:ASN:CB	3:B:487:HOH:O	2.65	0.45
1:A:222:LYS:HE2	1:A:259:TYR:CE2	2.53	0.44
1:B:225:TYR:HA	1:B:239:TRP:CH2	2.53	0.43
1:A:269:PHE:CE2	1:A:293:VAL:HG23	2.54	0.43
1:A:117:MET:HG3	1:A:179:LEU:CD2	2.49	0.42
1:A:302:PHE:O	1:A:334:LYS:HG2	2.21	0.41
1:A:108:ILE:HB	1:A:171:LYS:HD3	2.03	0.40
1:B:368:HIS:HD2	2:B:2:CIT:O1	2.04	0.40
1:A:117:MET:HG3	1:A:179:LEU:HD23	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	339/353 (96%)	332 (98%)	6 (2%)	1 (0%)	41	16
1	B	331/353 (94%)	327 (99%)	3 (1%)	1 (0%)	41	16
All	All	670/706 (95%)	659 (98%)	9 (1%)	2 (0%)	41	16

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	69	ASP
1	B	69	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	289/300 (96%)	285 (99%)	4 (1%)	67 30
1	B	287/300 (96%)	283 (99%)	4 (1%)	67 30
All	All	576/600 (96%)	568 (99%)	8 (1%)	67 30

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	56	ASN
1	A	96	ARG
1	A	127	GLN
1	A	142	PHE
1	B	96	ARG
1	B	127	GLN
1	B	142	PHE
1	B	339	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	56	ASN
1	A	182	GLN
1	B	85	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	CIT	B	1	-	3,12,12	1.34	0	3,17,17	1.40	1 (33%)
2	CIT	B	2	-	3,12,12	1.21	0	3,17,17	1.60	1 (33%)
2	CIT	B	3	-	3,12,12	1.40	0	3,17,17	0.79	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	CIT	B	1	-	-	0/6/16/16	-
2	CIT	B	2	-	-	1/6/16/16	-
2	CIT	B	3	-	-	0/6/16/16	-

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	2	CIT	C3-C2-C1	-2.48	111.01	114.98
2	B	1	CIT	C3-C4-C5	-2.41	111.13	114.98

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	2	CIT	C6-C3-C4-C5

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	2	CIT	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	336/353 (95%)	-0.07	9 (2%) 54 45	7, 11, 19, 30	0
1	B	330/353 (93%)	0.08	18 (5%) 25 19	8, 12, 23, 31	1 (0%)
All	All	666/706 (94%)	0.01	27 (4%) 37 30	7, 12, 21, 31	1 (0%)

All (27) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	302	PHE	7.1
1	A	298	ALA	6.6
1	B	365	HIS	6.4
1	B	366	HIS	5.7
1	B	364	GLY	5.4
1	B	304	TYR	4.9
1	B	367	HIS	4.7
1	B	363	GLU	4.2
1	A	301	SER	3.7
1	A	297	THR	3.6
1	A	299	ASN	3.4
1	B	334	LYS	3.2
1	B	353	ASN	3.1
1	B	270	GLN	3.0
1	A	296	GLN	2.8
1	A	353	ASN	2.8
1	A	352	TRP	2.6
1	A	80	ARG	2.6
1	B	324	TRP	2.6
1	B	355	ASP	2.5
1	B	325	ASN	2.5
1	B	368	HIS	2.4
1	A	300	GLY	2.3
1	B	274[A]	SER	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	102	ALA	2.2
1	B	208	GLN	2.2
1	B	339	LEU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	CIT	B	3	13/13	0.73	0.26	21,31,34,34	0
2	CIT	B	2	13/13	0.82	0.21	20,28,34,34	0
2	CIT	B	1	13/13	0.85	0.13	16,17,18,19	0

6.5 Other polymers [i](#)

There are no such residues in this entry.