



wwPDB X-ray Structure Validation Summary Report ⓘ

May 23, 2020 – 10:16 am BST

PDB ID : 4CBT
Title : Design, synthesis, and biological evaluation of potent and selective Class IIa HDAC inhibitors as a potential therapy for Huntington's disease
Authors : Burli, R.W.; Luckhurst, C.A.; Aziz, O.; Matthews, K.L.; Yates, D.; Lyons, K.A.; Beconi, M.; McAllister, G.; Breccia, P.; Stott, A.J.; Penrose, S.D.; Wall, M.; Lamers, M.; Leonard, P.; Mueller, I.; Richardson, C.M.; Jarvis, R.; Stones, L.; Hughes, S.; Wishart, G.; Haughan, A.F.; O'Connell, C.; Mead, T.; McNeil, H.; Vann, J.; Mangette, J.; Maillard, M.; Beaumont, V.; Munoz-Sanjuan, I.; Dominguez, C.
Deposited on : 2013-10-16
Resolution : 3.03 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.11
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.03 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2752 (3.08-3.00)
Clashscore	141614	3096 (3.08-3.00)
Ramachandran outliers	138981	2986 (3.08-3.00)
Sidechain outliers	138945	2988 (3.08-3.00)
RSRZ outliers	127900	2636 (3.08-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	395	77% 12% 12%
1	B	395	79% 10% • 11%
1	C	395	75% 12% • 12%

2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 7993 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HISTONE DEACETYLASE 4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	349	Total 2638	C 1660	N 465	O 494	S 19	0	0	0
1	B	352	Total 2645	C 1666	N 461	O 499	S 19	0	0	0
1	C	346	Total 2616	C 1650	N 462	O 485	S 19	0	0	0

There are 27 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	645	MET	-	expression tag	UNP P56524
A	646	GLY	-	expression tag	UNP P56524
A	647	SER	-	expression tag	UNP P56524
A	1034	HIS	-	expression tag	UNP P56524
A	1035	HIS	-	expression tag	UNP P56524
A	1036	HIS	-	expression tag	UNP P56524
A	1037	HIS	-	expression tag	UNP P56524
A	1038	HIS	-	expression tag	UNP P56524
A	1039	HIS	-	expression tag	UNP P56524
B	645	MET	-	expression tag	UNP P56524
B	646	GLY	-	expression tag	UNP P56524
B	647	SER	-	expression tag	UNP P56524
B	1035	HIS	-	expression tag	UNP P56524
B	1036	HIS	-	expression tag	UNP P56524
B	1037	HIS	-	expression tag	UNP P56524
B	1038	HIS	-	expression tag	UNP P56524
B	1039	HIS	-	expression tag	UNP P56524
B	1040	HIS	-	expression tag	UNP P56524
C	645	MET	-	expression tag	UNP P56524
C	646	GLY	-	expression tag	UNP P56524
C	647	SER	-	expression tag	UNP P56524
C	1034	HIS	-	expression tag	UNP P56524
C	1035	HIS	-	expression tag	UNP P56524

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	1036	HIS	-	expression tag	UNP P56524
C	1037	HIS	-	expression tag	UNP P56524
C	1038	HIS	-	expression tag	UNP P56524
C	1039	HIS	-	expression tag	UNP P56524

- Molecule 2 is (1R,2R,3R)-2-[4-(5-fluoranylpyrimidin-2-yl)phenyl]-N-oxidanyl-3-phenyl-cyclopropane-1-carboxamide (three-letter code: 9F4) (formula: C₂₀H₁₆FN₃O₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	F	N			O
2	A	1	Total	C	F	N	O	0	0
			26	20	1	3	2		
2	B	1	Total	C	F	N	O	0	0
			26	20	1	3	2		
2	C	1	Total	C	F	N	O	0	0
			26	20	1	3	2		

- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Zn		
3	B	2	Total	Zn	0	0
			2	2		
3	A	2	Total	Zn	0	0
			2	2		
3	C	2	Total	Zn	0	0
			2	2		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	5	Total O 5 5	0	0
4	B	4	Total O 4 4	0	0
4	C	1	Total O 1 1	0	0

Y927	D934	I968	D977	A980	I981	C982	L993	G994	I995	E996	LEU	ASP	PRO	LEU	PRO	GLU	K1003	V1020	I1023	Y1027	H1034	HIS	HIS	HIS	HIS	HIS
------	------	------	------	------	------	------	------	------	------	------	-----	-----	-----	-----	-----	-----	-------	-------	-------	-------	-------	-----	-----	-----	-----	-----

4 Data and refinement statistics

Property	Value	Source
Space group	P 32	Depositor
Cell constants a, b, c, α , β , γ	104.41Å 104.41Å 88.44Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	90.42 – 3.03 90.42 – 3.03	Depositor EDS
% Data completeness (in resolution range)	100.0 (90.42-3.03) 99.8 (90.42-3.03)	Depositor EDS
R_{merge}	0.16	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.51 (at 3.01Å)	Xtriage
Refinement program	REFMAC 5.5.0110	Depositor
R, R_{free}	0.211 , 0.273 0.211 , 0.266	Depositor DCC
R_{free} test set	1070 reflections (5.13%)	wwPDB-VP
Wilson B-factor (Å ²)	47.8	Xtriage
Anisotropy	0.016	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 24.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.011 for -h,-k,l 0.049 for h,-h-k,-l 0.032 for -k,-h,-l	Xtriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	7993	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 22.14 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 6.0533e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 9F4, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.48	0/2699	0.53	1/3661 (0.0%)
1	B	0.43	0/2707	0.53	2/3679 (0.1%)
1	C	0.52	0/2675	0.54	2/3628 (0.1%)
All	All	0.48	0/8081	0.53	5/10968 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	1

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	681	ARG	NE-CZ-NH1	7.10	123.85	120.30
1	C	681	ARG	NE-CZ-NH1	6.49	123.55	120.30
1	A	681	ARG	NE-CZ-NH1	6.16	123.38	120.30
1	B	681	ARG	NE-CZ-NH2	-5.92	117.34	120.30
1	C	681	ARG	NE-CZ-NH2	-5.30	117.65	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	899	ASP	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2638	0	2554	25	0
1	B	2645	0	2550	22	0
1	C	2616	0	2537	26	0
2	A	26	0	12	0	0
2	B	26	0	13	0	0
2	C	26	0	12	0	0
3	A	2	0	0	0	0
3	B	2	0	0	0	0
3	C	2	0	0	0	0
4	A	5	0	0	0	0
4	B	4	0	0	0	0
4	C	1	0	0	0	0
All	All	7993	0	7678	73	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 73 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:785:VAL:HG21	1:B:969:VAL:HG23	1.51	0.91
1:B:785:VAL:HG21	1:B:969:VAL:CG2	2.02	0.88
1:A:825:LEU:HD22	1:A:831:VAL:HG21	1.68	0.73
1:B:916:MET:HE1	1:B:960:LEU:HD23	1.74	0.69
1:B:682:ILE:HD13	1:B:798:ARG:HG2	1.76	0.66

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	343/395 (87%)	318 (93%)	23 (7%)	2 (1%)	25	60
1	B	346/395 (88%)	325 (94%)	18 (5%)	3 (1%)	17	52
1	C	336/395 (85%)	306 (91%)	28 (8%)	2 (1%)	25	60
All	All	1025/1185 (86%)	949 (93%)	69 (7%)	7 (1%)	22	57

5 of 7 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	652	PHE
1	C	900	PRO
1	A	942	PRO
1	B	994	GLY
1	A	941	THR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	279/322 (87%)	273 (98%)	6 (2%)	52	79
1	B	280/322 (87%)	274 (98%)	6 (2%)	53	80
1	C	275/322 (85%)	270 (98%)	5 (2%)	59	83
All	All	834/966 (86%)	817 (98%)	17 (2%)	55	81

5 of 17 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	802	HIS
1	B	839	TRP
1	C	802	HIS
1	B	726	ASN
1	C	808	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 14 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	869	ASN
1	B	947	ASN
1	C	863	HIS
1	B	843	HIS
1	C	843	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 6 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	9F4	A	2035	3	29,29,29	0.91	2 (6%)	39,41,41	1.03	4 (10%)
2	9F4	B	2034	3	29,29,29	0.85	2 (6%)	39,41,41	0.88	1 (2%)
2	9F4	C	2035	3	29,29,29	0.87	2 (6%)	39,41,41	0.94	3 (7%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	9F4	A	2035	3	-	6/18/27/27	0/4/4/4
2	9F4	B	2034	3	-	1/18/27/27	0/4/4/4
2	9F4	C	2035	3	-	2/18/27/27	0/4/4/4

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	2035	9F4	C16-N18	2.53	1.37	1.33
2	C	2035	9F4	C15-C16	-2.44	1.47	1.51
2	A	2035	9F4	C15-C16	-2.37	1.47	1.51
2	C	2035	9F4	C16-N18	2.27	1.36	1.33
2	B	2034	9F4	C15-C16	-2.23	1.47	1.51

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	2034	9F4	C23-C22-N21	3.20	123.38	121.46
2	A	2035	9F4	C6-C7-C8	-3.08	114.14	121.82
2	A	2035	9F4	C23-C22-N21	2.86	123.18	121.46
2	C	2035	9F4	C23-C22-N21	2.84	123.17	121.46
2	C	2035	9F4	C23-C24-N25	2.53	122.98	121.46

There are no chirality outliers.

5 of 9 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	B	2034	9F4	C7-C15-C16-O17
2	C	2035	9F4	C7-C15-C16-N18
2	C	2035	9F4	C7-C15-C16-O17
2	A	2035	9F4	C7-C15-C16-N18
2	A	2035	9F4	C7-C15-C16-O17



There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier.

Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	349/395 (88%)	-0.38	1 (0%) 94 83	23, 36, 50, 62	0
1	B	352/395 (89%)	-0.51	0 100 100	24, 34, 46, 48	0
1	C	346/395 (87%)	-0.30	0 100 100	30, 43, 61, 70	0
All	All	1047/1185 (88%)	-0.40	1 (0%) 95 89	23, 38, 54, 70	0

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	726	ASN	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	9F4	C	2035	26/26	0.95	0.23	33,35,38,38	0
3	ZN	B	2036	1/1	0.95	0.10	38,38,38,38	0


Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	9F4	A	2035	26/26	0.96	0.27	30,31,33,34	0
2	9F4	B	2034	26/26	0.97	0.20	27,28,33,34	0
3	ZN	C	2036	1/1	0.97	0.11	33,33,33,33	0
3	ZN	B	2035	1/1	0.97	0.11	25,25,25,25	0
3	ZN	C	2037	1/1	0.97	0.07	45,45,45,45	0
3	ZN	A	2037	1/1	0.98	0.07	40,40,40,40	0
3	ZN	A	2036	1/1	0.99	0.07	28,28,28,28	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers

There are no such residues in this entry.