



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 11, 2021 – 12:23 AM EDT

PDB ID : 3CDB
Title : Thermodynamic and structure guided design of statin hmg-coa reductase inhibitors
Authors : Pavlovsky, A.; Sarver, R.W.; Harris, M.S.; Finzel, B.C.
Deposited on : 2008-02-26
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.23.2
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.23.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	441	 7% 88% 6% • 5%
1	B	441	 2% 90% 5% 5%
1	C	441	 7% 89% 5% 6%
1	D	441	 6% 86% • 10%

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 13109 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 3-hydroxy-3-methylglutaryl-coenzyme A reductase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	421	3133	1951	551	601	30	0	0	0
1	B	421	3139	1954	552	602	31	0	2	0
1	C	415	3086	1923	541	591	31	0	1	0
1	D	396	2934	1826	518	561	29	0	0	0

There are 28 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	435	HIS	-	expression tag	UNP P04035
A	436	HIS	-	expression tag	UNP P04035
A	437	HIS	-	expression tag	UNP P04035
A	438	HIS	-	expression tag	UNP P04035
A	439	HIS	-	expression tag	UNP P04035
A	440	HIS	-	expression tag	UNP P04035
A	485	ILE	MET	engineered mutation	UNP P04035
B	435	HIS	-	expression tag	UNP P04035
B	436	HIS	-	expression tag	UNP P04035
B	437	HIS	-	expression tag	UNP P04035
B	438	HIS	-	expression tag	UNP P04035
B	439	HIS	-	expression tag	UNP P04035
B	440	HIS	-	expression tag	UNP P04035
B	485	ILE	MET	engineered mutation	UNP P04035
C	435	HIS	-	expression tag	UNP P04035
C	436	HIS	-	expression tag	UNP P04035
C	437	HIS	-	expression tag	UNP P04035
C	438	HIS	-	expression tag	UNP P04035
C	439	HIS	-	expression tag	UNP P04035
C	440	HIS	-	expression tag	UNP P04035
C	485	ILE	MET	engineered mutation	UNP P04035

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	435	HIS	-	expression tag	UNP P04035
D	436	HIS	-	expression tag	UNP P04035
D	437	HIS	-	expression tag	UNP P04035
D	438	HIS	-	expression tag	UNP P04035
D	439	HIS	-	expression tag	UNP P04035
D	440	HIS	-	expression tag	UNP P04035
D	485	ILE	MET	engineered mutation	UNP P04035

- Molecule 2 is (3R,5R)-7-{3-[(4-carbamoylphenyl)sulfamoyl]-4,5-bis(4-fluorophenyl)-2-(1-methylethylethyl)-1H-pyrrol-1-yl}-3,5-dihydroxyheptanoic acid (three-letter code: 9HI) (formula: C₃₃H₃₅F₂N₃O₇S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
			Total	C	F	N	O			S
2	A	1	Total	C	F	N	O	S	0	0
			46	33	2	3	7	1		
2	A	1	Total	C	F	N	O	S	0	0
			46	33	2	3	7	1		
2	C	1	Total	C	F	N	O	S	0	0
			46	33	2	3	7	1		
2	D	1	Total	C	F	N	O	S	0	0
			46	33	2	3	7	1		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	180	Total	O	0	0
			180	180		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	159	Total 159	O 159	0	0
3	C	122	Total 122	O 122	0	0
3	D	172	Total 172	O 172	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: 3-hydroxy-3-methylglutaryl-coenzyme A reductase




- Molecule 1: 3-hydroxy-3-methylglutaryl-coenzyme A reductase




- Molecule 1: 3-hydroxy-3-methylglutaryl-coenzyme A reductase



- Molecule 1: 3-hydroxy-3-methylglutaryl-coenzyme A reductase





4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	73.58Å 173.48Å 75.99Å 90.00° 118.68° 90.00°	Depositor
Resolution (Å)	50.00 – 2.30 43.68 – 2.29	Depositor EDS
% Data completeness (in resolution range)	92.9 (50.00-2.30) 92.4 (43.68-2.29)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.38 (at 2.29Å)	Xtrriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.202 , 0.239 0.204 , 0.238	Depositor DCC
R_{free} test set	3509 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å ²)	31.5	Xtrriage
Anisotropy	0.382	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 35.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.000 for -h-l,k,h 0.000 for l,k,-h-l 0.019 for h,-k,-h-l 0.021 for -h-l,-k,l 0.021 for l,-k,h	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	13109	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.26% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 9HI

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.33	0/3179	0.62	6/4298 (0.1%)
1	B	0.33	0/3195	0.62	5/4319 (0.1%)
1	C	0.32	0/3135	0.60	6/4236 (0.1%)
1	D	0.33	0/2975	0.61	3/4019 (0.1%)
All	All	0.33	0/12484	0.61	20/16872 (0.1%)

There are no bond length outliers.

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	623	ASP	CB-CG-OD2	6.47	124.13	118.30
1	A	623	ASP	CB-CG-OD2	5.89	123.60	118.30
1	A	547	ASP	CB-CG-OD2	5.74	123.46	118.30
1	C	547	ASP	CB-CG-OD2	5.72	123.45	118.30
1	D	547	ASP	CB-CG-OD2	5.71	123.44	118.30
1	C	516	ASP	CB-CG-OD2	5.70	123.42	118.30
1	A	516	ASP	CB-CG-OD2	5.61	123.35	118.30
1	B	547	ASP	CB-CG-OD2	5.55	123.29	118.30
1	C	790	ASP	CB-CG-OD2	5.49	123.24	118.30
1	D	653	ASP	CB-CG-OD2	5.39	123.15	118.30
1	D	790	ASP	CB-CG-OD2	5.39	123.15	118.30
1	B	516	ASP	CB-CG-OD2	5.35	123.12	118.30
1	B	829	ASP	CB-CG-OD2	5.34	123.10	118.30
1	C	653	ASP	CB-CG-OD2	5.29	123.06	118.30
1	C	623	ASP	CB-CG-OD2	5.28	123.05	118.30
1	B	601	ASP	CB-CG-OD2	5.26	123.03	118.30
1	C	829	ASP	CB-CG-OD2	5.23	123.01	118.30
1	A	653	ASP	CB-CG-OD2	5.15	122.94	118.30
1	A	767	ASP	CB-CG-OD2	5.08	122.87	118.30
1	A	690	ASP	CB-CG-OD2	5.02	122.82	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3133	0	3167	13	0
1	B	3139	0	3174	6	0
1	C	3086	0	3122	7	0
1	D	2934	0	2967	3	0
2	A	92	0	68	8	0
2	C	46	0	34	3	0
2	D	46	0	34	3	0
3	A	180	0	0	0	0
3	B	159	0	0	1	0
3	C	122	0	0	0	0
3	D	172	0	0	0	0
All	All	13109	0	12566	38	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (38) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:2:9HI:H13B	2:A:2:9HI:H7	1.71	0.70
1:A:655:MET:SD	1:A:657:MET:HG3	2.32	0.69
2:D:3:9HI:H13B	2:D:3:9HI:H7	1.76	0.67
2:A:1:9HI:H7	2:A:1:9HI:H13B	1.77	0.66
2:C:4:9HI:H13B	2:C:4:9HI:H7	1.79	0.64
1:C:681:LEU:HD22	1:D:731:VAL:HG22	1.79	0.64
1:C:771:ASN:OD1	1:C:775:SER:OG	2.15	0.63
2:A:1:9HI:C2	2:A:1:9HI:H26	2.31	0.60
2:A:1:9HI:H26	2:A:1:9HI:C12	2.32	0.59
1:A:861:HIS:HA	2:A:1:9HI:F2	1.96	0.55
2:D:3:9HI:H8A	2:D:3:9HI:H14B	1.93	0.51
2:C:4:9HI:O1	2:C:4:9HI:H6	2.10	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:655:MET:SD	1:B:657[B]:MET:HG3	2.51	0.51
1:A:441:GLU:N	1:A:442:PRO:CD	2.73	0.50
2:D:3:9HI:O1	2:D:3:9HI:H6	2.12	0.50
1:A:472:ASN:O	1:A:473:ALA:HB3	2.13	0.49
1:B:555:MET:HE3	1:B:563:VAL:HG22	1.94	0.49
2:A:2:9HI:O1	2:A:2:9HI:H6	2.14	0.48
1:C:657[B]:MET:HG3	1:C:658:ASN:N	2.28	0.48
1:A:672:HIS:CD2	1:A:676:PRO:HA	2.48	0.48
1:A:472:ASN:O	1:A:473:ALA:CB	2.62	0.48
1:A:638:ILE:O	1:D:796:THR:HG21	2.13	0.48
1:B:638:ILE:O	1:C:796:THR:HG21	2.14	0.47
1:A:474:LYS:O	1:A:475:HIS:HB2	2.15	0.46
1:A:477:PRO:O	1:A:478:ALA:HB2	2.17	0.45
1:A:596:LEU:HB3	1:A:601:ASP:HB2	1.99	0.45
1:B:590:ARG:HD2	3:B:984:HOH:O	2.17	0.45
1:B:702:ARG:O	1:B:799:SER:HA	2.17	0.45
1:A:468:ILE:HG12	1:A:498:LEU:CD1	2.48	0.44
1:C:702:ARG:O	1:C:799:SER:HA	2.17	0.43
1:A:771:ASN:OD1	1:A:775:SER:OG	2.36	0.43
1:B:796:THR:HG21	1:C:638:ILE:O	2.19	0.42
2:A:1:9HI:H14B	2:A:1:9HI:H8A	2.01	0.42
1:A:474:LYS:O	1:A:475:HIS:CB	2.67	0.42
2:C:4:9HI:H14B	2:C:4:9HI:H8A	2.02	0.41
1:D:468:ILE:HG21	1:D:501:LYS:NZ	2.36	0.41
2:A:1:9HI:C2	2:A:1:9HI:C26	2.99	0.41
1:C:655:MET:SD	1:C:657[A]:MET:HG3	2.61	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	419/441 (95%)	395 (94%)	20 (5%)	4 (1%)	15	17
1	B	421/441 (96%)	409 (97%)	12 (3%)	0	100	100
1	C	412/441 (93%)	396 (96%)	16 (4%)	0	100	100
1	D	392/441 (89%)	378 (96%)	13 (3%)	1 (0%)	41	50
All	All	1644/1764 (93%)	1578 (96%)	61 (4%)	5 (0%)	41	50

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	473	ALA
1	A	475	HIS
1	A	478	ALA
1	D	514	TYR
1	A	477	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	335/355 (94%)	323 (96%)	12 (4%)	35	49
1	B	337/355 (95%)	322 (96%)	15 (4%)	27	39
1	C	331/355 (93%)	320 (97%)	11 (3%)	38	53
1	D	313/355 (88%)	303 (97%)	10 (3%)	39	54
All	All	1316/1420 (93%)	1268 (96%)	48 (4%)	37	49

All (48) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	441	GLU
1	A	443	ARG
1	A	463	SER
1	A	482	GLU
1	A	486	GLU
1	A	498	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	512	LEU
1	A	630	ARG
1	A	649	SER
1	A	704	LYS
1	A	814	GLN
1	A	861	HIS
1	B	450	GLN
1	B	452	LEU
1	B	470	LEU
1	B	482	GLU
1	B	486	GLU
1	B	505	GLU
1	B	523	MET
1	B	634	LEU
1	B	657[A]	MET
1	B	657[B]	MET
1	B	752	HIS
1	B	771[A]	ASN
1	B	771[B]	ASN
1	B	828	LYS
1	B	829	ASP
1	C	450	GLN
1	C	470	LEU
1	C	505	GLU
1	C	523	MET
1	C	612	SER
1	C	649	SER
1	C	657[A]	MET
1	C	657[B]	MET
1	C	666	LYS
1	C	681	LEU
1	C	851	LEU
1	D	484	LEU
1	D	486	GLU
1	D	487	THR
1	D	657	MET
1	D	660	ILE
1	D	666	LYS
1	D	669	SER
1	D	791	LEU
1	D	828	LYS
1	D	851	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (17) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	472	ASN
1	A	510	GLN
1	A	518	ASN
1	A	567	ASN
1	A	632	GLN
1	A	819	GLN
1	B	450	GLN
1	B	472	ASN
1	B	529	ASN
1	C	469	GLN
1	C	472	ASN
1	C	679	GLN
1	D	472	ASN
1	D	518	ASN
1	D	632	GLN
1	D	642	ASN
1	D	861	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	9HI	C	4	-	42,49,49	1.64	3 (7%)	58,71,71	1.80	10 (17%)
2	9HI	A	1	-	42,49,49	1.56	2 (4%)	58,71,71	2.07	8 (13%)
2	9HI	D	3	-	42,49,49	1.58	3 (7%)	58,71,71	1.80	10 (17%)
2	9HI	A	2	-	42,49,49	1.60	3 (7%)	58,71,71	1.90	12 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	9HI	C	4	-	-	9/32/40/40	0/4/4/4
2	9HI	A	1	-	-	7/32/40/40	0/4/4/4
2	9HI	D	3	-	-	13/32/40/40	0/4/4/4
2	9HI	A	2	-	-	12/32/40/40	0/4/4/4

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	4	9HI	C2-S1	-8.68	1.65	1.79
2	A	1	9HI	C2-S1	-8.43	1.65	1.79
2	A	2	9HI	C2-S1	-8.38	1.66	1.79
2	D	3	9HI	C2-S1	-8.14	1.66	1.79
2	C	4	9HI	C32-N2	-4.03	1.36	1.43
2	D	3	9HI	C32-N2	-4.01	1.36	1.43
2	A	2	9HI	C32-N2	-3.81	1.37	1.43
2	A	1	9HI	C32-N2	-3.49	1.37	1.43
2	D	3	9HI	C1-C5	2.32	1.46	1.40
2	A	2	9HI	C1-C5	2.23	1.46	1.40
2	C	4	9HI	C1-C5	2.12	1.46	1.40

All (40) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1	9HI	O2-S1-O1	-10.93	106.11	119.55
2	D	3	9HI	O2-S1-O1	-9.40	108.00	119.55
2	A	2	9HI	O2-S1-O1	-9.37	108.03	119.55

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	4	9HI	O2-S1-O1	-9.27	108.15	119.55
2	A	1	9HI	C2-S1-N2	5.10	112.44	106.68
2	C	4	9HI	C1-C5-C27	-4.39	116.56	128.64
2	A	1	9HI	C1-C5-C27	-4.21	117.06	128.64
2	D	3	9HI	C1-C5-C27	-3.90	117.91	128.64
2	A	2	9HI	C1-C5-C27	-3.59	118.76	128.64
2	A	2	9HI	O1-S1-C2	3.10	113.35	108.74
2	D	3	9HI	C2-S1-N2	2.83	109.88	106.68
2	D	3	9HI	C19-C28-C22	-2.69	119.25	122.83
2	D	3	9HI	C25-C22-C28	2.61	121.06	118.36
2	A	1	9HI	O1-S1-C2	2.56	112.55	108.74
2	A	2	9HI	C7-N1-C5	-2.50	120.87	124.83
2	A	2	9HI	C2-C12-C6	-2.47	125.70	130.44
2	A	2	9HI	C7-N1-C12	2.45	128.82	124.72
2	A	2	9HI	C11-C10-C9	-2.44	110.47	114.18
2	A	2	9HI	C7-C8-C9	-2.43	108.30	113.31
2	A	1	9HI	C2-C12-C6	-2.38	125.87	130.44
2	A	1	9HI	C10-C11-C35	-2.38	108.45	113.19
2	D	3	9HI	O1-S1-C2	2.34	112.22	108.74
2	A	1	9HI	C5-N1-C12	2.27	110.27	108.78
2	A	2	9HI	C25-C22-C28	2.26	120.70	118.36
2	C	4	9HI	C10-C11-C35	-2.25	108.71	113.19
2	C	4	9HI	O1-S1-C2	2.24	112.08	108.74
2	C	4	9HI	C19-C28-C22	-2.24	119.85	122.83
2	D	3	9HI	C2-C12-C6	-2.23	126.16	130.44
2	D	3	9HI	C10-C11-C35	-2.23	108.75	113.19
2	A	1	9HI	C25-C31-C1	-2.22	117.11	120.79
2	A	2	9HI	C5-N1-C12	2.16	110.20	108.78
2	C	4	9HI	C2-C12-C6	-2.15	126.32	130.44
2	A	2	9HI	C26-C32-N2	2.14	124.73	120.09
2	A	2	9HI	C19-C28-C22	-2.13	119.99	122.83
2	C	4	9HI	C2-S1-N2	2.12	109.08	106.68
2	D	3	9HI	C7-N1-C12	2.10	128.25	124.72
2	C	4	9HI	C25-C22-C28	2.04	120.48	118.36
2	C	4	9HI	C24-C30-C15	-2.02	120.14	122.83
2	C	4	9HI	C5-N1-C12	2.01	110.10	108.78
2	D	3	9HI	C24-C30-C15	-2.00	120.16	122.83

There are no chirality outliers.

All (41) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	2	9HI	C20-C29-C3-N3
2	A	2	9HI	C23-C29-C3-N3
2	A	2	9HI	C20-C29-C3-O5
2	A	2	9HI	C23-C29-C3-O5
2	C	4	9HI	C20-C29-C3-N3
2	C	4	9HI	C20-C29-C3-O5
2	D	3	9HI	C20-C29-C3-N3
2	D	3	9HI	C23-C29-C3-N3
2	D	3	9HI	C20-C29-C3-O5
2	A	1	9HI	C32-N2-S1-O1
2	C	4	9HI	C23-C29-C3-N3
2	C	4	9HI	C23-C29-C3-O5
2	D	3	9HI	C23-C29-C3-O5
2	A	1	9HI	C32-N2-S1-C2
2	D	3	9HI	C32-N2-S1-O1
2	A	1	9HI	C18-C27-C5-N1
2	A	1	9HI	C32-N2-S1-O2
2	A	1	9HI	C21-C27-C5-N1
2	A	2	9HI	N1-C12-C6-C13
2	A	1	9HI	C18-C27-C5-C1
2	C	4	9HI	C18-C27-C5-C1
2	D	3	9HI	C18-C27-C5-C1
2	C	4	9HI	C21-C27-C5-N1
2	C	4	9HI	C18-C27-C5-N1
2	A	1	9HI	C21-C27-C5-C1
2	C	4	9HI	C21-C27-C5-C1
2	D	3	9HI	C18-C27-C5-N1
2	A	2	9HI	C2-C1-C31-C16
2	D	3	9HI	C2-C1-C31-C25
2	D	3	9HI	C21-C27-C5-C1
2	A	2	9HI	C18-C27-C5-C1
2	A	2	9HI	C2-C1-C31-C25
2	D	3	9HI	C2-C1-C31-C16
2	D	3	9HI	C21-C27-C5-N1
2	A	2	9HI	C21-C27-C5-C1
2	A	2	9HI	C5-C1-C31-C16
2	A	2	9HI	C5-C1-C31-C25
2	A	2	9HI	C18-C27-C5-N1
2	D	3	9HI	C5-C1-C31-C25
2	C	4	9HI	C5-C1-C31-C25
2	D	3	9HI	C5-C1-C31-C16


There are no ring outliers.


4 monomers are involved in 14 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	4	9HI	3	0
2	A	1	9HI	6	0
2	D	3	9HI	3	0
2	A	2	9HI	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	421/441 (95%)	0.27	32 (7%) 13 18	20, 32, 58, 65	0
1	B	421/441 (95%)	0.02	10 (2%) 59 66	21, 32, 44, 53	0
1	C	415/441 (94%)	0.30	29 (6%) 16 21	19, 34, 69, 91	0
1	D	396/441 (89%)	0.17	28 (7%) 16 21	20, 31, 68, 95	0
All	All	1653/1764 (93%)	0.19	99 (5%) 21 28	19, 32, 58, 95	0

All (99) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	861	HIS	9.6
1	D	461	PHE	7.2
1	A	479	TYR	7.0
1	D	861	HIS	6.9
1	C	461	PHE	6.9
1	B	861	HIS	6.7
1	D	475	HIS	6.6
1	D	474	LYS	6.5
1	C	452	LEU	6.2
1	A	456	GLU	6.1
1	C	460	LYS	6.1
1	A	483	THR	5.6
1	D	458	GLY	5.4
1	A	477	PRO	5.3
1	C	450	GLN	5.3
1	A	452	LEU	5.2
1	C	483	THR	5.0
1	D	486	GLU	4.9
1	C	451	ILE	4.8
1	C	462	LEU	4.8
1	C	444	PRO	4.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	451	ILE	4.6
1	D	471	VAL	4.6
1	C	861	HIS	4.4
1	A	450	GLN	4.4
1	A	480	LYS	4.3
1	A	475	HIS	4.2
1	A	482	GLU	4.2
1	D	485	ILE	4.2
1	D	463	SER	4.1
1	B	860	GLY	4.0
1	C	454	ASN	4.0
1	A	444	PRO	3.9
1	A	448	CYS	3.7
1	C	459	ALA	3.7
1	D	860	GLY	3.6
1	D	484	LEU	3.6
1	A	476	ILE	3.6
1	A	484	LEU	3.6
1	C	524	GLY	3.6
1	C	525	ALA	3.5
1	D	469	GLN	3.5
1	A	461	PHE	3.4
1	D	468	ILE	3.4
1	D	460	LYS	3.4
1	B	657[A]	MET	3.3
1	A	455	ALA	3.3
1	A	447	GLU	3.2
1	D	470	LEU	3.2
1	A	478	ALA	3.2
1	A	457	LYS	3.1
1	A	829	ASP	3.1
1	B	524	GLY	3.1
1	C	446	GLU	3.0
1	D	523	MET	3.0
1	D	459	ALA	3.0
1	A	446	GLU	3.0
1	A	474	LYS	2.9
1	A	772	VAL	2.8
1	B	446	GLU	2.8
1	D	465	ALA	2.7
1	A	473	ALA	2.7
1	B	450	GLN	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	486	GLU	2.7
1	C	523	MET	2.7
1	C	447	GLU	2.6
1	B	829	ASP	2.6
1	C	771	ASN	2.6
1	C	449	LEU	2.6
1	D	472	ASN	2.6
1	D	462	LEU	2.5
1	C	458	GLY	2.5
1	C	484	LEU	2.5
1	C	487	THR	2.5
1	B	479	TYR	2.4
1	A	454	ASN	2.4
1	D	487	THR	2.4
1	D	786	PRO	2.4
1	A	828	LYS	2.3
1	A	786	PRO	2.3
1	D	488	HIS	2.3
1	D	527	CYS	2.3
1	A	467	ILE	2.2
1	C	627	ARG	2.2
1	A	481	LEU	2.2
1	C	672	HIS	2.2
1	B	693	PRO	2.2
1	C	453	GLY	2.2
1	D	524	GLY	2.2
1	D	694	ALA	2.2
1	D	828	LYS	2.1
1	C	860	GLY	2.1
1	C	828	LYS	2.1
1	A	445	ASN	2.1
1	A	462	LEU	2.1
1	B	482	GLU	2.0
1	C	485	ILE	2.0
1	C	478	ALA	2.0
1	D	772	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	9HI	A	1	46/46	0.86	0.20	27,44,53,55	0
2	9HI	C	4	46/46	0.86	0.19	29,45,53,53	0
2	9HI	D	3	46/46	0.87	0.19	25,44,52,54	0
2	9HI	A	2	46/46	0.91	0.17	29,44,50,52	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around 9HI A 1:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around 9HI C 4:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around 9HI D 3:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.