



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 23, 2024 – 06:22 PM EDT

PDB ID : 3CFV
Title : Structural basis of the interaction of RbAp46/RbAp48 with histone H4
Authors : Pei, X.-Y.; Murzina, N.V.; Zhang, W.; McLaughlin, S.; Verreault, A.; Luisi, B.F.; Laue, E.D.
Deposited on : 2008-03-04
Resolution : 2.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3163 (2.60-2.60)
Clashscore	141614	3518 (2.60-2.60)
Ramachandran outliers	138981	3455 (2.60-2.60)
Sidechain outliers	138945	3455 (2.60-2.60)
RSRZ outliers	127900	3104 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	414	
1	B	414	
2	E	18	
2	F	18	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	ARS	A	412	-	-	X	-

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 6782 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Histone-binding protein RBBP7.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	B	393	3140	1984	538	605	5	8	0	0	0
1	A	390	3114	1969	533	599	5	8	0	0	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-2	HIS	-	cloning artifact	UNP Q16576
B	-1	MSE	-	cloning artifact	UNP Q16576
B	0	ALA	-	cloning artifact	UNP Q16576
A	-2	HIS	-	cloning artifact	UNP Q16576
A	-1	MSE	-	cloning artifact	UNP Q16576
A	0	ALA	-	cloning artifact	UNP Q16576

- Molecule 2 is a protein called Histone H4 peptide.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	E	18	142	86	33	23	0	0	0
2	F	14	109	67	27	15	0	0	0

- Molecule 3 is ARSENIC (three-letter code: ARS) (formula: As).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	As	0	0
			1	1		
3	A	1	Total	As	0	0
			1	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	125	Total 125	O 125	0	0
4	E	7	Total 7	O 7	0	0
4	A	141	Total 141	O 141	0	0
4	F	2	Total 2	O 2	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Histone-binding protein RBBP7



- Molecule 1: Histone-binding protein RBBP7




- Molecule 2: Histone H4 peptide

Chain E:  72% 17% 11%



• Molecule 2: Histone H4 peptide

Chain F:  17% 50% 22% 6% 22%



4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	108.66Å 44.79Å 109.59Å 90.00° 90.71° 90.00°	Depositor
Resolution (Å)	25.00 – 2.60 24.95 – 2.60	Depositor EDS
% Data completeness (in resolution range)	99.6 (25.00-2.60) 99.6 (24.95-2.60)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	0.14	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.02 (at 2.60Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.199 , 0.257 0.201 , 0.257	Depositor DCC
R_{free} test set	1577 reflections (4.78%)	wwPDB-VP
Wilson B-factor (Å ²)	46.1	Xtrriage
Anisotropy	0.213	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 68.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.074 for l,k,-h 0.022 for h,-k,-l 0.021 for l,-k,h	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	6782	wwPDB-VP
Average B, all atoms (Å ²)	51.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.02% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ARS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.23	0/3194	0.42	0/4340
1	B	0.23	0/3220	0.48	2/4374 (0.0%)
2	E	0.24	0/142	0.39	0/188
2	F	0.34	0/109	1.05	2/143 (1.4%)
All	All	0.23	0/6665	0.46	4/9045 (0.0%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	7	PHE	CB-CA-C	-10.64	89.12	110.40
2	F	30	THR	CB-CA-C	-8.19	89.48	111.60
1	B	7	PHE	N-CA-C	6.93	129.71	111.00
2	F	30	THR	N-CA-C	5.52	125.91	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3114	0	2973	77	0
1	B	3140	0	2989	81	0
2	E	142	0	156	4	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	F	109	0	127	15	0
3	A	1	0	0	2	0
3	B	1	0	0	1	0
4	A	141	0	0	0	0
4	B	125	0	0	1	0
4	E	7	0	0	0	0
4	F	2	0	0	0	0
All	All	6782	0	6245	175	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

The worst 5 of 175 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:F:29:ILE:N	2:F:30:THR:HB	1.43	1.32
1:B:6:MSE:HA	1:B:7:PHE:HB3	1.27	1.12
1:B:35:MSE:HE3	1:B:112:GLY:HA3	1.32	1.07
1:B:116:CYS:SG	3:B:412:ARS:AS	2.73	1.06
2:F:29:ILE:H	2:F:30:THR:CB	1.70	1.02

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	386/414 (93%)	368 (95%)	15 (4%)	3 (1%)	19	39
1	B	389/414 (94%)	370 (95%)	15 (4%)	4 (1%)	15	32
2	E	16/18 (89%)	16 (100%)	0	0	100	100
2	F	12/18 (67%)	10 (83%)	1 (8%)	1 (8%)	1	1

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	803/864 (93%)	764 (95%)	31 (4%)	8 (1%)	15	32

5 of 8 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	314	SER
1	B	5	GLU
1	B	406	ASN
1	A	314	SER
2	F	30	THR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	345/356 (97%)	321 (93%)	24 (7%)	15	30
1	B	348/356 (98%)	321 (92%)	27 (8%)	12	25
2	E	14/14 (100%)	12 (86%)	2 (14%)	3	5
2	F	10/14 (71%)	10 (100%)	0	100	100
All	All	717/740 (97%)	664 (93%)	53 (7%)	13	28

5 of 53 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	E	26	ILE
1	A	153	TYR
1	A	314	SER
1	A	34	VAL
1	A	62	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 36 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	225	HIS
1	A	399	GLN
1	A	249	GLN
1	A	321	GLN
1	B	315	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	382/414 (92%)	0.17	30 (7%) 12 9	17, 47, 84, 120	1 (0%)
1	B	385/414 (92%)	0.27	31 (8%) 12 8	14, 47, 86, 120	7 (1%)
2	E	18/18 (100%)	0.48	0 100 100	46, 77, 96, 99	0
2	F	14/18 (77%)	0.60	3 (21%) 0 0	46, 63, 94, 94	0
All	All	799/864 (92%)	0.23	64 (8%) 12 9	14, 48, 89, 120	8 (1%)

The worst 5 of 64 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	212	GLU	6.7
1	B	89	ASP	5.6
1	B	400	ILE	5.5
1	B	368	ILE	5.4
1	B	211	LYS	5.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	ARS	B	412	1/1	0.96	0.21	99,99,99,99	0
3	ARS	A	412	1/1	0.99	0.12	80,80,80,80	0

6.5 Other polymers [i](#)

There are no such residues in this entry.