



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 20, 2023 – 11:42 AM JST

PDB ID : 7CIG
Title : Crystal structure of L-methionine decarboxylase Q64A mutant from Streptomyces sp.590 in complexed with L- methionine methyl ester (geminal diamine form).
Authors : Okawa, A.; Shiba, T.; Hayashi, M.; Onoue, Y.; Murota, M.; Sato, D.; Inagaki, J.; Tamura, T.; Harada, S.; Inagaki, K.
Deposited on : 2020-07-07
Resolution : 1.45 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1156 (1.46-1.46)
Clashscore	141614	1202 (1.46-1.46)
Ramachandran outliers	138981	1178 (1.46-1.46)
Sidechain outliers	138945	1178 (1.46-1.46)
RSRZ outliers	127900	1139 (1.46-1.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	557	
1	B	557	

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 8809 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called L-methionine decarboxylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	518	Total 4100	C 2608	N 715	O 761	S 16	0	5	0
1	B	521	Total 4136	C 2631	N 720	O 769	S 16	0	7	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	64	ALA	GLN	engineered mutation	UNP A0A0G4DBU7
B	64	ALA	GLN	engineered mutation	UNP A0A0G4DBU7

- Molecule 2 is methyl (2S)-2-[[2-methyl-3-oxidanyl-5-(phosphonooxymethyl)pyridin-4-yl]methylamino]-4-methylsulfanyl-butanoate (three-letter code: G03) (formula: C₁₄H₂₃N₂O₇PS) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
			Total	C	N	O	P			S
2	A	1	25	14	2	7	1	1	0	0
2	B	1	25	14	2	7	1	1	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	278	Total	O	0	0
			278	278		
3	B	245	Total	O	0	0
			245	245		


3 Residue-property plots i

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: L-methionine decarboxylase



- Molecule 1: L-methionine decarboxylase



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	125.04Å 147.38Å 53.83Å 90.00° 100.01° 90.00°	Depositor
Resolution (Å)	19.87 – 1.45 19.87 – 1.45	Depositor EDS
% Data completeness (in resolution range)	99.8 (19.87-1.45) 99.8 (19.87-1.45)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.60 (at 1.45Å)	Xtrriage
Refinement program	REFMAC 5.8.0103	Depositor
R, R_{free}	0.175 , 0.194 0.181 , 0.198	Depositor DCC
R_{free} test set	8469 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	17.4	Xtrriage
Anisotropy	0.043	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 46.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	8809	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.58% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: G03

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.50	0/4219	0.74	3/5749 (0.1%)
1	B	0.49	0/4255	0.74	4/5800 (0.1%)
All	All	0.49	0/8474	0.74	7/11549 (0.1%)

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	470	ARG	NE-CZ-NH2	6.37	123.49	120.30
1	A	470	ARG	NE-CZ-NH2	6.09	123.35	120.30
1	B	394	LYS	CD-CE-NZ	5.98	125.45	111.70
1	A	394	LYS	CD-CE-NZ	5.65	124.70	111.70
1	B	38	ASP	CB-CG-OD1	5.21	122.99	118.30
1	B	57	ARG	NE-CZ-NH2	5.19	122.90	120.30
1	A	456	ARG	NE-CZ-NH1	5.04	122.82	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4100	0	3952	15	0
1	B	4136	0	3985	20	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	25	0	0	0	0
2	B	25	0	0	0	0
3	A	278	0	0	1	0
3	B	245	0	0	4	0
All	All	8809	0	7937	34	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (34) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:89:ASP:OD1	1:A:427:THR:HG21	1.71	0.90
1:A:516:VAL:HG22	1:A:522:THR:HG21	1.55	0.88
1:B:516:VAL:HG22	1:B:522:THR:HG21	1.66	0.75
1:A:258[B]:ASP:OD1	1:A:297[B]:ARG:NH1	2.29	0.66
1:B:258[B]:ASP:OD2	1:B:297[B]:ARG:NH2	2.32	0.63
1:B:284:PHE:O	1:B:528:HIS:HD2	1.82	0.63
1:A:202:VAL:HG13	1:A:207:VAL:O	1.99	0.62
1:A:393:HIS:H	1:A:393:HIS:CD2	2.17	0.61
1:A:284:PHE:O	1:A:528:HIS:HD2	1.83	0.61
1:A:494[A]:ARG:HG3	3:A:702:HOH:O	2.00	0.61
1:B:393:HIS:CD2	1:B:393:HIS:H	2.21	0.58
1:A:185:HIS:O	1:A:272:HIS:HD2	1.91	0.54
1:B:430:ALA:HB3	3:B:708:HOH:O	2.08	0.53
1:B:241:ARG:HH12	1:B:257:VAL:HG23	1.74	0.52
1:A:218:TYR:OH	1:A:272:HIS:HE1	1.94	0.51
1:B:200:LYS:HE3	3:B:874:HOH:O	2.11	0.50
1:B:218:TYR:OH	1:B:272:HIS:HE1	1.95	0.49
1:B:87:LEU:HD21	1:B:422:ILE:HG22	1.94	0.49
1:A:376:THR:CG2	1:A:382:VAL:HG23	2.43	0.48
1:A:72:THR:HB	3:B:701:HOH:O	2.13	0.47
1:B:425:PRO:HB2	1:B:427:THR:HG23	1.99	0.45
1:B:69:MET:HB3	1:B:72[B]:THR:HG23	1.99	0.45
1:B:72[A]:THR:OG1	1:B:73:ALA:N	2.50	0.45
1:B:185:HIS:O	1:B:272:HIS:HD2	2.00	0.45
1:B:265:GLU:OE1	1:B:304:ARG:NH1	2.51	0.43
1:B:516:VAL:HG23	1:B:519:ASP:HB3	2.00	0.43
1:A:516:VAL:HG23	1:A:519:ASP:HB3	2.01	0.43
1:B:146[B]:LEU:HD11	1:B:198:PHE:CD1	2.53	0.43
1:B:102:VAL:HG22	1:B:105:ARG:NH2	2.35	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:393:HIS:CD2	1:B:401:PRO:HA	2.55	0.42
1:A:75:ASP:OD2	1:A:447:ARG:NH2	2.54	0.41
1:A:62:GLY:HA3	1:A:508:LEU:HA	2.03	0.40
1:A:429:PHE:HE2	1:B:429:PHE:HE2	1.69	0.40
1:B:355:GLU:HG2	3:B:938:HOH:O	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	519/557 (93%)	501 (96%)	18 (4%)	0	100	100
1	B	524/557 (94%)	506 (97%)	17 (3%)	1 (0%)	47	22
All	All	1043/1114 (94%)	1007 (96%)	35 (3%)	1 (0%)	51	24

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	322	ARG

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	426/449 (95%)	416 (98%)	10 (2%)	50 17
1	B	430/449 (96%)	420 (98%)	10 (2%)	50 17
All	All	856/898 (95%)	836 (98%)	20 (2%)	50 17

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	119	ARG
1	A	224	LEU
1	A	292	ARG
1	A	313	VAL
1	A	422	ILE
1	A	436	PHE
1	A	494[A]	ARG
1	A	494[B]	ARG
1	A	520	GLU
1	A	528	HIS
1	B	92	GLN
1	B	179	ARG
1	B	180	ASN
1	B	224	LEU
1	B	309	GLN
1	B	317	CYS
1	B	352	GLU
1	B	436	PHE
1	B	494	ARG
1	B	528	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	182	ASN
1	A	272	HIS
1	A	279	ASN
1	A	393	HIS
1	A	528	HIS
1	B	160	GLN
1	B	182	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	272	HIS
1	B	279	ASN
1	B	393	HIS
1	B	528	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	G03	A	601	1	25,25,25	2.22	2 (8%)	31,34,34	1.91	5 (16%)
2	G03	B	601	1	25,25,25	2.13	3 (12%)	31,34,34	1.92	4 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	G03	A	601	1	-	8/21/21/21	0/1/1/1
2	G03	B	601	1	-	8/21/21/21	0/1/1/1

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	601	G03	CAF-CAE	-9.28	1.40	1.51
2	B	601	G03	CAF-CAE	-8.95	1.40	1.51
2	A	601	G03	CAX-NAY	4.03	1.42	1.34
2	B	601	G03	CAC-CAB	-3.22	1.37	1.40
2	B	601	G03	CAX-NAY	2.83	1.40	1.34

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	601	G03	CAF-N-CA	7.54	128.21	113.92
2	B	601	G03	CAF-N-CA	7.32	127.80	113.92
2	B	601	G03	C-CA-N	-4.61	103.16	112.11
2	A	601	G03	C-CA-N	-3.72	104.89	112.11
2	A	601	G03	CAE-CAF-N	2.57	118.86	111.78
2	A	601	G03	CG-CB-CA	-2.52	105.76	112.91
2	A	601	G03	CAO-OAN-C	2.39	121.34	115.94
2	B	601	G03	CAE-CAF-N	2.33	118.20	111.78
2	B	601	G03	OAS-PAT-OAV	-2.17	100.40	106.47

There are no chirality outliers.

All (16) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	601	G03	CB-CA-N-CAF
2	B	601	G03	CB-CA-N-CAF
2	A	601	G03	CA-C-OAN-CAO
2	A	601	G03	O-C-OAN-CAO
2	A	601	G03	CAQ-CAE-CAF-N
2	B	601	G03	CA-C-OAN-CAO
2	B	601	G03	O-C-OAN-CAO
2	B	601	G03	CAQ-CAE-CAF-N
2	B	601	G03	N-CA-CB-CG
2	A	601	G03	CAC-CAE-CAF-N
2	B	601	G03	CAC-CAE-CAF-N
2	B	601	G03	C-CA-CB-CG
2	A	601	G03	C-CA-N-CAF

Continued on next page...


Continued from previous page...


Mol	Chain	Res	Type	Atoms
2	B	601	G03	C-CA-N-CAF
2	A	601	G03	N-CA-CB-CG
2	A	601	G03	C-CA-CB-CG

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	518/557 (92%)	0.83	62 (11%) 4 4	11, 18, 48, 118	0
1	B	521/557 (93%)	0.90	64 (12%) 4 4	11, 19, 43, 149	0
All	All	1039/1114 (93%)	0.87	126 (12%) 4 4	11, 18, 46, 149	0

All (126) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	317	CYS	22.4
1	A	318	PRO	18.3
1	B	315	GLY	17.9
1	B	322	ARG	17.6
1	B	319	GLN	16.6
1	A	317	CYS	15.1
1	A	325	VAL	15.0
1	B	321	GLY	14.5
1	B	324	LEU	13.6
1	B	318	PRO	13.4
1	B	316	SER	12.8
1	B	325	VAL	12.1
1	B	161	PRO	11.7
1	A	418	GLN	11.1
1	A	424	ALA	10.7
1	B	320	THR	10.3
1	A	421	TYR	10.2
1	B	18	ALA	10.1
1	A	315	GLY	9.8
1	B	418	GLN	9.8
1	A	319	GLN	9.8
1	A	316	SER	9.6
1	A	422	ILE	9.4
1	A	314	TYR	9.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	314	TYR	9.2
1	A	321	GLY	9.0
1	A	323	PRO	8.7
1	B	313	VAL	8.6
1	A	420	ASP	8.3
1	A	320	THR	8.2
1	A	324	LEU	8.1
1	B	323	PRO	7.9
1	A	162	PRO	7.5
1	A	161	PRO	7.3
1	A	416	PRO	7.1
1	B	421	TYR	7.1
1	A	423	GLY	6.5
1	A	322	ARG	6.4
1	B	424	ALA	6.2
1	B	423	GLY	6.2
1	A	419	PRO	6.2
1	A	160	GLN	6.1
1	B	422	ILE	6.1
1	A	313	VAL	6.0
1	A	557	PRO	5.7
1	B	159	ILE	5.6
1	B	179	ARG	5.4
1	B	518	GLY	5.2
1	A	25	LEU	5.2
1	A	417	SER	5.0
1	B	555	GLY	5.0
1	A	179	ARG	4.9
1	A	518	GLY	4.6
1	B	160	GLN	4.6
1	B	420	ASP	4.5
1	A	425	PRO	4.2
1	B	312	VAL	4.2
1	B	327	VAL	4.1
1	A	312	VAL	4.1
1	A	415	PRO	4.0
1	B	19	VAL	4.0
1	B	419	PRO	3.8
1	B	520	GLU	3.8
1	A	159	ILE	3.8
1	A	24	GLU	3.6
1	B	416	PRO	3.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	26	ASP	3.6
1	A	93	SER	3.6
1	B	415	PRO	3.5
1	A	362	GLU	3.3
1	A	68	ASP	3.3
1	B	93	SER	3.3
1	B	68	ASP	3.2
1	B	380	GLY	3.2
1	B	417	SER	3.2
1	A	379	HIS	3.1
1	A	429	PHE	3.1
1	B	414	SER	3.1
1	A	284	PHE	3.0
1	A	327	VAL	3.0
1	B	196	TYR	2.9
1	B	26	ASP	2.9
1	B	23	PRO	2.9
1	A	414	SER	2.8
1	A	556	ALA	2.8
1	B	379	HIS	2.8
1	A	430	ALA	2.7
1	B	377	ALA	2.7
1	A	70	GLN	2.7
1	A	481	GLU	2.7
1	A	311	GLU	2.7
1	B	21	PRO	2.7
1	B	311	GLU	2.7
1	A	157	ALA	2.7
1	B	158	LEU	2.7
1	A	520	GLU	2.6
1	B	378	GLY	2.6
1	B	22	GLY	2.6
1	B	70	GLN	2.5
1	B	425	PRO	2.5
1	A	519	ASP	2.5
1	A	427	THR	2.5
1	B	25	LEU	2.4
1	A	92	GLN	2.4
1	B	24	GLU	2.3
1	B	157	ALA	2.3
1	A	366	PRO	2.3
1	A	480	VAL	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	216	GLU	2.2
1	A	514	LEU	2.2
1	B	547	LEU	2.2
1	B	284	PHE	2.2
1	B	366	PRO	2.1
1	B	514	LEU	2.1
1	A	180	ASN	2.1
1	A	216	GLU	2.1
1	A	181	PRO	2.1
1	A	355	GLU	2.1
1	B	44	ARG	2.1
1	B	523	ARG	2.1
1	B	376	THR	2.1
1	B	355	GLU	2.0
1	B	521	THR	2.0
1	B	184	HIS	2.0
1	A	156	LYS	2.0
1	A	517	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	G03	A	601	25/25	0.94	0.12	13,15,27,31	0
2	G03	B	601	25/25	0.94	0.13	13,15,32,37	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers

as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around G03 A 601:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers ⓘ

There are no such residues in this entry.