



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 21, 2024 – 04:26 PM EDT

PDB ID : 9CKJ
Title : Co-crystal structure of 53BP1 tandem Tudor domains in complex with UNC9512
Authors : Zeng, H.; Dong, A.; Shell, D.J.; Foley, C.; Arrowsmith, C.H.; Edwards, A.M.; Halabelian, L.; Structural Genomics Consortium (SGC)
Deposited on : 2024-07-09
Resolution : 2.25 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 1.20.1
EDS : **FAILED**
buster-report : 1.1.7 (2018)
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	180529	1919 (2.26-2.26)
Ramachandran outliers	177936	1884 (2.26-2.26)
Sidechain outliers	177891	1885 (2.26-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS failed to run properly.

Mol	Chain	Length	Quality of chain
1	A	124	
1	B	124	
1	C	124	
1	D	124	
1	E	124	
1	F	124	

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 5979 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called TP53-binding protein 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	118	Total 935	C 601	N 156	O 175	S 3	0	0	0
1	B	120	Total 929	C 599	N 152	O 175	S 3	0	0	0
1	C	119	Total 952	C 610	N 158	O 181	S 3	0	1	0
1	D	119	Total 903	C 581	N 148	O 171	S 3	0	0	0
1	E	118	Total 884	C 576	N 140	O 165	S 3	0	0	0
1	F	121	Total 936	C 605	N 150	O 178	S 3	0	0	0

- Molecule 2 is (3S)-1-[4-(4-methylpiperazin-1-yl)pyridine-2-carbonyl]-N-[(3M)-3-(3-oxo-2,3-dihydro-1H-isoindol-5-yl)phenyl]piperidine-3-carboxamide (three-letter code: A1AY8) (formula: C₃₁H₃₄N₆O₃) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			40	31	6	3		
2	B	1	Total	C	N	O	0	0
			40	31	6	3		
2	C	1	Total	C	N	O	0	0
			40	31	6	3		
2	D	1	Total	C	N	O	0	0
			40	31	6	3		
2	E	1	Total	C	N	O	0	0
			40	31	6	3		
2	F	1	Total	C	N	O	0	0
			40	31	6	3		

- Molecule 3 is UNKNOWN ATOM OR ION (three-letter code: UNX) (formula: X).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	F	2	Total	X	0	0
			2	2		

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	56	Total	O	0	0
			56	56		
4	B	40	Total	O	0	1
			41	41		
4	C	48	Total	O	0	1
			49	49		
4	D	19	Total	O	0	0
			19	19		
4	E	12	Total	O	0	0
			12	12		
4	F	21	Total	O	0	0
			21	21		

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS failed to run properly.


- Molecule 1: TP53-binding protein 1




- Molecule 1: TP53-binding protein 1




- Molecule 1: TP53-binding protein 1




- Molecule 1: TP53-binding protein 1



- Molecule 1: TP53-binding protein 1



- Molecule 1: TP53-binding protein 1

Chain F:  90% 6% ..



4 Data and refinement statistics

EDS failed to run properly - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	41.88Å 76.03Å 258.64Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.92 – 2.25	Depositor
% Data completeness (in resolution range)	97.5 (29.92-2.25)	Depositor
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.60 (at 2.27Å)	Xtrriage
Refinement program	BUSTER 2.10.3	Depositor
R, R_{free}	0.219 , 0.261	Depositor
Wilson B-factor (Å ²)	40.4	Xtrriage
Anisotropy	0.341	Xtrriage
L-test for twinning ²	$\langle L \rangle = 0.42$, $\langle L^2 \rangle = 0.25$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	5979	wwPDB-VP
Average B, all atoms (Å ²)	48.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.23% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: A1AY8, UNX

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.60	0/954	0.75	0/1284
1	B	0.60	0/948	0.73	0/1279
1	C	0.59	0/971	0.75	0/1306
1	D	0.52	0/922	0.70	0/1248
1	E	0.48	0/903	0.71	0/1223
1	F	0.52	0/957	0.69	0/1295
All	All	0.55	0/5655	0.72	0/7635

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	935	0	908	8	0
1	B	929	0	882	4	0
1	C	952	0	919	5	0
1	D	903	0	826	6	0
1	E	884	0	800	4	0
1	F	936	0	871	5	0
2	A	40	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	40	0	0	0	0
2	C	40	0	0	0	0
2	D	40	0	0	0	0
2	E	40	0	0	0	0
2	F	40	0	0	0	0
3	F	2	0	0	0	0
4	A	56	0	0	1	0
4	B	41	0	0	0	0
4	C	49	0	0	0	0
4	D	19	0	0	0	0
4	E	12	0	0	0	0
4	F	21	0	0	1	0
All	All	5979	0	5206	32	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (32) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:1562:ARG:HD2	4:F:2120:HOH:O	2.01	0.61
1:A:1544:VAL:HG21	1:A:1572:ILE:HD12	1.82	0.60
1:A:1490:ARG:HH11	1:A:1490:ARG:HB3	1.68	0.57
1:A:1544:VAL:CG2	1:A:1572:ILE:HD12	2.35	0.57
1:F:1492:VAL:HG12	1:F:1501:PHE:HB3	1.91	0.53
1:E:1490:ARG:HH11	1:E:1490:ARG:HB3	1.76	0.51
1:D:1544:VAL:CG2	1:D:1572:ILE:HD12	2.40	0.51
1:D:1544:VAL:HG21	1:D:1572:ILE:HD12	1.93	0.51
1:A:1501:PHE:HB2	1:A:1588:LEU:HB2	1.92	0.50
1:D:1492:VAL:HG12	1:D:1501:PHE:HB3	1.94	0.50
1:B:1572:ILE:HD12	1:B:1581:TYR:CD1	2.49	0.48
1:C:1501:PHE:HB2	1:C:1588:LEU:HB2	1.95	0.48
1:F:1562:ARG:HE	1:F:1564:GLU:HG3	1.79	0.48
1:E:1492:VAL:HG12	1:E:1501:PHE:HB3	1.96	0.47
1:A:1508:ARG:HH21	1:A:1516:LYS:HE3	1.78	0.47
1:D:1501:PHE:HB2	1:D:1588:LEU:HB2	1.97	0.47
1:A:1492:VAL:HG12	1:A:1501:PHE:HB3	1.96	0.47
1:B:1492:VAL:HG12	1:B:1501:PHE:HB3	1.96	0.47
1:B:1501:PHE:HB2	1:B:1588:LEU:HB2	1.98	0.45
1:C:1572:ILE:HD12	1:C:1581:TYR:CD1	2.52	0.45
1:E:1501:PHE:HB2	1:E:1588:LEU:HB2	1.99	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:1572:ILE:HD12	1:E:1581:TYR:CD1	2.52	0.44
1:A:1490:ARG:HG2	1:A:1505:LYS:HG2	2.00	0.43
1:A:1562:ARG:HB2	4:A:2155:HOH:O	2.17	0.43
1:F:1501:PHE:HB2	1:F:1588:LEU:HB2	2.01	0.43
1:B:1500:TYR:CD1	1:B:1589:SER:HA	2.54	0.43
1:F:1572:ILE:HD12	1:F:1581:TYR:CD1	2.55	0.41
1:C:1583[B]:ARG:HE	1:C:1583[B]:ARG:HB3	1.51	0.41
1:C:1492:VAL:HG12	1:C:1501:PHE:HB3	2.01	0.41
1:D:1493:ALA:HB3	1:D:1517:LEU:HD22	2.02	0.40
1:C:1500:TYR:CD2	1:C:1589:SER:HA	2.56	0.40
1:D:1486:PHE:HA	1:D:1489:LEU:HD12	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	116/124 (94%)	114 (98%)	2 (2%)	0	100	100
1	B	118/124 (95%)	113 (96%)	5 (4%)	0	100	100
1	C	118/124 (95%)	115 (98%)	3 (2%)	0	100	100
1	D	117/124 (94%)	113 (97%)	4 (3%)	0	100	100
1	E	116/124 (94%)	112 (97%)	4 (3%)	0	100	100
1	F	119/124 (96%)	117 (98%)	2 (2%)	0	100	100
All	All	704/744 (95%)	684 (97%)	20 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	94/104 (90%)	92 (98%)	2 (2%)	48	57
1	B	91/104 (88%)	86 (94%)	5 (6%)	18	18
1	C	97/104 (93%)	96 (99%)	1 (1%)	73	80
1	D	83/104 (80%)	80 (96%)	3 (4%)	30	37
1	E	77/104 (74%)	74 (96%)	3 (4%)	27	33
1	F	89/104 (86%)	86 (97%)	3 (3%)	32	40
All	All	531/624 (85%)	514 (97%)	17 (3%)	34	42

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1498	ASN
1	A	1517	LEU
1	B	1498	ASN
1	B	1514	LYS
1	B	1517	LEU
1	B	1564	GLU
1	B	1583	ARG
1	C	1498	ASN
1	D	1498	ASN
1	D	1517	LEU
1	D	1563	LYS
1	E	1490	ARG
1	E	1498	ASN
1	E	1517	LEU
1	F	1498	ASN
1	F	1517	LEU
1	F	1564	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1498	ASN
1	B	1498	ASN
1	C	1498	ASN
1	D	1498	ASN
1	E	1498	ASN
1	F	1498	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 2 are unknown - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	A1AY8	C	2001	-	45,45,45	0.49	0	63,64,64	0.53	1 (1%)
2	A1AY8	D	2001	-	45,45,45	0.43	0	63,64,64	0.50	0
2	A1AY8	E	2001	-	45,45,45	0.45	0	63,64,64	0.47	0
2	A1AY8	B	2001	-	45,45,45	0.49	0	63,64,64	0.56	1 (1%)
2	A1AY8	F	2001	-	45,45,45	0.44	0	63,64,64	0.54	0
2	A1AY8	A	2001	-	45,45,45	0.49	0	63,64,64	0.55	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	A1AY8	C	2001	-	-	0/24/53/53	0/6/6/6
2	A1AY8	D	2001	-	-	1/24/53/53	0/6/6/6
2	A1AY8	E	2001	-	-	1/24/53/53	0/6/6/6
2	A1AY8	B	2001	-	-	1/24/53/53	0/6/6/6
2	A1AY8	F	2001	-	-	1/24/53/53	0/6/6/6
2	A1AY8	A	2001	-	-	0/24/53/53	0/6/6/6

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	2001	A1AY8	C6-N7-C8	2.15	123.96	118.11
2	C	2001	A1AY8	C6-N7-C8	2.02	123.60	118.11

There are no chirality outliers.


All (4) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	B	2001	A1AY8	C37-C22-N21-C20
2	E	2001	A1AY8	C37-C22-N21-C20
2	F	2001	A1AY8	C37-C22-N21-C20
2	D	2001	A1AY8	C37-C22-N21-C20


There are no ring outliers.


No monomer is involved in short contacts.


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.














5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

EDS failed to run properly - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS failed to run properly - this section is therefore empty.


6.3 Carbohydrates [i](#)

EDS failed to run properly - this section is therefore empty.

6.4 Ligands [i](#)




EDS failed to run properly - this section is therefore empty.

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around A1AY8 B 2001:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around A1AY8 C 2001:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around A1AY8 D 2001:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around A1AY8 E 2001:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

EDS failed to run properly - this section is therefore empty.