



Full wwPDB X-ray Structure Validation Report ⓘ

Dec 3, 2023 – 04:51 pm GMT

PDB ID : 2CM6
Title : crystal structure of the C2B domain of rabphilin3A
Authors : Schlicker, C.; Montaville, P.; Sheldrick, G.M.; Becker, S.
Deposited on : 2006-05-04
Resolution : 1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtrriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36

PERCENTILES INFOmissingINFO

1 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 2397 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called RABPHILIN-3A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	149	1153	743	193	211	6	0	0	0
1	B	148	1159	742	198	213	6	0	0	0

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Ca	0	0
			2	2		
2	B	2	Total	Ca	0	0
			2	2		

- Molecule 3 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	O	P	0	0
			5	4	1		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	39	Total	O	0	0
			39	39		
4	B	37	Total	O	0	0
			37	37		

SEQUENCE-PLOTS INFOmissingINFO

2 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	40.07Å 59.61Å 66.39Å 90.00° 103.89° 90.00°	Depositor
Resolution (Å)	43.76 – 1.85 43.76 – 1.85	Depositor EDS
% Data completeness (in resolution range)	97.2 (43.76-1.85) 96.6 (43.76-1.85)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.36 (at 1.86Å)	Xtrriage
Refinement program	SHELXL-97	Depositor
R, R_{free}	0.198 , 0.268 0.195 , 0.262	Depositor DCC
R_{free} test set	1270 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	21.6	Xtrriage
Anisotropy	0.303	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 74.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	2397	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.66% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

3 Model quality [i](#)

3.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.33	0/1177	0.94	1/1584 (0.1%)
1	B	0.32	0/1184	0.90	0/1594
All	All	0.32	0/2361	0.92	1/3178 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	533	GLU	C-N-CA	6.13	137.02	121.70

There are no chirality outliers.

There are no planarity outliers.

3.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1153	0	1072	23	0
1	B	1159	0	1100	19	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
3	B	5	0	0	0	0
4	A	39	0	0	2	0
4	B	37	0	0	1	0
All	All	2397	0	2172	42	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (42) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:526:LEU:HD21	1:A:634:ILE:HG22	1.55	0.89
1:B:1565:VAL:HA	1:B:1604:ASN:ND2	2.14	0.63
1:B:1565:VAL:HA	1:B:1604:ASN:HD22	1.69	0.58
1:B:1565:VAL:HG13	1:B:1566:HIS:CD2	2.41	0.55
1:B:1588:MET:SD	1:B:1623:LYS:HE3	2.46	0.55
1:B:1565:VAL:HG22	1:B:1604:ASN:HD22	1.72	0.54
1:A:566:HIS:H	1:A:604:ASN:ND2	2.06	0.54
1:B:1526:LEU:HD21	1:B:1634:ILE:HG13	1.88	0.53
1:B:1554:GLN:HG3	1:B:1558:ILE:HD11	1.91	0.53
1:A:543:LYS:HG2	1:A:565:VAL:HG12	1.91	0.52
1:B:1566:HIS:H	1:B:1604:ASN:ND2	2.08	0.52
1:B:1578:PRO:HA	1:B:1630:TRP:O	2.10	0.51
1:A:545:LEU:HG	1:A:562:ILE:HD12	1.91	0.51
1:A:632:TYR:OH	1:A:634:ILE:HD11	2.10	0.51
1:B:1617:HIS:CD2	1:B:1665:LYS:HD3	2.46	0.51
1:A:565:VAL:HG23	1:A:604:ASN:HD22	1.76	0.50
1:B:1600:LYS:HB3	1:B:1600:LYS:NZ	2.27	0.50
1:B:1526:LEU:O	1:B:1529:GLU:HB2	2.12	0.50
1:B:1565:VAL:CG2	1:B:1604:ASN:HD22	2.24	0.49
1:A:551:SER:OG	1:A:554:GLN:HB2	2.12	0.49
1:A:526:LEU:HD21	1:A:634:ILE:CG2	2.35	0.47
1:A:565:VAL:HG22	1:A:566:HIS:CD2	2.50	0.46
1:A:565:VAL:HA	1:A:604:ASN:HD22	1.79	0.46
1:B:1636:LYS:NZ	1:B:1637:SER:O	2.50	0.45
1:B:1664:ASN:ND2	4:B:2035:HOH:O	2.50	0.44
1:A:556:GLY:HA3	1:A:612:PHE:CZ	2.53	0.44
1:B:1565:VAL:HG22	1:B:1604:ASN:ND2	2.31	0.44
1:A:620:LEU:HD22	1:A:658:TRP:CZ2	2.53	0.43
1:B:1595:LYS:HB3	1:B:1595:LYS:HE3	1.57	0.43
1:A:650:ALA:O	1:A:655:LEU:HB2	2.19	0.43
1:A:549:MET:HG3	1:A:667:LYS:C	2.39	0.43
1:A:665:LYS:O	1:A:666:ASP:HB2	2.19	0.42
1:A:634:ILE:HD13	1:A:634:ILE:HA	1.77	0.42
1:A:595:LYS:HG3	1:A:596:THR:O	2.19	0.42
1:A:565:VAL:HG23	1:A:604:ASN:ND2	2.34	0.42
1:A:581:LYS:C	1:A:582:LEU:HD23	2.40	0.42
1:A:665:LYS:HG3	4:A:2037:HOH:O	2.18	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:664:ASN:ND2	4:A:2036:HOH:O	2.52	0.42
1:A:543:LYS:HG3	1:A:672:TRP:CE3	2.55	0.41
1:A:620:LEU:HD23	1:A:620:LEU:HA	1.88	0.41
1:B:1636:LYS:HZ3	1:B:1636:LYS:HG2	1.69	0.41
1:B:1566:HIS:H	1:B:1604:ASN:HD21	1.68	0.40

There are no symmetry-related clashes.

3.3 Torsion angles [i](#)

3.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	143/166 (86%)	137 (96%)	4 (3%)	2 (1%)	11	3
1	B	144/166 (87%)	139 (96%)	4 (3%)	1 (1%)	22	9
All	All	287/332 (86%)	276 (96%)	8 (3%)	3 (1%)	15	5

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	1540	GLU
1	A	540	GLU
1	A	534	ARG

3.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	108/144 (75%)	101 (94%)	7 (6%)	17	4
1	B	115/144 (80%)	106 (92%)	9 (8%)	12	2
All	All	223/288 (77%)	207 (93%)	16 (7%)	14	3

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	524	MET
1	A	620	LEU
1	A	634	ILE
1	A	637	SER
1	A	668	LYS
1	A	670	GLU
1	A	671	ARG
1	B	1531	GLN
1	B	1565	VAL
1	B	1595	LYS
1	B	1616	LYS
1	B	1620	LEU
1	B	1631	ASP
1	B	1651	LYS
1	B	1653	GLU
1	B	1671	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	594	HIS
1	A	604	ASN
1	A	608	ASN
1	A	664	ASN
1	B	1566	HIS
1	B	1604	ASN
1	B	1664	ASN

3.3.3 RNA [i](#)

There are no RNA molecules in this entry.

3.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

3.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

3.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 4 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	PO4	B	2681	-	4,4,4	0.50	0	6,6,6	0.36	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

3.7 Other polymers [i](#)

There are no such residues in this entry.

3.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

4 Fit of model and data [i](#)

4.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	149/166 (89%)	-0.41	0 100 100	18, 27, 44, 58	2 (1%)
1	B	148/166 (89%)	-0.49	1 (0%) 87 88	16, 26, 43, 69	1 (0%)
All	All	297/332 (89%)	-0.45	1 (0%) 94 93	16, 26, 44, 69	3 (1%)

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	1539	GLU	3.4

4.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

4.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

4.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	PO4	B	2681	5/5	0.98	0.12	28,31,53,62	0
2	CA	B	2679	1/1	0.99	0.07	21,21,21,21	0
2	CA	A	1680	1/1	0.99	0.05	18,18,18,18	0
2	CA	B	2680	1/1	1.00	0.07	20,20,20,20	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	CA	A	1679	1/1	1.00	0.08	18,18,18,18	0

4.5 Other polymers [i](#)

There are no such residues in this entry.