



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 31, 2023 – 11:06 AM JST

PDB ID : 5CN1
Title : Crystal structure of yeast GGA1_GAE domain-P21
Authors : Zhang, F.; Song, Y.; Li, X.; Teng, M.K.
Deposited on : 2015-07-17
Resolution : 2.31 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.31 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5974 (2.34-2.30)
Clashscore	141614	6604 (2.34-2.30)
Ramachandran outliers	138981	6523 (2.34-2.30)
Sidechain outliers	138945	6523 (2.34-2.30)
RSRZ outliers	127900	5855 (2.34-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	125	 67% 21% 10%
1	B	125	 71% 18% 10%
1	C	125	 71% 17% 10%
1	D	125	 70% 16% 6% 9%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 3441 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ADP-ribosylation factor-binding protein GGA1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	113	Total 851	C 530	N 148	O 170	S 3	0	0	0
1	B	113	Total 834	C 518	N 146	O 168	S 2	0	0	0
1	C	112	Total 847	C 527	N 149	O 168	S 3	0	0	0
1	D	114	Total 860	C 537	N 151	O 169	S 3	0	0	0

- Molecule 2 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	8	Total 8	O 8	0	0
2	B	18	Total 18	O 18	0	0
2	C	12	Total 12	O 12	0	0
2	D	11	Total 11	O 11	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: ADP-ribosylation factor-binding protein GGA1

Chain A: 



- Molecule 1: ADP-ribosylation factor-binding protein GGA1

Chain B: 



- Molecule 1: ADP-ribosylation factor-binding protein GGA1

Chain C: 



- Molecule 1: ADP-ribosylation factor-binding protein GGA1

Chain D: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	67.01Å 53.59Å 71.55Å 90.00° 103.57° 90.00°	Depositor
Resolution (Å)	42.48 – 2.31 42.45 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.0 (42.48-2.31) 99.0 (42.45-2.30)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.70 (at 2.32Å)	Xtrriage
Refinement program	REFMAC 5.7.0032	Depositor
R, R_{free}	0.212 , 0.270 0.216 , 0.272	Depositor DCC
R_{free} test set	1124 reflections (5.13%)	wwPDB-VP
Wilson B-factor (Å ²)	40.2	Xtrriage
Anisotropy	0.367	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 41.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3441	wwPDB-VP
Average B, all atoms (Å ²)	41.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 20.76 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 8.1124e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.56	0/862	0.81	1/1170 (0.1%)
1	B	0.59	0/847	0.80	1/1152 (0.1%)
1	C	0.57	0/859	0.79	1/1163 (0.1%)
1	D	0.57	0/871	0.81	0/1179
All	All	0.57	0/3439	0.80	3/4664 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	1

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	487	LEU	CA-CB-CG	5.60	128.18	115.30
1	B	487	LEU	CA-CB-CG	5.12	127.08	115.30
1	A	541	THR	CB-CA-C	-5.06	97.93	111.60

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	443	GLN	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	851	0	826	16	0
1	B	834	0	783	10	0
1	C	847	0	827	14	0
1	D	860	0	846	16	0
2	A	8	0	0	0	0
2	B	18	0	0	0	0
2	C	12	0	0	0	0
2	D	11	0	0	0	0
All	All	3441	0	3282	49	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (49) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:483:ASN:HB3	1:B:544:ASP:HA	1.65	0.78
1:C:523:ILE:O	1:C:525:VAL:O	2.09	0.71
1:A:461:ARG:NH2	1:A:464:MET:O	2.28	0.65
1:D:494:THR:HG21	1:D:520:VAL:HG13	1.80	0.64
1:A:489:ALA:HB2	1:C:487:LEU:HD22	1.80	0.62
1:A:544:ASP:OD2	1:D:551:THR:HG21	2.00	0.62
1:D:447:LEU:HD21	1:D:535:TRP:NE1	2.14	0.62
1:D:488:VAL:HG13	1:D:498:LEU:HD11	1.82	0.60
1:A:480:PRO:O	1:A:541:THR:HG23	2.00	0.60
1:D:488:VAL:HG21	1:D:533:LEU:HD21	1.84	0.59
1:B:487:LEU:HD22	1:D:489:ALA:HB2	1.84	0.59
1:C:494:THR:CG2	1:C:520:VAL:HG13	2.36	0.56
1:B:485:ALA:HA	1:B:501:GLN:HE22	1.74	0.53
1:A:539:TYR:OH	1:A:548:GLU:OE2	2.24	0.52
1:D:494:THR:HG21	1:D:520:VAL:CG1	2.39	0.52
1:B:475:ASN:O	1:B:509:ASN:HA	2.11	0.50
1:B:499:GLN:HA	1:D:495:THR:HB	1.94	0.49
1:C:454:ARG:NH1	1:C:456:ASP:OD2	2.43	0.49
1:C:478:SER:C	1:C:508:SER:HG	2.16	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:475:ASN:HB2	1:D:506:LEU:HB3	1.95	0.48
1:A:488:VAL:CG1	1:A:533:LEU:HD22	2.44	0.47
1:C:494:THR:HG23	1:C:520:VAL:HG13	1.95	0.47
1:A:535:TRP:HE1	1:A:552:THR:HG23	1.80	0.47
1:A:533:LEU:O	1:A:551:THR:HA	2.15	0.47
1:A:534:LYS:HG3	1:A:551:THR:HG22	1.97	0.47
1:C:491:PRO:O	1:C:494:THR:HB	2.16	0.46
1:C:478:SER:C	1:C:508:SER:OG	2.54	0.46
1:D:504:ASN:ND2	1:D:504:ASN:C	2.70	0.46
1:A:531:ILE:HD11	1:A:554:LEU:HD12	1.98	0.45
1:C:494:THR:HG21	1:C:520:VAL:HG13	1.98	0.45
1:D:481:ILE:HG12	1:D:541:THR:OG1	2.17	0.45
1:B:449:GLN:OE1	1:B:454:ARG:NH1	2.48	0.45
1:C:456:ASP:OD1	1:C:472:PHE:HB2	2.17	0.44
1:A:491:PRO:O	1:A:494:THR:HG23	2.17	0.43
1:D:488:VAL:CG2	1:D:533:LEU:HD21	2.47	0.43
1:A:488:VAL:HG13	1:A:533:LEU:HD22	2.01	0.43
1:D:488:VAL:HG22	1:D:496:LEU:CD2	2.49	0.43
1:C:457:TYR:CD2	1:C:469:LEU:CD2	3.02	0.42
1:C:538:ASN:HB3	1:D:545:SER:HB3	2.00	0.42
1:C:525:VAL:O	1:C:526:ASN:CB	2.67	0.42
1:D:447:LEU:CD2	1:D:552:THR:HG21	2.49	0.42
1:C:498:LEU:HD22	1:C:516:GLN:OE1	2.20	0.42
1:B:539:TYR:OH	1:B:548:GLU:OE2	2.11	0.42
1:A:465:THR:O	1:A:521:GLU:HA	2.20	0.42
1:B:489:ALA:HB2	1:D:487:LEU:HD22	2.01	0.42
1:A:445:HIS:O	1:A:456:ASP:HB2	2.20	0.41
1:B:499:GLN:HB3	1:B:500:PRO:CD	2.51	0.41
1:A:487:LEU:HD12	1:A:487:LEU:N	2.36	0.40
1:B:481:ILE:HB	1:B:506:LEU:HB2	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	111/125 (89%)	108 (97%)	3 (3%)	0	100	100
1	B	111/125 (89%)	107 (96%)	4 (4%)	0	100	100
1	C	108/125 (86%)	102 (94%)	6 (6%)	0	100	100
1	D	112/125 (90%)	108 (96%)	4 (4%)	0	100	100
All	All	442/500 (88%)	425 (96%)	17 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	92/113 (81%)	84 (91%)	8 (9%)	10	12
1	B	87/113 (77%)	77 (88%)	10 (12%)	5	6
1	C	94/113 (83%)	87 (93%)	7 (7%)	13	17
1	D	94/113 (83%)	79 (84%)	15 (16%)	2	2
All	All	367/452 (81%)	327 (89%)	40 (11%)	6	7

All (40) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	443	GLN
1	A	494	THR
1	A	504	ASN
1	A	525	VAL
1	A	527	LEU
1	A	541	THR
1	A	547	GLU
1	A	552	THR
1	B	443	GLN
1	B	453	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	459	LEU
1	B	487	LEU
1	B	495	THR
1	B	496	LEU
1	B	498	LEU
1	B	504	ASN
1	B	524	SER
1	B	552	THR
1	C	443	GLN
1	C	459	LEU
1	C	465	THR
1	C	487	LEU
1	C	546	LYS
1	C	551	THR
1	C	554	LEU
1	D	443	GLN
1	D	447	LEU
1	D	461	ARG
1	D	469	LEU
1	D	487	LEU
1	D	488	VAL
1	D	490	SER
1	D	495	THR
1	D	504	ASN
1	D	509	ASN
1	D	510	SER
1	D	521	GLU
1	D	527	LEU
1	D	533	LEU
1	D	551	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (17) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	449	GLN
1	A	501	GLN
1	A	504	ASN
1	A	516	GLN
1	B	443	GLN
1	B	499	GLN
1	B	501	GLN
1	B	504	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	509	ASN
1	B	516	GLN
1	C	443	GLN
1	C	501	GLN
1	C	507	GLN
1	D	443	GLN
1	D	501	GLN
1	D	504	ASN
1	D	516	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	113/125 (90%)	0.06	0	100 100	25, 40, 59, 72	0
1	B	113/125 (90%)	0.01	0	100 100	21, 37, 65, 70	0
1	C	112/125 (89%)	0.05	1 (0%)	84 88	26, 41, 60, 69	0
1	D	114/125 (91%)	0.04	1 (0%)	84 88	24, 43, 64, 71	0
All	All	452/500 (90%)	0.04	2 (0%)	92 95	21, 41, 63, 72	0

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	460	THR	2.3
1	D	528	GLY	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.