wwPDB EM Validation Summary Report

Nov 6, 2022 – 09:28 PM EST

PDB ID	:	6CP7
EMDB ID	:	EMD-7549
Title	:	Monomer yeast ATP synthase Fo reconstituted in nanodisc generated from masked refinement.
Authors	:	Srivastava, A.P.; Luo, M.; Symersky, J.; Liao, M.F.; Mueller, D.M.
Deposited on	:	2018-03-13
Resolution	:	4.10 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/EMValidationReportHelp
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references ⓘ) were used in the production of this report:

EMDB validation analysis	:	0.0.1.dev43
Mogul	:	1.8.5 (274361), CSD as541be (2020)
MolProbity	:	4.02b-467
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
MapQ	:	1.9.9
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.31.2

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 4.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion < 40%). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	K	76	49% 64% 32%
1	L	76	39% 49% 42% 7%
1	M	76	39% 57% 39%
1	N	76	16% 75% 21%
1	O	76	46% 61% 34%
1	P	76	36% 80% 14%
1	Q	76	41% 63% 28% 5%
1	R	76	41% 63% 32%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	S	76	
1	T	76	
2	8	48	
3	X	249	
4	Z	209	
5	7	173	
6	U	95	
7	J	37	

2 Entry composition (i)

There are 7 unique types of molecules in this entry. The entry contains 9164 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ATP synthase subunit 9, mitochondrial.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	K	74	Total	C	N	O	S	0	0
			534	358	82	90	4		
1	L	74	Total	C	N	O	S	0	0
			529	353	82	91	3		
1	M	75	Total	C	N	O	S	0	0
			535	358	83	91	3		
1	N	74	Total	C	N	O	S	0	0
			532	356	82	91	3		
1	O	74	Total	C	N	O	S	0	0
			533	357	82	91	3		
1	P	74	Total	C	N	O	S	0	0
			529	352	82	91	4		
1	Q	74	Total	C	N	O	S	0	0
			522	347	82	90	3		
1	R	75	Total	C	N	O	S	0	0
			534	356	83	92	3		
1	S	74	Total	C	N	O	S	0	0
			532	356	82	91	3		
1	T	74	Total	C	N	O	S	0	0
			533	357	82	91	3		

- Molecule 2 is a protein called ATP synthase protein 8.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	8	42	Total	C	N	O	S	0	0
			364	256	52	53	3		

- Molecule 3 is a protein called ATP synthase subunit a.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	X	224	Total	C	N	O	S	0	0
			1760	1198	265	287	10		

- Molecule 4 is a protein called ATP synthase subunit 4, mitochondrial.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	Z	54	Total	C	N	O	S	0	0
			393	253	64	75	1		

- Molecule 5 is a protein called ATP synthase subunit d, mitochondrial.

Mol	Chain	Residues	Atoms					AltConf	Trace
5	7	67	Total	C	N	O	S	0	0
			520	329	86	103	2		

- Molecule 6 is a protein called ATP synthase subunit f, mitochondrial.

Mol	Chain	Residues	Atoms					AltConf	Trace
6	U	68	Total	C	N	O	S	0	0
			522	346	89	86	1		

- Molecule 7 is a protein called ATP synthase subunit J, mitochondrial.

Mol	Chain	Residues	Atoms					AltConf	Trace
7	J	37	Total	C	N	O	S	0	0
			292	197	45	48	2		

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: ATP synthase subunit 9, mitochondrial

Chain K:



• Molecule 1: ATP synthase subunit 9, mitochondrial

Chain L:



• Molecule 1: ATP synthase subunit 9, mitochondrial

Chain M:



• Molecule 1: ATP synthase subunit 9, mitochondrial

Chain N:





• Molecule 1: ATP synthase subunit 9, mitochondrial

Chain O:





• Molecule 1: ATP synthase subunit 9, mitochondrial

Chain P:



• Molecule 1: ATP synthase subunit 9, mitochondrial

Chain Q:





• Molecule 1: ATP synthase subunit 9, mitochondrial

Chain R:



• Molecule 1: ATP synthase subunit 9, mitochondrial

Chain S:





• Molecule 1: ATP synthase subunit 9, mitochondrial

Chain T:



• Molecule 2: ATP synthase protein 8

Chain 8:



• Molecule 3: ATP synthase subunit a

Chain X:







• Molecule 4: ATP synthase subunit 4, mitochondrial

Chain Z:





• Molecule 5: ATP synthase subunit d, mitochondrial

Chain 7:



• Molecule 6: ATP synthase subunit f, mitochondrial

Chain U:



• Molecule 7: ATP synthase subunit J, mitochondrial

Chain J:





4 Experimental information (i)

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C1	Depositor
Number of particles used	109206	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI POLARA 300	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/Å^2$)	8	Depositor
Minimum defocus (nm)	1000	Depositor
Maximum defocus (nm)	2600	Depositor
Magnification	31000	Depositor
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor
Maximum map value	0.180	Depositor
Minimum map value	-0.101	Depositor
Average map value	-0.000	Depositor
Map value standard deviation	0.006	Depositor
Recommended contour level	0.0475	Depositor
Map size (Å)	393.6, 393.6, 393.6	wwPDB
Map dimensions	320, 320, 320	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.23, 1.23, 1.23	Depositor

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: FME

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#$\|Z\|>5$	RMSZ	#$\|Z\|>5$
1	K	0.45	0/532	0.74	0/721
1	L	0.40	0/530	0.72	0/718
1	M	0.41	0/535	0.77	2/724 (0.3%)
1	N	0.40	0/533	0.79	1/722 (0.1%)
1	O	0.41	0/533	0.81	2/722 (0.3%)
1	P	0.41	0/527	0.78	2/714 (0.3%)
1	Q	0.38	0/523	0.72	0/709
1	R	0.42	0/534	0.76	0/723
1	S	0.42	0/533	0.83	3/722 (0.4%)
1	T	0.45	0/533	0.79	1/722 (0.1%)
2	8	0.42	0/374	0.72	0/503
3	X	0.35	0/1807	0.65	0/2466
4	Z	0.31	0/398	0.56	0/538
5	7	0.33	0/527	0.57	1/709 (0.1%)
6	U	0.32	0/540	0.58	0/731
7	J	0.35	0/302	0.61	0/410
All	All	0.39	0/9261	0.71	12/12554 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand.A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	R	0	1
6	U	0	1
All	All	0	2

There are no bond length outliers.

The worst 5 of 12 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	P	20	LEU	CA-CB-CG	7.02	131.44	115.30
1	O	57	LEU	CA-CB-CG	6.65	130.59	115.30
1	N	57	LEU	CA-CB-CG	6.04	129.20	115.30
1	S	53	LEU	CA-CB-CG	5.79	128.62	115.30
1	S	66	LEU	CA-CB-CG	5.76	128.55	115.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	R	19	LEU	Peptide
6	U	75	ALA	Peptide

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	K	534	0	575	30	0
1	L	529	0	562	45	0
1	M	535	0	571	28	0
1	N	532	0	571	13	0
1	O	533	0	573	23	0
1	P	529	0	560	9	0
1	Q	522	0	541	33	0
1	R	534	0	567	33	0
1	S	532	0	571	34	0
1	T	533	0	573	17	0
2	8	364	0	390	9	0
3	X	1760	0	1847	48	0
4	Z	393	0	377	0	0
5	7	520	0	511	8	0
6	U	522	0	483	14	0
7	J	292	0	298	3	0
All	All	9164	0	9570	262	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

The worst 5 of 262 close contacts within the same asymmetric unit are listed below, sorted by

their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:R:58:SER:HB3	1:S:26:ILE:CD1	1.39	1.51
1:Q:18:GLY:HA3	1:Q:65:CYS:SG	1.56	1.43
1:R:58:SER:CB	1:S:26:ILE:HD11	1.54	1.35
3:X:164:LEU:O	3:X:168:ALA:N	1.58	1.34
1:Q:24:ILE:HG22	1:Q:25:GLY:N	1.37	1.24

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	K	72/76 (95%)	66 (92%)	6 (8%)	0	100	100
1	L	72/76 (95%)	67 (93%)	4 (6%)	1 (1%)	11	45
1	M	73/76 (96%)	65 (89%)	7 (10%)	1 (1%)	11	45
1	N	72/76 (95%)	68 (94%)	4 (6%)	0	100	100
1	O	72/76 (95%)	65 (90%)	6 (8%)	1 (1%)	11	45
1	P	72/76 (95%)	70 (97%)	2 (3%)	0	100	100
1	Q	72/76 (95%)	68 (94%)	1 (1%)	3 (4%)	3	24
1	R	73/76 (96%)	68 (93%)	5 (7%)	0	100	100
1	S	72/76 (95%)	67 (93%)	5 (7%)	0	100	100
1	T	72/76 (95%)	67 (93%)	5 (7%)	0	100	100
2	8	40/48 (83%)	38 (95%)	2 (5%)	0	100	100
3	X	222/249 (89%)	204 (92%)	18 (8%)	0	100	100
4	Z	52/209 (25%)	52 (100%)	0	0	100	100
5	7	65/173 (38%)	63 (97%)	2 (3%)	0	100	100
6	U	66/95 (70%)	57 (86%)	9 (14%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
7	J	35/37 (95%)	32 (91%)	3 (9%)	0	100 100
All	All	1202/1571 (76%)	1117 (93%)	79 (7%)	6 (0%)	32 67

5 of 6 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	M	59	GLU
1	Q	25	GLY
1	Q	24	ILE
1	L	29	VAL
1	Q	17	ILE

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	K	53/55 (96%)	52 (98%)	1 (2%)	57 75
1	L	53/55 (96%)	45 (85%)	8 (15%)	3 17
1	M	53/55 (96%)	48 (91%)	5 (9%)	8 31
1	N	54/55 (98%)	52 (96%)	2 (4%)	34 59
1	O	54/55 (98%)	50 (93%)	4 (7%)	13 41
1	P	52/55 (94%)	47 (90%)	5 (10%)	8 30
1	Q	50/55 (91%)	45 (90%)	5 (10%)	7 29
1	R	53/55 (96%)	49 (92%)	4 (8%)	13 40
1	S	54/55 (98%)	53 (98%)	1 (2%)	57 75
1	T	54/55 (98%)	53 (98%)	1 (2%)	57 75
2	8	41/47 (87%)	41 (100%)	0	100 100
3	X	190/217 (88%)	185 (97%)	5 (3%)	46 67
4	Z	37/182 (20%)	37 (100%)	0	100 100
5	7	56/158 (35%)	54 (96%)	2 (4%)	35 60
6	U	45/76 (59%)	45 (100%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
7	J	30/30 (100%)	30 (100%)	0	100 100
All	All	929/1260 (74%)	886 (95%)	43 (5%)	31 54

5 of 43 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	Q	66	LEU
1	T	67	MET
1	Q	67	MET
1	R	66	LEU
3	X	163	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
3	X	71	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

10 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
1	FME	N	1	1	5,6,10	1.41	1 (20%)	3,6,11	1.85	1 (33%)
1	FME	K	1	1	8,9,10	0.95	0	7,9,11	0.90	0
1	FME	O	1	1	6,7,10	0.94	0	6,7,11	1.33	1 (16%)
1	FME	P	1	1	8,9,10	0.91	0	7,9,11	0.85	0
1	FME	R	1	1	6,7,10	1.02	0	6,7,11	1.24	1 (16%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\|>2$	Counts	RMSZ	$\#\|Z\|>2$
1	FME	S	1	1	5,6,10	1.51	1 (20%)	3,6,11	1.78	1 (33%)
1	FME	L	1	1	5,6,10	1.30	1 (20%)	3,6,11	1.65	1 (33%)
1	FME	T	1	1	6,7,10	0.97	0	6,7,11	1.26	1 (16%)
1	FME	Q	1	1	5,6,10	1.33	1 (20%)	3,6,11	1.51	1 (33%)
1	FME	M	1	1	6,7,10	0.98	0	6,7,11	1.47	1 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	FME	N	1	1	-	0/2/5/11	-
1	FME	K	1	1	-	6/7/9/11	-
1	FME	O	1	1	-	3/5/7/11	-
1	FME	P	1	1	-	4/7/9/11	-
1	FME	R	1	1	-	1/5/7/11	-
1	FME	S	1	1	-	0/2/5/11	-
1	FME	L	1	1	-	0/2/5/11	-
1	FME	T	1	1	-	3/5/7/11	-
1	FME	Q	1	1	-	0/2/5/11	-
1	FME	M	1	1	-	3/5/7/11	-

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	S	1	FME	CA-N	-2.73	1.43	1.46
1	N	1	FME	CA-N	-2.48	1.44	1.46
1	Q	1	FME	CA-N	-2.10	1.44	1.46
1	L	1	FME	CA-N	-2.07	1.44	1.46

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	S	1	FME	CB-CA-N	2.97	112.91	109.61
1	N	1	FME	CB-CA-N	2.96	112.90	109.61
1	M	1	FME	CB-CA-C	-2.85	107.21	111.29
1	O	1	FME	CB-CA-C	-2.70	107.42	111.29
1	L	1	FME	CB-CA-N	2.53	112.43	109.61

There are no chirality outliers.

5 of 20 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	K	1	FME	O-C-CA-CB
1	K	1	FME	CA-CB-CG-SD
1	M	1	FME	C-CA-CB-CG
1	O	1	FME	CB-CA-N-CN
1	O	1	FME	C-CA-CB-CG

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	L	1	FME	1	0

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

There are no ligands in this entry.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Map visualisation

This section contains visualisations of the EMDB entry EMD-7549. These allow visual inspection of the internal detail of the map and identification of artifacts.

No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections

6.1.1 Primary map



X



Y



Z

The images above show the map projected in three orthogonal directions.

6.2 Central slices

6.2.1 Primary map



X Index: 160



Y Index: 160



Z Index: 160

The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices (i)

6.3.1 Primary map



X Index: 159



Y Index: 145



Z Index: 115

The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal surface views (i)

6.4.1 Primary map

X

Y

Z

The images above show the 3D surface view of the map at the recommended contour level 0.0475. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5 Mask visualisation (i)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis

This section contains the results of statistical analysis of the map.

7.1 Map-value distribution



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.

7.2 Volume estimate (i)



The volume at the recommended contour level is 51 nm^3; this corresponds to an approximate mass of 47 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum (i)



*Reported resolution corresponds to spatial frequency of 0.244 Å^{-1}

8 Fourier-Shell correlation

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC



*Reported resolution corresponds to spatial frequency of 0.244 Å^{-1}

8.2 Resolution estimates (i)

Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	4.10	-	-
Author-provided FSC curve	4.04	5.19	4.08
Unmasked-calculated*	-	-	-

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps.

9 Map-model fit

This section contains information regarding the fit between EMDB map EMD-7549 and PDB model 6CP7. Per-residue inclusion information can be found in section 3 on page 6.

9.1 Map-model overlay



X



Y



Z

The images above show the 3D surface view of the map at the recommended contour level 0.0475 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model (i)



X Y Z

The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model (i)



X Y Z

The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.0475).

9.4 Atom inclusion (i)



At the recommended contour level, 59% of all backbone atoms, 38% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.0475) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.3783	0.3110
7	0.3774	0.2710
8	0.3473	0.2800
J	0.2203	0.2830
K	0.4143	0.3190
L	0.4734	0.3450
M	0.4756	0.3540
N	0.5728	0.3570
O	0.4321	0.3360
P	0.4867	0.3450
Q	0.4605	0.3410
R	0.4444	0.3530
S	0.4329	0.3700
T	0.4943	0.3820
U	0.1468	0.2320
X	0.2444	0.2540
Z	0.2005	0.2410