



Full wwPDB X-ray Structure Validation Report ⓘ

May 22, 2020 – 04:26 pm BST

PDB ID : 3CQH
Title : Crystal Structure of L-xylulose-5-phosphate 3-epimerase UlaE from the Anaerobic L-ascorbate Utilization Pathway of Escherichia coli
Authors : Shi, R.; Matte, A.; Cygler, M.; Montreal-Kingston Bacterial Structural Genomics Initiative (BSGI)
Deposited on : 2008-04-03
Resolution : 2.08 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.08 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	6189 (2.10-2.06)
Clashscore	141614	6738 (2.10-2.06)
Ramachandran outliers	138981	6663 (2.10-2.06)
Sidechain outliers	138945	6664 (2.10-2.06)
RSRZ outliers	127900	6057 (2.10-2.06)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	295	 11% 71% 14% • 13%
1	B	295	 10% 71% 13% • 15%

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 4235 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called L-ribulose-5-phosphate 3-epimerase ulaE.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	257	2042	1293	357	377	5	10	0	1	0
1	B	252	1999	1264	350	373	4	8	0	1	0

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-10	MSE	-	expression tag	UNP Q8XDI5
A	-9	GLY	-	expression tag	UNP Q8XDI5
A	-8	SER	-	expression tag	UNP Q8XDI5
A	-7	SER	-	expression tag	UNP Q8XDI5
A	-6	HIS	-	expression tag	UNP Q8XDI5
A	-5	HIS	-	expression tag	UNP Q8XDI5
A	-4	HIS	-	expression tag	UNP Q8XDI5
A	-3	HIS	-	expression tag	UNP Q8XDI5
A	-2	HIS	-	expression tag	UNP Q8XDI5
A	-1	HIS	-	expression tag	UNP Q8XDI5
A	0	GLY	-	expression tag	UNP Q8XDI5
A	1	SER	-	expression tag	UNP Q8XDI5
B	-10	MSE	-	expression tag	UNP Q8XDI5
B	-9	GLY	-	expression tag	UNP Q8XDI5
B	-8	SER	-	expression tag	UNP Q8XDI5
B	-7	SER	-	expression tag	UNP Q8XDI5
B	-6	HIS	-	expression tag	UNP Q8XDI5
B	-5	HIS	-	expression tag	UNP Q8XDI5
B	-4	HIS	-	expression tag	UNP Q8XDI5
B	-3	HIS	-	expression tag	UNP Q8XDI5
B	-2	HIS	-	expression tag	UNP Q8XDI5
B	-1	HIS	-	expression tag	UNP Q8XDI5
B	0	GLY	-	expression tag	UNP Q8XDI5
B	1	SER	-	expression tag	UNP Q8XDI5

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	91	Total	O	0	0
			91	91		
3	B	78	Total	O	0	0
			78	78		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: L-ribulose-5-phosphate 3-epimerase ulaE



- Molecule 1: L-ribulose-5-phosphate 3-epimerase ulaE



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	110.39Å 110.39Å 103.02Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 2.08 33.06 – 2.08	Depositor EDS
% Data completeness (in resolution range)	99.8 (50.00-2.08) 99.8 (33.06-2.08)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.38 (at 2.08Å)	Xtrriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.212 , 0.242 0.211 , 0.239	Depositor DCC
R_{free} test set	1939 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	34.7	Xtrriage
Anisotropy	0.247	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 52.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4235	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.01% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.58	0/2074	0.65	1/2784 (0.0%)
1	B	0.54	0/2029	0.66	2/2724 (0.1%)
All	All	0.56	0/4103	0.65	3/5508 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	B	112	ARG	NE-CZ-NH2	-11.50	114.55	120.30
1	B	112	ARG	NE-CZ-NH1	6.13	123.37	120.30
1	A	249	LEU	CA-CB-CG	-5.01	103.77	115.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	5	GLN	Peptide

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2042	0	2022	41	1
1	B	1999	0	1974	36	0
2	A	10	0	0	0	0
2	B	15	0	0	1	0
3	A	91	0	0	2	0
3	B	78	0	0	6	0
All	All	4235	0	3996	74	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (74) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:10:ILE:HD11	1:A:252:MSE:CE	1.72	1.20
1:A:5:GLN:N	1:B:147:ARG:HD3	1.76	1.01
1:A:149:GLN:O	1:B:112:ARG:HD2	1.62	0.99
1:B:225:PHE:HE2	1:B:252:MSE:CE	1.75	0.98
1:A:115:GLN:HE22	1:A:211:HIS:CE1	1.85	0.95
1:A:10:ILE:HD11	1:A:252:MSE:HE3	1.51	0.92
1:A:10:ILE:CD1	1:A:252:MSE:CE	2.47	0.91
1:A:10:ILE:CD1	1:A:252:MSE:HE2	2.13	0.78
1:B:6:ILE:HD11	1:B:275:MSE:SE	2.35	0.76
1:A:10:ILE:HD11	1:A:252:MSE:HE2	1.70	0.73
1:A:115:GLN:NE2	1:A:211:HIS:CE1	2.57	0.72
1:B:252:MSE:HE1	1:B:267:ALA:CB	2.22	0.70
1:B:225:PHE:CE2	1:B:252:MSE:CE	2.67	0.68
1:A:30:THR:HG22	3:A:451:HOH:O	1.93	0.68
1:B:252:MSE:HE1	1:B:267:ALA:HB1	1.76	0.68
1:B:225:PHE:HE2	1:B:252:MSE:HE1	1.56	0.67
1:B:225:PHE:HE2	1:B:252:MSE:HE3	1.62	0.65
1:A:217:PRO:HD3	1:A:229:VAL:HG11	1.79	0.64
1:A:259:ASP:HB2	3:A:465:HOH:O	2.00	0.62
1:A:104:GLN:HA	1:A:107:GLN:HE21	1.66	0.61
1:A:42:GLU:OE2	1:A:80:ARG:HB2	2.01	0.60

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:10:ILE:HD11	1:A:252:MSE:HE1	1.78	0.60
1:A:149:GLN:O	1:B:112:ARG:CD	2.45	0.60
1:B:16:PRO:HG3	1:B:260:PRO:HG2	1.85	0.59
1:B:225:PHE:CE2	1:B:252:MSE:HE1	2.34	0.58
1:B:221:LYS:HE3	1:B:253:TRP:CD2	2.38	0.58
1:B:147:ARG:NH2	3:B:478:HOH:O	2.39	0.56
1:A:189:LEU:HB3	1:A:196:VAL:HG13	1.88	0.56
1:A:6:ILE:HG13	1:A:6:ILE:O	2.06	0.55
1:B:80:ARG:HG3	3:B:479:HOH:O	2.07	0.55
1:B:6:ILE:CD1	1:B:275:MSE:SE	3.05	0.55
1:A:218:GLY:HA2	1:A:220:PHE:CE1	2.44	0.53
1:B:99:MSE:HE1	1:B:116:LEU:HD22	1.91	0.53
1:A:217:PRO:HD3	1:A:229:VAL:CG1	2.40	0.52
1:A:156:ILE:HG21	1:A:199:GLU:HB3	1.91	0.51
1:B:248:TYR:OH	1:B:280:MSE:HE1	2.11	0.51
1:B:204:ILE:HG22	3:B:442:HOH:O	2.10	0.50
1:B:30:THR:HG23	3:B:452:HOH:O	2.11	0.50
1:A:253:TRP:O	1:A:256:THR:HG22	2.13	0.49
1:B:204:ILE:HG12	1:B:242:SER:HB3	1.94	0.49
1:A:49:ARG:HA	1:A:52:TRP:CD1	2.49	0.48
1:A:156:ILE:HG12	1:A:182:LEU:HD13	1.96	0.48
1:A:156:ILE:O	1:A:157:MSE:CB	2.62	0.47
1:A:35:PHE:HB2	1:A:72:PRO:HD3	1.96	0.47
1:A:190:SER:HA	1:A:196:VAL:HG21	1.97	0.47
1:B:221:LYS:HE3	1:B:253:TRP:CE2	2.51	0.46
1:A:188:ASN:OD1	1:A:220:PHE:HB3	2.16	0.46
1:A:156:ILE:O	1:A:157:MSE:HB2	2.15	0.46
1:A:204:ILE:HD11	1:A:244:TYR:HB2	1.98	0.45
1:A:219:VAL:O	1:A:219:VAL:HG12	2.16	0.45
1:B:229:VAL:O	1:B:229:VAL:HG22	2.16	0.45
1:B:49:ARG:HA	1:B:52:TRP:CD1	2.52	0.45
1:B:142:VAL:HG21	1:B:175:LEU:HD13	1.97	0.45
1:A:156:ILE:HA	1:A:182:LEU:HD22	1.99	0.45
1:B:185:ASP:HB2	3:B:426:HOH:O	2.17	0.44
1:A:221:LYS:HE3	1:A:253:TRP:CD2	2.52	0.44
1:B:236:PHE:O	1:B:280:MSE:HE3	2.18	0.44
1:A:191:ALA:HB1	1:A:218:GLY:H	1.83	0.43
1:B:109:VAL:HG23	1:B:111:ILE:HG13	2.01	0.43
1:B:241:GLN:HG2	3:B:450:HOH:O	2.19	0.42
1:A:141:SER:HA	1:A:144:MSE:HE2	2.01	0.42
1:B:43:THR:O	1:B:47:LEU:HB2	2.20	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:216:LYS:O	1:A:217:PRO:C	2.57	0.41
1:A:83:PRO:HG2	1:A:86:SER:HB2	2.03	0.41
1:A:14:ALA:CB	1:A:252:MSE:HE3	2.51	0.41
1:A:14:ALA:HB1	1:A:252:MSE:HE3	2.02	0.41
1:A:5:GLN:HB2	1:A:6:ILE:H	1.74	0.41
1:B:225:PHE:CE2	1:B:252:MSE:HE3	2.50	0.41
1:A:10:ILE:CD1	1:A:252:MSE:HE1	2.44	0.41
1:B:6:ILE:HG12	1:B:6:ILE:O	2.20	0.40
1:B:274:ARG:HD3	1:B:274:ARG:HA	1.81	0.40
1:B:221:LYS:HB3	1:B:221:LYS:HE2	1.88	0.40
1:B:231[A]:ASP:HA	2:B:404:SO4:O3	2.22	0.40
1:B:37:GLU:HB2	1:B:249:LEU:HD13	2.03	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:217:PRO:O	1:A:258:GLU:OE1[3_544]	1.79	0.41

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	252/295 (85%)	246 (98%)	5 (2%)	1 (0%)	34	31
1	B	245/295 (83%)	238 (97%)	5 (2%)	2 (1%)	19	14
All	All	497/590 (84%)	484 (97%)	10 (2%)	3 (1%)	25	20

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	176	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	258	GLU
1	B	222	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	214/232 (92%)	203 (95%)	11 (5%)	24	21
1	B	209/232 (90%)	202 (97%)	7 (3%)	38	39
All	All	423/464 (91%)	405 (96%)	18 (4%)	29	28

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	5	GLN
1	A	6	ILE
1	A	42	GLU
1	A	44	ASP
1	A	87	GLU
1	A	124	GLN
1	A	176	ASN
1	A	183	TYR
1	A	199	GLU
1	A	220	PHE
1	A	221	LYS
1	B	6	ILE
1	B	81	ARG
1	B	89	ASP
1	B	115	GLN
1	B	156	ILE
1	B	183	TYR
1	B	222	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	107	GLN
1	A	115	GLN
1	B	62	ASN
1	B	176	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	A	405	-	4,4,4	0.17	0	6,6,6	0.27	0
2	SO4	B	403	-	4,4,4	0.13	0	6,6,6	0.15	0
2	SO4	B	401	-	4,4,4	0.14	0	6,6,6	0.26	0
2	SO4	B	404	-	4,4,4	0.17	0	6,6,6	0.16	0
2	SO4	A	402	-	4,4,4	0.18	0	6,6,6	0.14	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	404	SO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	247/295 (83%)	0.76	31 (12%) 3 4	20, 31, 47, 59	2 (0%)
1	B	243/295 (82%)	0.82	30 (12%) 4 5	22, 35, 52, 63	1 (0%)
All	All	490/590 (83%)	0.79	61 (12%) 4 5	20, 33, 51, 63	3 (0%)

All (61) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	219	VAL	7.9
1	B	219	VAL	6.7
1	B	256	THR	6.6
1	A	220	PHE	6.2
1	A	192	TRP	6.1
1	B	257	ALA	5.8
1	B	218	GLY	5.7
1	A	222	ASN	5.5
1	B	222	ASN	4.7
1	A	256	THR	4.2
1	A	258	GLU	4.1
1	A	217	PRO	4.1
1	A	90	ALA	4.0
1	B	89	ASP	3.9
1	B	17	ALA	3.9
1	A	175	LEU	3.8
1	B	217	PRO	3.7
1	A	257	ALA	3.6
1	B	202	ALA	3.6
1	B	156	ILE	3.6
1	A	113	VAL	3.4
1	A	114	ILE	3.3
1	B	175	LEU	3.2
1	B	72	PRO	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	220	PHE	3.1
1	B	129	GLU	3.1
1	A	218	GLY	3.1
1	B	227	GLU	2.9
1	B	258	GLU	2.9
1	A	176	ASN	2.9
1	B	90	ALA	2.9
1	A	202	ALA	2.8
1	B	91	VAL	2.7
1	B	87	GLU	2.7
1	A	201	GLN	2.7
1	B	196	VAL	2.7
1	A	216	LYS	2.6
1	B	221	LYS	2.5
1	A	87	GLU	2.5
1	A	187	GLY	2.5
1	B	197	GLN	2.5
1	A	150	VAL	2.4
1	A	42	GLU	2.4
1	A	111	ILE	2.4
1	A	73	SER	2.4
1	B	111	ILE	2.4
1	A	183	TYR	2.4
1	A	93	ALA	2.3
1	B	201	GLN	2.3
1	B	113	VAL	2.2
1	B	241	GLN	2.2
1	B	259	ASP	2.2
1	B	114	ILE	2.2
1	A	133	ARG	2.1
1	B	96	LEU	2.1
1	A	80	ARG	2.1
1	B	132	ARG	2.1
1	A	281	VAL	2.1
1	A	221	LYS	2.1
1	A	112	ARG	2.0
1	A	204	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	SO4	B	404	5/5	0.85	0.25	88,88,88,88	0
2	SO4	A	405	5/5	0.95	0.21	62,63,63,64	0
2	SO4	B	403	5/5	0.97	0.11	54,55,56,57	0
2	SO4	B	401	5/5	0.97	0.09	50,50,50,51	0
2	SO4	A	402	5/5	0.99	0.07	55,56,57,58	0

6.5 Other polymers [i](#)

There are no such residues in this entry.