



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 19, 2024 – 12:26 pm GMT

PDB ID : 8CRA
Title : Structure of the keratin-like domain of SEPALLATA3 and AGAMOUS from Arabidopsis thaliana
Authors : Zubieta, C.; Hugouvieux, V.
Deposited on : 2023-03-08
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtrriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3907 (2.40-2.40)
Clashscore	141614	4398 (2.40-2.40)
Ramachandran outliers	138981	4318 (2.40-2.40)
Sidechain outliers	138945	4319 (2.40-2.40)
RSRZ outliers	127900	3811 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	96	
1	B	96	
1	C	96	
1	D	96	
2	E	98	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	98	 <p>2% 85% 8% • 5%</p>
2	G	98	 <p>6% 69% 17% • 10%</p>
2	H	98	 <p>6% 80% 9% • 10%</p>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 6163 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Floral homeotic protein AGAMOUS.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	96	Total 788	C 480	N 151	O 154	S 3	0	0	0
1	B	91	Total 745	C 456	N 141	O 145	S 3	0	0	0
1	C	92	Total 758	C 463	N 144	O 148	S 3	0	0	0
1	D	93	Total 769	C 469	N 148	O 149	S 3	0	0	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	191	ARG	-	expression tag	UNP P17839
A	192	ASN	-	expression tag	UNP P17839
B	191	ARG	-	expression tag	UNP P17839
B	192	ASN	-	expression tag	UNP P17839
C	191	ARG	-	expression tag	UNP P17839
C	192	ASN	-	expression tag	UNP P17839
D	191	ARG	-	expression tag	UNP P17839
D	192	ASN	-	expression tag	UNP P17839

- Molecule 2 is a protein called Developmental protein SEPALLATA 3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	E	96	Total 787	C 485	N 145	O 155	S 2	0	0	0
2	F	93	Total 765	C 471	N 141	O 151	S 2	0	0	0
2	G	88	Total 727	C 449	N 133	O 143	S 2	0	0	0
2	H	88	Total 709	C 437	N 127	O 143	S 2	0	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	21	Total O 21 21	0	0
3	B	13	Total O 13 13	0	0
3	C	24	Total O 24 24	0	0
3	D	7	Total O 7 7	0	0
3	E	15	Total O 15 15	0	0
3	F	19	Total O 19 19	0	0
3	G	2	Total O 2 2	0	0
3	H	14	Total O 14 14	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Floral homeotic protein AGAMOUS




- Molecule 1: Floral homeotic protein AGAMOUS





- Molecule 1: Floral homeotic protein AGAMOUS





- Molecule 1: Floral homeotic protein AGAMOUS



- Molecule 2: Developmental protein SEPALLATA 3




- Molecule 2: Developmental protein SEPALLATA 3



- Molecule 2: Developmental protein SEPALLATA 3



- Molecule 2: Developmental protein SEPALLATA 3



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	101.33Å 138.37Å 180.23Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.40 180.23 – 2.40	Depositor EDS
% Data completeness (in resolution range)	99.4 (20.00-2.40) 99.6 (180.23-2.40)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.35 (at 2.40Å)	Xtrriage
Refinement program	REFMAC 5.8.0405	Depositor
R, R_{free}	0.237 , 0.279 0.244 , 0.279	Depositor DCC
R_{free} test set	2422 reflections (4.89%)	wwPDB-VP
Wilson B-factor (Å ²)	55.2	Xtrriage
Anisotropy	0.225	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 43.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	6163	wwPDB-VP
Average B, all atoms (Å ²)	83.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.75% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.43	0/793	0.70	0/1057
1	B	0.38	0/750	0.68	0/1002
1	C	0.39	0/763	0.64	0/1018
1	D	0.36	0/774	0.65	0/1032
2	E	0.38	0/792	0.68	0/1061
2	F	0.38	0/769	0.68	0/1028
2	G	0.31	0/730	0.61	0/976
2	H	0.44	0/712	0.72	1/954 (0.1%)
All	All	0.39	0/6083	0.67	1/8128 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	B	0	2
1	C	0	3
2	E	0	2
2	G	0	1
2	H	0	1
All	All	0	11

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	H	174	ARG	O-C-N	-8.24	109.52	122.70

There are no chirality outliers.

All (11) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	146	ARG	Sidechain
1	A	191	ARG	Sidechain
1	B	114	ARG	Sidechain
1	B	146	ARG	Sidechain
1	C	125	ARG	Sidechain
1	C	146	ARG	Sidechain
1	C	184	ARG	Sidechain
2	E	104	ARG	Sidechain
2	E	85	ARG	Sidechain
2	G	110	ARG	Sidechain
2	H	174	ARG	Mainchain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	788	0	798	13	0
1	B	745	0	755	7	2
1	C	758	0	769	6	0
1	D	769	0	782	7	0
2	E	787	0	818	5	2
2	F	765	0	796	8	0
2	G	727	0	759	14	0
2	H	709	0	719	7	0
3	A	21	0	0	4	0
3	B	13	0	0	0	0
3	C	24	0	0	0	0
3	D	7	0	0	0	0
3	E	15	0	0	0	0
3	F	19	0	0	0	0
3	G	2	0	0	1	0
3	H	14	0	0	0	0
All	All	6163	0	6196	53	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (53) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:189:GLU:O	1:A:192:ASN:HB2	1.56	1.05
2:F:174:ARG:O	2:F:175:LEU:HD22	1.78	0.83
1:A:189:GLU:O	1:A:192:ASN:CB	2.27	0.81
1:A:191:ARG:NH2	3:A:201:HOH:O	2.21	0.72
1:C:121:GLN:HB3	1:C:125:ARG:HH12	1.58	0.68
1:A:183:LEU:HB2	2:H:171:LEU:HD23	1.78	0.65
2:F:105:TYR:CE2	2:F:109:GLN:NE2	2.65	0.65
1:A:189:GLU:HG3	1:A:190:ASN:H	1.61	0.65
1:A:97:SER:N	3:A:202:HOH:O	2.29	0.64
1:C:165:GLU:HG3	2:G:154:LEU:HD11	1.82	0.62
1:D:128:MET:CE	2:G:134:GLN:HE22	2.12	0.62
1:C:177:HIS:CE1	2:G:164:LEU:HD11	2.37	0.59
3:A:215:HOH:O	2:E:125:THR:CG2	2.51	0.58
2:F:174:ARG:O	2:F:174:ARG:HG3	2.04	0.58
1:B:165:GLU:HG2	2:E:154:LEU:HD11	1.88	0.56
2:G:104:ARG:O	2:G:108:LEU:HD23	2.06	0.55
1:A:189:GLU:CG	1:A:190:ASN:H	2.20	0.55
1:C:121:GLN:HB3	1:C:125:ARG:NH1	2.21	0.55
2:G:143:ARG:O	2:G:147:THR:HG23	2.09	0.52
2:F:85:ARG:HA	2:F:88:LEU:HD12	1.91	0.52
1:D:128:MET:HE3	2:G:134:GLN:HE22	1.74	0.51
1:C:144:GLU:OE1	2:F:146:ARG:NH1	2.36	0.51
2:H:166:GLU:O	2:H:170:THR:HG23	2.12	0.50
2:H:92:LEU:C	2:H:92:LEU:HD13	2.32	0.50
1:D:128:MET:HE2	2:G:134:GLN:HE22	1.77	0.49
1:A:165:GLU:HG3	2:H:154:LEU:HD11	1.95	0.48
2:G:95:GLN:NE2	3:G:201:HOH:O	2.46	0.48
1:A:191:ARG:HG2	1:A:191:ARG:O	2.13	0.48
1:D:120:ILE:HG21	2:G:108:LEU:HB3	1.96	0.48
1:A:97:SER:O	1:A:100:GLU:HG2	2.14	0.47
1:B:177:HIS:NE2	2:E:164:LEU:HD21	2.29	0.47
1:B:106:TYR:HB2	2:H:94:SER:OG	2.14	0.47
2:F:114:ASN:HB3	2:F:131:LEU:HD11	1.97	0.46
2:F:174:ARG:HG2	2:F:174:ARG:NH1	2.30	0.46
2:G:107:ALA:HA	2:G:110:ARG:NH1	2.31	0.46
2:F:174:ARG:HG2	2:F:174:ARG:HH11	1.81	0.45
2:G:95:GLN:O	2:G:99:LEU:HD22	2.15	0.45
3:A:215:HOH:O	2:E:125:THR:HG21	2.14	0.45
1:B:187:ILE:N	2:E:174:ARG:HH22	2.15	0.45
2:H:146:ARG:HE	2:H:146:ARG:HB3	1.52	0.44
1:D:121:GLN:O	1:D:125:ARG:HG3	2.17	0.44
2:G:160:LYS:O	2:G:164:LEU:HD23	2.19	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:154:ILE:HD13	2:G:131:LEU:HB3	2.01	0.42
1:B:97:SER:O	1:B:101:ILE:HG23	2.19	0.42
2:G:166:GLU:HA	2:G:169:LYS:HG3	2.02	0.42
1:B:181:GLN:HB3	1:B:184:ARG:NH2	2.35	0.42
1:A:189:GLU:CG	1:A:190:ASN:N	2.84	0.41
1:B:162:LEU:HD23	1:B:162:LEU:HA	1.92	0.41
1:A:184:ARG:HA	1:A:187:ILE:HG22	2.03	0.41
1:C:162:LEU:HD23	1:C:162:LEU:HA	1.92	0.40
1:D:183:LEU:HD23	1:D:183:LEU:HA	1.93	0.40
1:A:190:ASN:O	1:A:191:ARG:C	2.59	0.40
2:H:126:LYS:HA	2:H:126:LYS:HD3	1.99	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:142:ASN:ND2	2:E:85:ARG:NH1[8_457]	1.46	0.74
1:B:142:ASN:ND2	2:E:85:ARG:CZ[8_457]	2.13	0.07

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	94/96 (98%)	93 (99%)	1 (1%)	0	100	100
1	B	89/96 (93%)	89 (100%)	0	0	100	100
1	C	90/96 (94%)	90 (100%)	0	0	100	100
1	D	91/96 (95%)	91 (100%)	0	0	100	100
2	E	94/98 (96%)	94 (100%)	0	0	100	100
2	F	91/98 (93%)	90 (99%)	1 (1%)	0	100	100
2	G	86/98 (88%)	86 (100%)	0	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	H	86/98 (88%)	85 (99%)	1 (1%)	0	100	100
All	All	721/776 (93%)	718 (100%)	3 (0%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	87/88 (99%)	85 (98%)	2 (2%)	50	70
1	B	83/88 (94%)	79 (95%)	4 (5%)	25	41
1	C	84/88 (96%)	81 (96%)	3 (4%)	35	54
1	D	85/88 (97%)	82 (96%)	3 (4%)	36	55
2	E	90/91 (99%)	87 (97%)	3 (3%)	38	57
2	F	87/91 (96%)	84 (97%)	3 (3%)	37	56
2	G	83/91 (91%)	76 (92%)	7 (8%)	11	16
2	H	79/91 (87%)	78 (99%)	1 (1%)	69	84
All	All	678/716 (95%)	652 (96%)	26 (4%)	33	51

All (26) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	120	ILE
1	A	189	GLU
1	B	104	GLN
1	B	118	ILE
1	B	120	ILE
1	B	181	GLN
1	C	156	SER
1	C	186	LYS
1	C	189	GLU
1	D	102	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	115	GLN
1	D	186	LYS
2	E	93	SER
2	E	168	ASN
2	E	171	LEU
2	F	85	ARG
2	F	169	LYS
2	F	175	LEU
2	G	92	LEU
2	G	96	GLN
2	G	99	LEU
2	G	112	GLN
2	G	156	ASP
2	G	169	LYS
2	G	173	LEU
2	H	164	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	159	ASN
1	B	159	ASN
1	B	178	ASN
1	C	116	GLN
1	C	177	HIS
1	D	115	GLN
1	D	159	ASN
1	D	181	GLN
2	F	109	GLN
2	G	134	GLN
2	H	96	GLN
2	H	134	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	96/96 (100%)	0.52	4 (4%) 36 35	39, 65, 148, 187	0
1	B	91/96 (94%)	0.52	6 (6%) 18 17	41, 78, 155, 187	0
1	C	92/96 (95%)	0.32	1 (1%) 80 79	42, 67, 108, 143	0
1	D	93/96 (96%)	0.39	6 (6%) 18 17	43, 84, 161, 183	0
2	E	96/98 (97%)	0.24	2 (2%) 63 61	35, 62, 144, 180	0
2	F	93/98 (94%)	0.33	2 (2%) 62 60	39, 59, 108, 134	0
2	G	88/98 (89%)	0.56	6 (6%) 17 15	61, 96, 204, 222	0
2	H	88/98 (89%)	0.42	6 (6%) 17 15	41, 74, 155, 170	0
All	All	737/776 (94%)	0.41	33 (4%) 33 31	35, 72, 161, 222	0

All (33) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	188	ALA	8.1
2	H	88	LEU	6.4
2	G	90	VAL	5.4
2	H	171	LEU	4.4
1	D	102	ASN	4.3
1	A	187	ILE	3.8
1	A	192	ASN	3.6
2	E	175	LEU	3.5
2	G	173	LEU	3.5
2	G	94	SER	3.3
2	H	90	VAL	3.3
1	B	98	VAL	3.1
1	B	185	ALA	3.1
1	D	105	TYR	3.0
1	D	100	GLU	2.7
2	H	170	THR	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	F	126	LYS	2.7
2	G	88	LEU	2.6
1	C	99	ALA	2.5
2	H	92	LEU	2.5
1	D	103	ALA	2.5
1	B	97	SER	2.4
2	G	91	GLU	2.4
2	H	173	LEU	2.4
1	B	186	LYS	2.4
1	B	141	ARG	2.4
2	E	171	LEU	2.3
2	G	104	ARG	2.3
1	A	191	ARG	2.1
1	D	101	ILE	2.1
1	B	100	GLU	2.1
2	F	173	LEU	2.1
1	D	127	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.