



# Full wwPDB X-ray Structure Validation Report ⓘ

Dec 12, 2023 – 05:45 pm GMT

PDB ID : 4CTE  
Title : Crystal structure of the catalytic domain of the modular laminarinase ZgLamC mutant E142S in complex with a thio-oligosaccharide  
Authors : Labourel, A.; Jam, M.; Legentil, L.; Sylla, B.; Ficko-Blean, E.; Hehemann, J.H.; Ferrieres, V.; Czjzek, M.; Michel, G.  
Deposited on : 2014-03-13  
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5950 (1.80-1.80)
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)
RSRZ outliers	127900	5850 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	233	 5% 90% 9%
1	B	233	 2% 90% 9%
2	C	2	 50% 50%
3	D	3	 67% 33%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GS1	D	1	-	-	-	X
7	GOL	A	284	-	-	X	-
7	GOL	B	284	-	-	X	-

## 2 Entry composition [i](#)

There are 10 unique types of molecules in this entry. The entry contains 4178 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ENDO-1,3-BETA-GLUCANASE, FAMILY GH16.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	231	1878	1190	316	365	7	0	7	0
1	B	231	1881	1189	319	366	7	0	7	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	142	SER	GLU	engineered mutation	UNP G0L2L9
B	142	SER	GLU	engineered mutation	UNP G0L2L9

- Molecule 2 is an oligosaccharide called 1-thio-beta-D-glucopyranose-(1-3)-1-thio-beta-D-glucopyranose.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	O	S			
2	C	2	23	12	9	2	0	0	0

- Molecule 3 is an oligosaccharide called beta-D-glucopyranose-(1-3)-1-thio-beta-D-glucopyranose-(1-3)-1-thio-beta-D-glucopyranose.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	O	S			
3	D	3	34	18	14	2	0	0	0


- Molecule 4 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Ca		
4	A	1	1	1	0	0
4	B	1	1	1	0	0

- Molecule 5 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total Cl 1 1	0	0
5	B	1	Total Cl 1 1	0	0

- Molecule 6 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total C O 4 2 2	0	0
6	B	1	Total C O 4 2 2	0	0

- Molecule 7 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			6	3	3		
7	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 8 is ACETATE ION (three-letter code: ACT) (formula:  $C_2H_3O_2^-$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 9 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	B	1	Total 1	Na 1	0	0


- Molecule 10 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	A	140	Total 140	O 140	0	0
10	B	193	Total 193	O 193	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: ENDO-1,3-BETA-GLUCANASE, FAMILY GH16




- Molecule 1: ENDO-1,3-BETA-GLUCANASE, FAMILY GH16



- Molecule 2: 1-thio-beta-D-glucopyranose-(1-3)-1-thio-beta-D-glucopyranose



- Molecule 3: beta-D-glucopyranose-(1-3)-1-thio-beta-D-glucopyranose-(1-3)-1-thio-beta-D-glucopyranose





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	56.01Å 93.93Å 142.28Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.11 – 1.80 48.11 – 1.80	Depositor EDS
% Data completeness (in resolution range)	99.8 (48.11-1.80) 99.8 (48.11-1.80)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.23 (at 1.81Å)	Xtrriage
Refinement program	REFMAC 5.6.0117	Depositor
R, $R_{free}$	0.177 , 0.222 0.218 , 0.241	Depositor DCC
$R_{free}$ test set	3536 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	21.8	Xtrriage
Anisotropy	0.273	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 47.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	4178	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	33.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.53% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, EDO, CA, NA, GS1, BGC, ACT, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	1.02	6/1950 (0.3%)	0.95	2/2650 (0.1%)
1	B	1.18	7/1950 (0.4%)	1.01	6/2650 (0.2%)
All	All	1.11	13/3900 (0.3%)	0.98	8/5300 (0.2%)

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	230	TRP	CD2-CE2	6.32	1.49	1.41
1	A	192	TRP	CD2-CE2	5.94	1.48	1.41
1	B	29	TRP	CD2-CE2	5.86	1.48	1.41
1	A	156	TRP	CD2-CE2	5.80	1.48	1.41
1	B	52	TRP	NE1-CE2	-5.79	1.30	1.37
1	B	41	TRP	CD2-CE2	5.78	1.48	1.41
1	B	121	TRP	CD2-CE2	5.71	1.48	1.41
1	A	185	TRP	CD2-CE2	5.44	1.47	1.41
1	B	174	ASP	CB-CG	5.42	1.63	1.51
1	B	132	TRP	CD2-CE2	5.40	1.47	1.41
1	A	29	TRP	CD2-CE2	5.39	1.47	1.41
1	A	52	TRP	CD2-CE2	5.16	1.47	1.41
1	B	192	TRP	CD2-CE2	5.12	1.47	1.41

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	186	ASP	CB-CG-OD1	6.99	124.59	118.30
1	B	90	ARG	NE-CZ-NH1	5.86	123.23	120.30
1	A	96	ARG	NE-CZ-NH2	-5.84	117.38	120.30
1	B	247	ASP	CB-CG-OD2	-5.80	113.08	118.30
1	B	62	ARG	NE-CZ-NH1	5.62	123.11	120.30
1	B	247	ASP	CB-CG-OD1	5.44	123.20	118.30

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	158	ASP	CB-CG-OD1	5.27	123.04	118.30
1	B	34	ASP	CB-CG-OD1	5.16	122.94	118.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1878	0	1726	10	0
1	B	1881	0	1722	8	0
2	C	23	0	21	2	0
3	D	34	0	30	1	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	1	0	0	0	0
5	B	1	0	0	1	0
6	A	4	0	6	0	0
6	B	4	0	6	0	0
7	A	6	0	8	5	0
7	B	6	0	8	4	0
8	B	4	0	3	0	0
9	B	1	0	0	0	0
10	A	140	0	0	3	0
10	B	193	0	0	4	0
All	All	4178	0	3530	24	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (24) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:157:SER:O	10:B:2127:HOH:O	2.11	0.68
1:B:134:ALA:O	1:B:213:GLU:HG3	1.96	0.65

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
7:A:284:GOL:H2	2:C:1:GS1:H61	1.82	0.61
1:B:56:GLU:OE2	7:B:284:GOL:H11	2.01	0.61
1:A:100:LYS:HE3	1:A:215:GLN:O	2.03	0.59
7:B:284:GOL:H12	10:B:2041:HOH:O	2.03	0.57
7:B:284:GOL:C1	10:B:2041:HOH:O	2.52	0.57
1:A:239:LYS:NZ	10:A:2133:HOH:O	2.33	0.56
1:B:166:TYR:CD2	1:B:202:VAL:HG12	2.44	0.53
1:A:117:TRP:HZ2	2:C:1:GS1:O5	1.92	0.52
1:B:185:TRP:CD1	1:B:185:TRP:C	2.86	0.49
1:A:56:GLU:OE2	7:A:284:GOL:H11	2.12	0.49
1:B:137:GLU:OE2	1:B:139[A]:ASP:OD1	2.32	0.47
1:A:117:TRP:CE2	7:A:284:GOL:H31	2.50	0.46
1:B:52:TRP:HA	5:B:281:CL:CL	2.54	0.45
1:A:137:GLU:OE2	1:A:139:ASP:OD1	2.35	0.45
7:A:284:GOL:H12	10:A:2032:HOH:O	2.15	0.44
1:A:100:LYS:HB2	1:A:217:GLU:HG2	2.01	0.43
7:A:284:GOL:C1	10:A:2032:HOH:O	2.67	0.42
7:B:284:GOL:H11	10:B:2041:HOH:O	2.18	0.42
1:A:74:ILE:HG23	1:A:89:ALA:HB3	2.02	0.42
1:A:25:TYR:HB3	1:A:252:TYR:HB3	2.04	0.40
1:B:53:GLY:HA3	3:D:2:GS1:O3	2.21	0.40
1:A:117:TRP:CD1	1:A:117:TRP:C	2.94	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	236/233 (101%)	228 (97%)	8 (3%)	0	100	100
1	B	236/233 (101%)	226 (96%)	10 (4%)	0	100	100
All	All	472/466 (101%)	454 (96%)	18 (4%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	197/191 (103%)	196 (100%)	1 (0%)	88	87
1	B	197/191 (103%)	196 (100%)	1 (0%)	88	87
All	All	394/382 (103%)	392 (100%)	2 (0%)	88	87

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	169	ASP
1	B	185	TRP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	114	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

5 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	GS1	C	1	2	10,11,12	1.32	2 (20%)	12,15,17	2.97	7 (58%)
2	GS1	C	2	2	11,12,12	1.65	3 (27%)	15,17,17	1.35	3 (20%)
3	GS1	D	1	3	10,11,12	0.42	0	12,15,17	0.90	1 (8%)
3	GS1	D	2	3	11,12,12	0.57	0	15,17,17	1.60	3 (20%)
3	BGC	D	3	3	11,11,12	0.67	0	15,15,17	1.37	2 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GS1	C	1	2	-	2/2/18/22	0/1/1/1
2	GS1	C	2	2	-	0/2/22/22	0/1/1/1
3	GS1	D	1	3	-	1/2/18/22	0/1/1/1
3	GS1	D	2	3	-	0/2/22/22	0/1/1/1
3	BGC	D	3	3	-	0/2/19/22	0/1/1/1

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	1	GS1	C3-C2	-2.82	1.47	1.52
2	C	2	GS1	O5-C1	2.80	1.46	1.42
2	C	2	GS1	C4-C3	2.75	1.59	1.52
2	C	1	GS1	C3-C4	-2.37	1.48	1.52
2	C	2	GS1	O3-C3	2.18	1.48	1.43

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	1	GS1	C3-C4-C5	-5.61	103.49	110.77
2	C	1	GS1	O5-C5-C4	-5.05	103.09	110.06
2	C	1	GS1	O5-C5-C6	3.70	115.63	106.44
3	D	2	GS1	C1-C2-C3	-3.52	103.63	110.59
3	D	3	BGC	C1-C2-C3	3.46	113.92	109.67

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	2	GS1	O5-C5-C4	-2.88	104.46	109.69
2	C	1	GS1	O4-C4-C3	-2.68	103.28	109.94
2	C	1	GS1	O2-C2-C1	2.66	115.41	110.20
2	C	2	GS1	O5-C1-C2	2.51	113.47	110.31
2	C	2	GS1	O3-C3-C4	2.44	115.98	110.35
3	D	2	GS1	O5-C5-C6	2.41	112.42	106.44
2	C	1	GS1	C3-C2-C1	-2.39	101.39	110.84
2	C	1	GS1	C1-O5-C5	-2.19	108.55	112.57
2	C	2	GS1	C1-C2-C3	-2.10	106.45	110.59
3	D	3	BGC	O5-C1-C2	2.03	113.90	110.77
3	D	1	GS1	C6-C5-C4	-2.02	110.72	113.54

There are no chirality outliers.

All (3) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	C	1	GS1	O5-C5-C6-O6
2	C	1	GS1	C4-C5-C6-O6
3	D	1	GS1	O5-C5-C6-O6

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	1	GS1	2	0
3	D	2	GS1	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



## 5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 5 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul



statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
7	GOL	A	284	-	5,5,5	0.43	0	5,5,5	0.91	0
6	EDO	A	263	-	3,3,3	0.53	0	2,2,2	0.75	0
7	GOL	B	284	-	5,5,5	0.51	0	5,5,5	1.19	0
6	EDO	B	283	-	3,3,3	0.55	0	2,2,2	1.07	0
8	ACT	B	282	4	3,3,3	1.62	1 (33%)	3,3,3	1.96	1 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	GOL	A	284	-	-	4/4/4/4	-
6	EDO	A	263	-	-	0/1/1/1	-
7	GOL	B	284	-	-	4/4/4/4	-
6	EDO	B	283	-	-	1/1/1/1	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
8	B	282	ACT	O-C	2.45	1.33	1.22

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
8	B	282	ACT	OXT-C-O	2.86	132.58	122.05

There are no chirality outliers.

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
7	A	284	GOL	O1-C1-C2-C3
7	A	284	GOL	C1-C2-C3-O3
7	B	284	GOL	O1-C1-C2-C3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms
7	B	284	GOL	C1-C2-C3-O3
7	A	284	GOL	O1-C1-C2-O2
7	B	284	GOL	O1-C1-C2-O2
7	A	284	GOL	O2-C2-C3-O3
7	B	284	GOL	O2-C2-C3-O3
6	B	283	EDO	O1-C1-C2-O2

There are no ring outliers.

2 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	A	284	GOL	5	0
7	B	284	GOL	4	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	231/233 (99%)	0.24	12 (5%) 27 22	22, 37, 65, 91	2 (0%)
1	B	231/233 (99%)	-0.46	4 (1%) 70 66	16, 24, 36, 63	1 (0%)
All	All	462/466 (99%)	-0.11	16 (3%) 44 38	16, 29, 60, 91	3 (0%)

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	49	TYR	4.6
1	A	35[A]	ASP	4.4
1	A	159	GLN	3.5
1	A	24	ASP	3.4
1	A	205	SER	2.8
1	A	160	ASN	2.7
1	A	208	VAL	2.6
1	A	117	TRP	2.6
1	B	24	ASP	2.5
1	A	34	ASP	2.5
1	B	197[A]	GLN	2.2
1	B	166	TYR	2.1
1	A	206	ASN	2.1
1	A	230	TRP	2.1
1	A	193	PHE	2.1
1	B	230	TRP	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.


### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	GS1	D	1	11/12	0.34	0.46	29,32,38,53	11
2	GS1	C	1	11/12	0.68	0.38	29,31,39,57	11
3	BGC	D	3	11/12	0.76	0.19	34,39,41,47	11
2	GS1	C	2	12/12	0.78	0.17	38,45,52,57	1
3	GS1	D	2	12/12	0.89	0.16	26,28,36,38	12

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
6	EDO	A	263	4/4	0.90	0.14	35,36,39,43	0
7	GOL	A	284	6/6	0.90	0.14	25,30,35,42	6
7	GOL	B	284	6/6	0.90	0.18	21,27,33,41	6
6	EDO	B	283	4/4	0.91	0.13	30,31,32,37	0
8	ACT	B	282	4/4	0.91	0.16	25,25,31,41	0
5	CL	B	281	1/1	0.92	0.07	43,43,43,43	0
5	CL	A	261	1/1	0.93	0.10	43,43,43,43	0
4	CA	A	260	1/1	0.97	0.05	39,39,39,39	0
9	NA	B	285	1/1	0.98	0.06	31,31,31,31	0
4	CA	B	280	1/1	1.00	0.07	21,21,21,21	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.