



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 18, 2024 – 01:33 AM EST

PDB ID : 4CTS  
Title : CRYSTAL STRUCTURE ANALYSIS AND MOLECULAR MODEL OF A COMPLEX OF CITRATE SYNTHASE WITH OXALOACETATE AND S-ACETONYL-COENZYME A  
Authors : Remington, S.; Wiegand, G.; Huber, R.  
Deposited on : 1984-01-27  
Resolution : 2.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtrriage (Phenix) : **NOT EXECUTED**  
EDS : **NOT EXECUTED**  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	2172 (2.90-2.90)
Ramachandran outliers	138981	2115 (2.90-2.90)
Sidechain outliers	138945	2117 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	437	
1	B	437	

## 2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 7002 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CITRATE SYNTHASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	437	Total 3444	C 2200	N 591	O 634	S 19	227	0	0
1	B	437	Total 3444	C 2200	N 591	O 634	S 19	230	0	0

- Molecule 2 is OXALOACETATE ION (three-letter code: OAA) (formula:  $C_4H_3O_5$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
2	A	1	Total 9	C 4	O 5	0	0
2	B	1	Total 9	C 4	O 5	0	0

- Molecule 3 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	49	Total O 49 49	0	0
3	B	47	Total O 47 47	0	0

### 3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


Note EDS was not executed.

- Molecule 1: CITRATE SYNTHASE



- Molecule 1: CITRATE SYNTHASE





## 4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	101.54Å 101.54Å 224.60Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	5.00 – 2.90	Depositor
% Data completeness (in resolution range)	(Not available) (5.00-2.90)	Depositor
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
Refinement program	EREF	Depositor
R, $R_{free}$	0.182 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	7002	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	20.0	wwPDB-VP

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: OAA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	1.22	11/3528 (0.3%)	1.77	41/4786 (0.9%)
1	B	1.22	11/3528 (0.3%)	1.77	54/4786 (1.1%)
All	All	1.22	22/7056 (0.3%)	1.77	95/9572 (1.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	1	105
1	B	2	104
All	All	3	209

The worst 5 of 22 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	114	TRP	NE1-CE2	-8.77	1.26	1.37
1	B	109	TRP	NE1-CE2	-8.18	1.26	1.37
1	A	114	TRP	NE1-CE2	-7.93	1.27	1.37
1	A	168	TRP	NE1-CE2	-7.85	1.27	1.37
1	A	371	TRP	NE1-CE2	-7.65	1.27	1.37

The worst 5 of 95 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	117	ARG	NE-CZ-NH2	-17.05	111.78	120.30
1	B	65	ARG	NE-CZ-NH1	14.64	127.62	120.30
1	A	191	ARG	NE-CZ-NH2	-11.17	114.72	120.30
1	A	67	ARG	NE-CZ-NH1	-10.86	114.87	120.30

*Continued on next page...*



Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	20	ARG	NE-CZ-NH2	-9.80	115.40	120.30

All (3) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
1	A	37	THR	CB
1	B	37	THR	CB
1	B	307	ASN	CA

5 of 209 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	11	ALA	Mainchain
1	A	18	GLN	Sidechain
1	A	25	ARG	Sidechain
1	A	26	GLN	Sidechain
1	A	32	VAL	Mainchain

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3444	0	3436	291	17
1	B	3444	0	3436	283	27
2	A	9	0	2	0	0
2	B	9	0	2	0	0
3	A	49	0	0	4	0
3	B	47	0	0	0	2
All	All	7002	0	6876	502	38

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 39.

The worst 5 of 502 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:268:GLY:HA2	1:B:251:VAL:HG13	1.47	0.96
1:A:419:LEU:HD11	1:B:243:VAL:HG22	1.51	0.92
1:B:58:LEU:HD12	1:B:63:GLY:HA2	1.53	0.90
1:A:280:GLU:HA	1:A:283:VAL:HB	1.54	0.90
1:A:75:GLN:HA	1:A:87:PRO:HG3	1.56	0.88

The worst 5 of 38 symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:203:ILE:O	3:B:476:HOH:O[6_455]	0.90	1.30
1:B:153:ASN:OD1	1:B:306:TRP:CH2[3_544]	0.98	1.22
1:A:206:LYS:CD	1:B:191:ARG:NH2[6_455]	1.22	0.98
1:B:156:ARG:CZ	1:B:303:ASP:CB[3_544]	1.41	0.79
1:B:153:ASN:CG	1:B:306:TRP:CH2[3_544]	1.42	0.78

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	435/437 (100%)	359 (82%)	55 (13%)	21 (5%)	<b>2</b> <b>8</b>
1	B	435/437 (100%)	357 (82%)	59 (14%)	19 (4%)	<b>2</b> <b>10</b>
All	All	870/874 (100%)	716 (82%)	114 (13%)	40 (5%)	<b>2</b> <b>9</b>

5 of 40 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	166	LYS
1	A	205	SER
1	A	239	GLU
1	B	166	LYS
1	B	239	GLU

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	371/371 (100%)	311 (84%)	60 (16%)	2	7
1	B	371/371 (100%)	315 (85%)	56 (15%)	3	9
All	All	742/742 (100%)	626 (84%)	116 (16%)	2	8

5 of 116 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	413	ARG
1	B	413	ARG
1	B	112	LYS
1	B	405	VAL
1	B	343	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 33 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	246	HIS
1	B	267	ASN
1	B	391	ASN
1	A	333	GLN
1	A	279	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	OAA	B	438	-	8,8,8	4.21	1 (12%)	9,10,10	1.50	2 (22%)
2	OAA	A	438	-	8,8,8	4.58	1 (12%)	9,10,10	1.27	1 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	OAA	B	438	-	-	1/8/8/8	-
2	OAA	A	438	-	-	2/8/8/8	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	438	OAA	C3-C4	-12.58	1.36	1.53
2	B	438	OAA	C3-C4	-11.33	1.38	1.53

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	438	OAA	O5-C4-C3	2.49	120.80	113.97
2	A	438	OAA	O2-C1-O1	-2.41	117.28	123.30
2	B	438	OAA	O5-C4-O4	-2.22	118.53	123.61

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	438	OAA	C1-C2-C3-O3
2	A	438	OAA	C1-C2-C3-O3
2	A	438	OAA	C1-C2-C3-C4

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

EDS was not executed - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS was not executed - this section is therefore empty.

### 6.3 Carbohydrates [i](#)

EDS was not executed - this section is therefore empty.

### 6.4 Ligands [i](#)

EDS was not executed - this section is therefore empty.

### 6.5 Other polymers [i](#)

EDS was not executed - this section is therefore empty.