

Full wwPDB X-ray Structure Validation Report (i)

Dec 16, 2023 – 08:50 pm GMT

PDB ID	:	4CVX
Title	:	COMPLEX OF A B2 CHICKEN MHC CLASS I MOLECULE AND A 9MER
		CHICKEN PEPTIDE
Authors	:	Chappell, P.E.; Roversi, P.; Harrison, M.C.; Mears, L.E.; Kaufman, J.F.; Lea,
		S.M.
Deposited on		
Resolution	:	3.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:


Refmac	: : :	1.132.3620191225.v01 (using entries in the PDB archive December 25th 2019)5.8.0158
CCP4 Ideal geometry (proteins) Ideal geometry (DNA, RNA) Validation Pipeline (wwPDB-VP)	:	Parkinson et al. (1996)

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY \, DIFFRACTION$

The reported resolution of this entry is 3.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
R_{free}	130704	1149 (3.34-3.26)
Clashscore	141614	1205 (3.34-3.26)
Ramachandran outliers	138981	1183 (3.34-3.26)
Sidechain outliers	138945	1182 (3.34-3.26)
RSRZ outliers	127900	1115 (3.34-3.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	А	310	85%	12%
1	D	310	85% •	12%
2	В	98	98%	·
2	Е	98	97%	•••
3	С	9	89%	11%



Continued from previous page...

Mol	Chain	Length	Quality of chain
3	F	9	100%



4CVX

2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 6060 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace	
1	٨	272	Total	С	Ν	0	S	0	0	0
			2193	1385	388	412	8			
1	Л	272	Total	С	Ν	0	S	0	0	0
			2193	1385	388	412	8	0	0	0

• Molecule 1 is a protein called MHC CLASS I ALPHA CHAIN 2.

Chain	Residue	Modelled	Actual	Comment	Reference
А	273	ARG	-	expression tag	UNP O46789
А	274	SER	-	expression tag	UNP O46789
А	275	GLY	-	expression tag	UNP O46789
А	276	GLY	-	expression tag	UNP O46789
А	277	GLY	-	expression tag	UNP O46789
А	278	LEU	-	expression tag	UNP O46789
А	279	ASN	-	expression tag	UNP O46789
А	280	ASP	-	expression tag	UNP O46789
А	281	ILE	-	expression tag	UNP O46789
А	282	PHE	-	expression tag	UNP O46789
A	283	GLU	-	expression tag	UNP O46789
А	284	ALA	-	expression tag	UNP O46789
А	285	GLN	-	expression tag	UNP O46789
A	286	LYS	-	expression tag	UNP O46789
А	287	ILE	-	expression tag	UNP O46789
А	288	GLU	-	expression tag	UNP O46789
А	289	TRP	-	expression tag	UNP O46789
А	290	HIS	-	expression tag	UNP O46789
А	291	GLU	-	expression tag	UNP O46789
А	292	ASN	-	expression tag	UNP O46789
А	293	SER	-	expression tag	UNP O46789
А	294	SER	-	expression tag	UNP O46789
А	295	SER	-	expression tag	UNP O46789
А	296	VAL	-	expression tag	UNP O46789
А	297	ASP	-	expression tag	UNP O46789

There are 76 discrepancies between the modelled and reference sequences:



Chain	Residue	Modelled	Actual	Comment	Reference
А	298	LYS	-	expression tag	UNP O46789
А	299	LEU	-	expression tag	UNP O46789
А	300	ALA	-	expression tag	UNP O46789
А	301	ALA	-	expression tag	UNP O46789
А	302	ALA	-	expression tag	UNP O46789
А	303	LEU	-	expression tag	UNP O46789
А	304	GLU	-	expression tag	UNP O46789
А	305	HIS	-	expression tag	UNP O46789
А	306	HIS	-	expression tag	UNP O46789
А	307	HIS	-	expression tag	UNP O46789
А	308	HIS	-	expression tag	UNP O46789
А	309	HIS	-	expression tag	UNP O46789
А	310	HIS	-	expression tag	UNP O46789
D	273	ARG	-	expression tag	UNP O46789
D	274	SER	-	expression tag	UNP O46789
D	275	GLY	-	expression tag	UNP O46789
D	276	GLY	-	expression tag	UNP O46789
D	277	GLY	-	expression tag	UNP O46789
D	278	LEU	-	expression tag	UNP O46789
D	279	ASN	-	expression tag	UNP O46789
D	280	ASP	-	expression tag	UNP O46789
D	281	ILE	-	expression tag	UNP O46789
D	282	PHE	-	expression tag	UNP O46789
D	283	GLU	-	expression tag	UNP O46789
D	284	ALA	-	expression tag	UNP O46789
D	285	GLN	-	expression tag	UNP O46789
D	286	LYS	-	expression tag	UNP O46789
D	287	ILE	-	expression tag	UNP O46789
D	288	GLU	-	expression tag	UNP O46789
D	289	TRP	-	expression tag	UNP O46789
D	290	HIS	-	expression tag	UNP O46789
D	291	GLU	-	expression tag	UNP O46789
D	292	ASN	-	expression tag	UNP O46789
D	293	SER	-	expression tag	UNP O46789
D	294	SER	-	expression tag	UNP O46789
D	295	SER	-	expression tag	UNP O46789
D	296	VAL	-	expression tag	UNP O46789
D	297	ASP	-	expression tag	UNP O46789
D	298	LYS	-	expression tag	UNP O46789
D	299	LEU	-	expression tag	UNP O46789
D	300	ALA	-	expression tag	UNP O46789
D	301	ALA	-	expression tag	UNP O46789

 α . . : 1 0



Contentia	Continucu from previous page								
Chain	Residue	Modelled	Actual	Comment	Reference				
D	302	ALA	-	expression tag	UNP O46789				
D	303	LEU	-	expression tag	UNP O46789				
D	304	GLU	-	expression tag	UNP O46789				
D	305	HIS	-	expression tag	UNP O46789				
D	306	HIS	-	expression tag	UNP O46789				
D	307	HIS	-	expression tag	UNP O46789				
D	308	HIS	-	expression tag	UNP O46789				
D	309	HIS	-	expression tag	UNP O46789				
D	310	HIS	-	expression tag	UNP O46789				

Continued from previous page...

• Molecule 2 is a protein called BETA-2-MICROGLOBULIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	В	96	Total	С	Ν	0	S	0	0	0
			759	486	124	144	5			
0	F	06	Total	С	Ν	0	S	0	0	0
	96	759	486	124	144	5	0	0	0	

• Molecule 3 is a protein called SELF-PEPTIDE.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
3	3 C 9	Total C N O	0	0	0	
5		9	74 50 10 14	0	0	U
2	о Б	0	Total C N O	0	0	0
ЭГ	9	74 50 10 14	0	0	0	

• Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	А	4	Total O 4 4	0	0
4	В	1	Total O 1 1	0	0
4	D	2	Total O 2 2	0	0
4	F	1	Total O 1 1	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: MHC CLASS I ALPHA CHAIN 2

Chain A:	85%	12%	I
E1 F33 V34 N76 L80 M128 M128	A160 A160 A160 A160 A160 A17 A17 A16 A17 A17 A17 A17 A17 A17 A17 A14 A14 A14 A14 A14 A14 A14 A14 A14 A14	GLU HIS HIS HIS HIS	HIS
• Molecule 1:	MHC CLASS I ALPHA CHAIN 2		
Chain D:	85%	12%	I
E1 F33 V34 N76 L80 M128 M128	K143 A150 A150 A150 A150 C12 A12 A12 A12 A12 A12 A12 A12 A12 A12 A	ALA ALA LEU GLU HIS HIS	SIH SIH SIH
• Molecule 2:	BETA-2-MICROGLOBULIN		
Chain B:	98%	·	
ASP L2 E97 PHE			
• Molecule 2:	BETA-2-MICROGLOBULIN		
Chain E:	97%		
ASP L2 P13 E97 PHE			
• Molecule 3:	SELF-PEPTIDE		
Chain C:	89%	11%	
ri L9			
• Molecule 3:	SELF-PEPTIDE		
Chain F:	100%		



There are no outlier residues recorded for this chain.



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 65	Depositor
Cell constants	173.85Å 173.85Å 87.51Å	Deperitor
a, b, c, α , β , γ	90.00° 90.00° 120.00°	Depositor
Resolution (Å)	75.66 - 3.30	Depositor
Resolution (A)	$75.66 \ - \ 3.30$	EDS
% Data completeness	99.9 (75.66-3.30)	Depositor
(in resolution range)	$99.9\ (75.66\text{-}3.30)$	EDS
R _{merge}	0.16	Depositor
R _{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	3.02 (at 3.33Å)	Xtriage
Refinement program	REFMAC 5.7.0032	Depositor
D D	0.237 , 0.262	Depositor
R, R_{free}	0.224 , 0.249	DCC
R_{free} test set	1171 reflections (5.12%)	wwPDB-VP
Wilson B-factor $(Å^2)$	67.8	Xtriage
Anisotropy	0.553	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.30 , 13.5	EDS
L-test for twinning ²	$< L > = 0.38, < L^2 > = 0.21$	Xtriage
Estimated twinning fraction	0.357 for h,-h-k,-l	Xtriage
Reported twinning fraction	0.615 for H, K, L	Depositor
Reported twinning fraction	0.385 for K, H, -L	Depositor
Outliers	0 of 22858 reflections	Xtriage
F_o, F_c correlation	0.87	EDS
Total number of atoms	6060	wwPDB-VP
Average B, all atoms $(Å^2)$	77.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 2.76% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles	
	Ullalli	RMSZ # Z > 5		RMSZ	# Z > 5
1	А	0.27	0/2258	0.43	0/3071
1	D	0.27	0/2258	0.44	0/3071
2	В	0.27	0/783	0.42	0/1065
2	Е	0.27	0/783	0.42	0/1065
3	С	0.34	0/77	0.50	0/105
3	F	0.34	0/77	0.43	0/105
All	All	0.27	0/6236	0.43	0/8482

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	2193	0	2062	5	0
1	D	2193	0	2062	6	0
2	В	759	0	721	0	0
2	Е	759	0	721	1	0
3	С	74	0	72	1	0
3	F	74	0	72	0	0
4	А	4	0	0	0	0
4	В	1	0	0	0	0
4	D	2	0	0	0	0
4	F	1	0	0	0	0



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	6060	0	5710	10	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (10) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:143:LYS:HE2	1:D:143:LYS:HG2	1.85	0.57
1:D:128:MET:SD	1:D:150:ALA:HB1	2.53	0.48
1:A:128:MET:SD	1:A:150:ALA:HB1	2.54	0.47
1:A:143:LYS:CE	1:D:143:LYS:HG2	2.46	0.44
1:D:185:ARG:HD2	2:E:13:PRO:HB3	1.99	0.43
1:A:76:ASN:O	1:A:80:LEU:HD12	2.19	0.42
3:C:9:LEU:C	3:C:9:LEU:HD12	2.40	0.42
1:D:76:ASN:O	1:D:80:LEU:HD12	2.19	0.41
1:D:33:PHE:CD2	1:D:34:VAL:HG23	2.55	0.41
1:A:33:PHE:CD2	1:A:34:VAL:HG23	2.55	0.41

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	А	270/310~(87%)	260~(96%)	10 (4%)	0	100 100
1	D	270/310~(87%)	260 (96%)	10 (4%)	0	100 100
2	В	94/98~(96%)	89~(95%)	5(5%)	0	100 100
2	Ε	94/98~(96%)	89~(95%)	5 (5%)	0	100 100
3	С	7/9~(78%)	6 (86%)	1 (14%)	0	100 100



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
3	F	7/9~(78%)	6 (86%)	1 (14%)	0	100	100
All	All	742/834~(89%)	710 (96%)	32~(4%)	0	100	100

Continued from previous page...

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent side chain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	А	222/253~(88%)	221 (100%)	1 (0%)	88 93	
1	D	222/253~(88%)	220~(99%)	2(1%)	78 87	
2	В	84/86~(98%)	84 (100%)	0	100 100	
2	Ε	84/86~(98%)	84 (100%)	0	100 100	
3	С	8/8 (100%)	8 (100%)	0	100 100	
3	F	8/8 (100%)	8 (100%)	0	100 100	
All	All	628/694~(90%)	625~(100%)	3(0%)	88 93	

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	А	175	GLU
1	D	223	ASP
1	D	271	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.



5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

There are no ligands in this entry.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	$\langle RSRZ \rangle$	ŧ	₽RSR	Z>2	$OWAB(A^2)$	Q<0.9
1	А	272/310~(87%)	0.19	0	100	100	67, 76, 85, 89	0
1	D	272/310 (87%)	0.21	0	100	100	66, 76, 98, 100	0
2	В	96/98~(97%)	0.11	0	100	100	74, 78, 84, 84	0
2	Е	96/98~(97%)	0.11	0	100	100	74, 78, 83, 87	0
3	С	9/9~(100%)	0.66	0	100	100	73, 74, 75, 76	0
3	F	9/9~(100%)	0.07	0	100	100	70, 70, 71, 71	0
All	All	754/834~(90%)	0.18	0	100	100	66, 77, 93, 100	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

There are no ligands in this entry.

6.5 Other polymers (i)

There are no such residues in this entry.

