



# wwPDB X-ray Structure Validation Summary Report ⓘ

Sep 26, 2023 – 12:46 PM EDT

PDB ID : 6CWD  
Title : Hepatitis B core-antigen in complex with scFv e13  
Authors : Eren, E.; Steven, A.C.; Wingfield, P.T.  
Deposited on : 2018-03-30  
Resolution : 3.33 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.35.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.33 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1060 (3.38-3.30)
Clashscore	141614	1111 (3.38-3.30)
Ramachandran outliers	138981	1090 (3.38-3.30)
Sidechain outliers	138945	1089 (3.38-3.30)
RSRZ outliers	127900	1028 (3.38-3.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	261	
1	B	261	
1	E	261	
1	G	261	
2	C	149	

Continued on next page...

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
2	D	149	 <p>5% 56% 31% • 11%</p>
2	F	149	 <p>% 56% 28% • 15%</p>
2	H	149	 <p>% 57% 29% • 12%</p>

## 2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 10852 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Single chain variable fragment (scFv) e13.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	B	231	1687	1060	282	340	5	0	0	0
1	G	231	1687	1060	282	340	5	0	0	0
1	A	231	1687	1060	282	340	5	0	0	0
1	E	229	1674	1051	280	338	5	0	0	0

- Molecule 2 is a protein called Capsid protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	D	133	1050	687	174	187	2	0	0	0
2	H	131	1031	673	172	184	2	0	0	0
2	C	129	1026	671	170	183	2	0	0	0
2	F	127	1010	661	168	179	2	0	0	0

There are 8 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
D	48	ALA	CYS	engineered mutation	UNP P03147
D	107	ALA	CYS	engineered mutation	UNP P03147
H	48	ALA	CYS	engineered mutation	UNP P03147
H	107	ALA	CYS	engineered mutation	UNP P03147
C	48	ALA	CYS	engineered mutation	UNP P03147
C	107	ALA	CYS	engineered mutation	UNP P03147
F	48	ALA	CYS	engineered mutation	UNP P03147
F	107	ALA	CYS	engineered mutation	UNP P03147

### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Single chain variable fragment (scFv) e13

Chain B:  62% 25% 11%




- Molecule 1: Single chain variable fragment (scFv) e13

Chain G:  2% 66% 23% 11%




- Molecule 1: Single chain variable fragment (scFv) e13


Chain A:  75% 13% 11%




• Molecule 1: Single chain variable fragment (scFv) e13




• Molecule 2: Capsid protein



• Molecule 2: Capsid protein




• Molecule 2: Capsid protein



• Molecule 2: Capsid protein





## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	219.54Å 65.10Å 148.21Å 90.00° 101.65° 90.00°	Depositor
Resolution (Å)	19.92 – 3.33 19.92 – 3.33	Depositor EDS
% Data completeness (in resolution range)	97.7 (19.92-3.33) 84.7 (19.92-3.33)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.42 (at 3.36Å)	Xtrriage
Refinement program	PHENIX (1.10.1_2155: ???)	Depositor
R, $R_{free}$	0.242 , 0.282 0.242 , 0.280	Depositor DCC
$R_{free}$ test set	1959 reflections (6.64%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	47.4	Xtrriage
Anisotropy	0.721	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.27 , 31.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.45$ , $\langle L^2 \rangle = 0.28$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.85	EDS
Total number of atoms	10852	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	57.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.02% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.28	0/1719	0.58	1/2338 (0.0%)
1	B	0.29	0/1719	0.56	0/2338
1	E	0.27	0/1706	0.55	0/2320
1	G	0.28	0/1719	0.55	0/2338
2	C	0.33	0/1058	0.57	0/1450
2	D	0.48	1/1083 (0.1%)	0.63	0/1485
2	F	0.32	0/1041	0.57	0/1426
2	H	0.33	0/1063	0.60	0/1457
All	All	0.32	1/11108 (0.0%)	0.57	1/15152 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	C	0	1
2	H	0	1
All	All	0	2

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	44	SER	C-N	11.03	1.55	1.34

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	146	LEU	CA-CB-CG	6.41	130.05	115.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	C	46	GLU	Peptide
2	H	44	SER	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1687	0	1668	26	0
1	B	1687	0	1668	41	0
1	E	1674	0	1655	34	0
1	G	1687	0	1668	36	0
2	C	1026	0	1011	37	1
2	D	1050	0	1031	37	0
2	F	1010	0	998	36	0
2	H	1031	0	1013	34	1
All	All	10852	0	10712	263	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

The worst 5 of 263 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:35:LEU:HD11	1:A:90:CYS:HB2	1.56	0.88
2:C:15:LEU:HB3	2:C:119:LEU:HD13	1.58	0.86
2:D:86:VAL:HG13	2:D:87:SER:H	1.41	0.84
2:C:57:GLN:HG2	2:F:57:GLN:HG2	1.64	0.78
1:E:79:GLY:O	1:E:81:GLN:NE2	2.18	0.77

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:18:PHE:O	2:C:132:TYR:OH[2_7510]	2.11	0.09

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	227/261 (87%)	215 (95%)	12 (5%)	0	100	100
1	B	227/261 (87%)	213 (94%)	13 (6%)	1 (0%)	34	68
1	E	225/261 (86%)	210 (93%)	14 (6%)	1 (0%)	34	68
1	G	227/261 (87%)	215 (95%)	12 (5%)	0	100	100
2	C	125/149 (84%)	109 (87%)	16 (13%)	0	100	100
2	D	129/149 (87%)	109 (84%)	16 (12%)	4 (3%)	4	26
2	F	123/149 (83%)	106 (86%)	16 (13%)	1 (1%)	19	53
2	H	127/149 (85%)	111 (87%)	15 (12%)	1 (1%)	19	53
All	All	1410/1640 (86%)	1288 (91%)	114 (8%)	8 (1%)	25	60

5 of 8 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	D	46	GLU
2	H	9	PHE
2	F	43	GLU
2	D	86	VAL
2	D	47	HIS

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	183/202 (91%)	183 (100%)	0	100	100

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	183/202 (91%)	180 (98%)	3 (2%)	62	81
1	E	182/202 (90%)	182 (100%)	0	100	100
1	G	183/202 (91%)	182 (100%)	1 (0%)	88	93
2	C	111/130 (85%)	109 (98%)	2 (2%)	59	79
2	D	113/130 (87%)	111 (98%)	2 (2%)	59	79
2	F	109/130 (84%)	106 (97%)	3 (3%)	43	71
2	H	111/130 (85%)	109 (98%)	2 (2%)	59	79
All	All	1175/1328 (88%)	1162 (99%)	13 (1%)	73	86

5 of 13 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	H	92	ASN
2	C	46	GLU
2	F	112	ARG
2	F	61	CYS
2	F	95	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	G	81	GLN
2	H	47	HIS
1	E	81	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	231/261 (88%)	-0.17	0 <span style="border: 1px solid blue; padding: 2px;">100</span> <span style="border: 1px solid blue; padding: 2px;">100</span>	28, 56, 81, 102	0
1	B	231/261 (88%)	-0.30	1 (0%) <span style="border: 1px solid blue; padding: 2px;">92</span> <span style="border: 1px solid blue; padding: 2px;">94</span>	25, 47, 69, 80	0
1	E	229/261 (87%)	-0.14	2 (0%) <span style="border: 1px solid blue; padding: 2px;">84</span> <span style="border: 1px solid blue; padding: 2px;">85</span>	45, 64, 89, 112	0
1	G	231/261 (88%)	-0.21	5 (2%) <span style="border: 1px solid blue; padding: 2px;">62</span> <span style="border: 1px solid blue; padding: 2px;">61</span>	27, 50, 78, 97	0
2	C	129/149 (86%)	-0.09	4 (3%) <span style="border: 1px solid gray; padding: 2px;">49</span> <span style="border: 1px solid gray; padding: 2px;">48</span>	31, 58, 97, 112	0
2	D	133/149 (89%)	-0.08	7 (5%) <span style="border: 1px solid red; padding: 2px;">26</span> <span style="border: 1px solid red; padding: 2px;">27</span>	30, 55, 99, 112	0
2	F	127/149 (85%)	-0.14	1 (0%) <span style="border: 1px solid blue; padding: 2px;">86</span> <span style="border: 1px solid blue; padding: 2px;">87</span>	36, 60, 99, 129	0
2	H	131/149 (87%)	-0.16	2 (1%) <span style="border: 1px solid blue; padding: 2px;">73</span> <span style="border: 1px solid blue; padding: 2px;">72</span>	31, 56, 107, 119	0
All	All	1442/1640 (87%)	-0.17	22 (1%) <span style="border: 1px solid blue; padding: 2px;">73</span> <span style="border: 1px solid blue; padding: 2px;">72</span>	25, 56, 87, 129	0

The worst 5 of 22 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	H	144	PRO	4.2
2	D	145	GLU	3.4
2	C	42	LEU	3.2
1	G	249	SER	3.1
1	G	130	GLU	2.8

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands

There are no ligands in this entry.

## 6.5 Other polymers

There are no such residues in this entry.