



wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 14, 2023 – 10:46 PM EDT

PDB ID : 8CX3
Title : Crystal structure of full-length mesothelin
Authors : Zhan, J.; Esser, L.; Xia, D.
Deposited on : 2022-05-19
Resolution : 3.61 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.61 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1257 (3.70-3.50)
Clashscore	141614	1353 (3.70-3.50)
Ramachandran outliers	138981	1307 (3.70-3.50)
Sidechain outliers	138945	1307 (3.70-3.50)
RSRZ outliers	127900	1161 (3.70-3.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	312	 5% 65% 13% 21%
1	B	312	 2% 66% 13% 21%
1	C	312	 3% 67% 12% 21%
1	D	312	 5% 66% 13% 21%
1	E	312	 5% 65% 14% 21%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	312	 8% 64% 14% 21%
2	G	2	 100%
3	H	3	 100%

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 11960 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Isoform 4 of Mesothelin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	245	1942	1245	318	369	10	0	0	0
1	B	248	1963	1258	323	372	10	0	0	0
1	C	248	1963	1258	323	372	10	0	0	0
1	D	248	1963	1258	323	372	10	0	0	0
1	E	248	1963	1258	323	372	10	0	0	0
1	F	248	1963	1258	323	372	10	0	0	0

There are 48 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	295	MET	-	initiating methionine	UNP Q13421
A	593	VAL	MET	conflict	UNP Q13421
A	601	HIS	-	expression tag	UNP Q13421
A	602	HIS	-	expression tag	UNP Q13421
A	603	HIS	-	expression tag	UNP Q13421
A	604	HIS	-	expression tag	UNP Q13421
A	605	HIS	-	expression tag	UNP Q13421
A	606	HIS	-	expression tag	UNP Q13421
B	295	MET	-	initiating methionine	UNP Q13421
B	593	VAL	MET	conflict	UNP Q13421
B	601	HIS	-	expression tag	UNP Q13421
B	602	HIS	-	expression tag	UNP Q13421
B	603	HIS	-	expression tag	UNP Q13421
B	604	HIS	-	expression tag	UNP Q13421
B	605	HIS	-	expression tag	UNP Q13421
B	606	HIS	-	expression tag	UNP Q13421
C	295	MET	-	initiating methionine	UNP Q13421

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
C	593	VAL	MET	conflict	UNP Q13421
C	601	HIS	-	expression tag	UNP Q13421
C	602	HIS	-	expression tag	UNP Q13421
C	603	HIS	-	expression tag	UNP Q13421
C	604	HIS	-	expression tag	UNP Q13421
C	605	HIS	-	expression tag	UNP Q13421
C	606	HIS	-	expression tag	UNP Q13421
D	295	MET	-	initiating methionine	UNP Q13421
D	593	VAL	MET	conflict	UNP Q13421
D	601	HIS	-	expression tag	UNP Q13421
D	602	HIS	-	expression tag	UNP Q13421
D	603	HIS	-	expression tag	UNP Q13421
D	604	HIS	-	expression tag	UNP Q13421
D	605	HIS	-	expression tag	UNP Q13421
D	606	HIS	-	expression tag	UNP Q13421
E	295	MET	-	initiating methionine	UNP Q13421
E	593	VAL	MET	conflict	UNP Q13421
E	601	HIS	-	expression tag	UNP Q13421
E	602	HIS	-	expression tag	UNP Q13421
E	603	HIS	-	expression tag	UNP Q13421
E	604	HIS	-	expression tag	UNP Q13421
E	605	HIS	-	expression tag	UNP Q13421
E	606	HIS	-	expression tag	UNP Q13421
F	295	MET	-	initiating methionine	UNP Q13421
F	593	VAL	MET	conflict	UNP Q13421
F	601	HIS	-	expression tag	UNP Q13421
F	602	HIS	-	expression tag	UNP Q13421
F	603	HIS	-	expression tag	UNP Q13421
F	604	HIS	-	expression tag	UNP Q13421
F	605	HIS	-	expression tag	UNP Q13421
F	606	HIS	-	expression tag	UNP Q13421

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace	
			Total	C	N				O
2	G	2	28	16	2	10	0	0	0

- Molecule 3 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	H	3	39	22	2	15	0	0	0

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
4	A	1	14	8	1	5	0	0
4	B	1	14	8	1	5	0	0
4	C	1	14	8	1	5	0	0
4	E	1	14	8	1	5	0	0

- Molecule 5 is PHOSPHATE ION (three-letter code: PO4) (formula: O_4P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	O	P	0	0
			5	4	1		
5	B	1	Total	O	P	0	0
			5	4	1		
5	B	1	Total	O	P	0	0
			5	4	1		
5	C	1	Total	O	P	0	0
			5	4	1		
5	D	1	Total	O	P	0	0
			5	4	1		
5	E	1	Total	O	P	0	0
			5	4	1		
5	E	1	Total	O	P	0	0
			5	4	1		
5	E	1	Total	O	P	0	0
			5	4	1		
5	F	1	Total	O	P	0	0
			5	4	1		
5	F	1	Total	O	P	0	0
			5	4	1		

- Molecule 6 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			6	3	3		
6	B	1	Total	C	O	0	0
			6	3	3		
6	C	1	Total	C	O	0	0
			6	3	3		
6	D	1	Total	C	O	0	0
			6	3	3		
6	F	1	Total	C	O	0	0
			6	3	3		


3 Residue-property plots i

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Isoform 4 of Mesothelin




- Molecule 1: Isoform 4 of Mesothelin




- Molecule 1: Isoform 4 of Mesothelin




- Molecule 1: Isoform 4 of Mesothelin




- Molecule 1: Isoform 4 of Mesothelin



- Molecule 1: Isoform 4 of Mesothelin



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 3: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain H:  100%

3AG1
3AG2
3AG3

4 Data and refinement statistics i

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	197.17Å 279.32Å 170.05Å 90.00° 125.09° 90.00°	Depositor
Resolution (Å)	49.29 – 3.61 49.29 – 3.61	Depositor EDS
% Data completeness (in resolution range)	91.0 (49.29-3.61) 99.1 (49.29-3.61)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.96 (at 3.57Å)	Xtriage
Refinement program	REFMAC 5.8.0350	Depositor
R, R_{free}	0.291 , 0.310 0.268 , 0.294	Depositor DCC
R_{free} test set	4341 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	99.8	Xtriage
Anisotropy	0.269	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 50.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.41$, $\langle L^2 \rangle = 0.23$	Xtriage
Estimated twinning fraction	0.059 for $-1/2^*h+1/2^*k-1,3/2^*h+1/2^*k+1,1/2^*h-1/2^*k$ 0.056 for $1/2^*h-1/2^*k+1,-1/2^*h+1/2^*k+1,-h$ 0.057 for $1/2^*h+1/2^*k+1,1/2^*h+1/2^*k-1,-h$ 0.054 for $-1/2^*h-1/2^*k-1,-3/2^*h+1/2^*k-1,1/2^*h+1/2^*k$ 0.065 for $-h,-h-2^*1,1/2^*h-1/2^*k$ 0.054 for $-h,h+2^*1,1/2^*h+1/2^*k$ 0.104 for $1/2^*h+1/2^*k+1,3/2^*h-1/2^*k+1,-l$ 0.114 for $1/2^*h-1/2^*k+1,-3/2^*h-1/2^*k-1,-l$ 0.049 for $-1/2^*h+1/2^*k-1,1/2^*h-1/2^*k-1,-1/2^*h-1/2^*k$ 0.065 for $-1/2^*h-1/2^*k-1,-1/2^*h-1/2^*k+1,-1/2^*h+1/2^*k$ 0.064 for $h,-k,-h-l$	Xtriage
Reported twinning fraction	0.318 for H, K, L 0.329 for $1/2H-1/2K+L$, $-3/2H-1/2K-L$, $-L$ 0.298 for $1/2H+1/2K+L$, $3/2H-1/2K+L$, $-L$ 0.055 for H, $-K$, $-H-L$	Depositor
Outliers	0 of 85403 reflections	Xtriage

Continued on next page...

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

Continued from previous page...

Property	Value	Source
F _o ,F _c correlation	0.89	EDS
Total number of atoms	11960	wwPDB-VP
Average B, all atoms (Å ²)	131.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.53% of the height of the origin peak. No significant pseudotranslation is detected.*

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, BMA, NAG, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.26	0/1983	0.45	0/2691
1	B	0.27	0/2006	0.46	0/2723
1	C	0.27	0/2006	0.44	0/2723
1	D	0.27	0/2006	0.45	0/2723
1	E	0.27	0/2006	0.46	0/2723
1	F	0.27	0/2006	0.46	0/2723
All	All	0.27	0/12013	0.45	0/16306

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
1	E	0	1
1	F	0	1
All	All	0	4

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	338	ARG	Sidechain
1	B	338	ARG	Sidechain
1	E	338	ARG	Sidechain

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Group
1	F	417	ARG	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1942	0	1956	25	0
1	B	1963	0	1973	25	0
1	C	1963	0	1973	23	0
1	D	1963	0	1973	39	0
1	E	1963	0	1973	37	0
1	F	1963	0	1973	38	0
2	G	28	0	25	0	0
3	H	39	0	34	0	0
4	A	14	0	13	0	0
4	B	14	0	13	1	0
4	C	14	0	13	0	0
4	E	14	0	13	0	0
5	A	5	0	0	0	0
5	B	10	0	0	0	0
5	C	5	0	0	0	0
5	D	5	0	0	0	0
5	E	15	0	0	0	0
5	F	10	0	0	0	0
6	A	6	0	8	0	0
6	B	6	0	8	0	0
6	C	6	0	8	0	0
6	D	6	0	8	0	0
6	F	6	0	8	0	0
All	All	11960	0	11972	175	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 175 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:535:THR:HG22	1:F:535:THR:HG22	1.49	0.95
1:E:362:GLN:H	1:E:362:GLN:HE21	1.24	0.84
1:A:310:GLU:OE1	1:A:338:ARG:NH1	2.10	0.80
1:D:449:LEU:O	1:D:479:LYS:HE3	1.85	0.77
1:D:497:GLN:NE2	1:D:498:SER:H	1.83	0.76

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	243/312 (78%)	229 (94%)	14 (6%)	0	100	100
1	B	246/312 (79%)	233 (95%)	13 (5%)	0	100	100
1	C	246/312 (79%)	231 (94%)	15 (6%)	0	100	100
1	D	246/312 (79%)	232 (94%)	14 (6%)	0	100	100
1	E	246/312 (79%)	230 (94%)	15 (6%)	1 (0%)	34	71
1	F	246/312 (79%)	230 (94%)	16 (6%)	0	100	100
All	All	1473/1872 (79%)	1385 (94%)	87 (6%)	1 (0%)	51	83

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	E	488	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	217/274 (79%)	211 (97%)	6 (3%)	43	72
1	B	219/274 (80%)	214 (98%)	5 (2%)	50	76
1	C	219/274 (80%)	214 (98%)	5 (2%)	50	76
1	D	219/274 (80%)	213 (97%)	6 (3%)	44	73
1	E	219/274 (80%)	211 (96%)	8 (4%)	34	66
1	F	219/274 (80%)	212 (97%)	7 (3%)	39	70
All	All	1312/1644 (80%)	1275 (97%)	37 (3%)	43	72

5 of 37 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	543	LEU
1	F	417	ARG
1	E	546	HIS
1	F	336	MET
1	C	402	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 7 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	497	GLN
1	E	340	ASN
1	F	540	GLN
1	E	540	GLN
1	B	340	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates i

5 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	G	1	2,1	14,14,15	0.40	0	17,19,21	0.66	0
2	NAG	G	2	2	14,14,15	0.39	0	17,19,21	0.58	0
3	NAG	H	1	3,1	14,14,15	0.40	0	17,19,21	0.56	0
3	NAG	H	2	3	14,14,15	0.40	0	17,19,21	0.64	0
3	BMA	H	3	3	11,11,12	0.25	0	15,15,17	0.50	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	G	1	2,1	-	2/6/23/26	0/1/1/1
2	NAG	G	2	2	-	2/6/23/26	0/1/1/1
3	NAG	H	1	3,1	-	3/6/23/26	0/1/1/1
3	NAG	H	2	3	-	3/6/23/26	0/1/1/1
3	BMA	H	3	3	-	1/2/19/22	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.


5 of 11 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	G	1	NAG	O7-C7-N2-C2
3	H	2	NAG	C3-C2-N2-C7
3	H	2	NAG	C8-C7-N2-C2
3	H	2	NAG	O7-C7-N2-C2
2	G	1	NAG	C8-C7-N2-C2

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

19 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	PO4	A	702	-	4,4,4	0.62	0	6,6,6	0.48	0
6	GOL	B	704	-	5,5,5	0.09	0	5,5,5	0.27	0
4	NAG	A	701	1	14,14,15	0.40	0	17,19,21	0.49	0
6	GOL	D	702	-	5,5,5	0.09	0	5,5,5	0.27	0
6	GOL	A	703	-	5,5,5	0.11	0	5,5,5	0.29	0
5	PO4	B	702	-	4,4,4	0.60	0	6,6,6	0.44	0
6	GOL	F	703	-	5,5,5	0.09	0	5,5,5	0.28	0
5	PO4	B	703	-	4,4,4	0.55	0	6,6,6	0.47	0
5	PO4	F	702	-	4,4,4	0.66	0	6,6,6	0.43	0
5	PO4	F	701	-	4,4,4	0.59	0	6,6,6	0.48	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	PO4	C	702	-	4,4,4	0.63	0	6,6,6	0.43	0
5	PO4	E	702	-	4,4,4	0.68	0	6,6,6	0.41	0
4	NAG	C	701	1	14,14,15	0.39	0	17,19,21	0.62	0
4	NAG	E	701	1	14,14,15	0.39	0	17,19,21	0.65	0
5	PO4	E	703	-	4,4,4	0.61	0	6,6,6	0.45	0
6	GOL	C	703	-	5,5,5	0.10	0	5,5,5	0.27	0
5	PO4	D	701	-	4,4,4	0.67	0	6,6,6	0.43	0
4	NAG	B	701	1	14,14,15	0.42	0	17,19,21	0.61	0
5	PO4	E	704	-	4,4,4	0.66	0	6,6,6	0.44	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	GOL	A	703	-	-	0/4/4/4	-
6	GOL	F	703	-	-	0/4/4/4	-
4	NAG	E	701	1	-	4/6/23/26	0/1/1/1
6	GOL	C	703	-	-	0/4/4/4	-
4	NAG	B	701	1	-	5/6/23/26	0/1/1/1
4	NAG	A	701	1	-	2/6/23/26	0/1/1/1
4	NAG	C	701	1	-	0/6/23/26	0/1/1/1
6	GOL	B	704	-	-	0/4/4/4	-
6	GOL	D	702	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 13 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	701	NAG	C8-C7-N2-C2
4	A	701	NAG	O7-C7-N2-C2
4	B	701	NAG	C3-C2-N2-C7
4	B	701	NAG	C8-C7-N2-C2
4	B	701	NAG	O7-C7-N2-C2

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	701	NAG	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	245/312 (78%)	0.49	4 (1%) 72 57	90, 112, 131, 141	0
1	B	248/312 (79%)	0.54	6 (2%) 59 42	100, 113, 139, 166	0
1	C	248/312 (79%)	0.50	8 (3%) 47 32	98, 120, 137, 172	0
1	D	248/312 (79%)	0.63	17 (6%) 16 10	111, 143, 171, 184	0
1	E	248/312 (79%)	0.52	16 (6%) 18 11	107, 136, 158, 171	0
1	F	248/312 (79%)	0.73	24 (9%) 7 4	136, 164, 182, 191	0
All	All	1485/1872 (79%)	0.57	75 (5%) 28 17	90, 128, 173, 191	0

The worst 5 of 75 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	499	PHE	3.6
1	D	476	LEU	3.4
1	F	425	LEU	3.4
1	C	375	LEU	3.3
1	E	476	LEU	3.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.


6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	NAG	H	1	14/15	0.86	0.28	175,177,180,180	0
3	BMA	H	3	11/12	0.88	0.17	176,176,177,177	0
2	NAG	G	2	14/15	0.91	0.21	164,164,166,167	0
3	NAG	H	2	14/15	0.92	0.18	174,176,178,178	0
2	NAG	G	1	14/15	0.93	0.18	164,166,166,167	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	NAG	C	701	14/15	0.78	0.22	117,122,124,125	0
4	NAG	B	701	14/15	0.84	0.21	124,126,128,129	0
4	NAG	E	701	14/15	0.84	0.15	149,151,154,154	0
4	NAG	A	701	14/15	0.88	0.22	113,118,121,122	0
6	GOL	B	704	6/6	0.91	0.28	49,51,51,52	0
6	GOL	A	703	6/6	0.95	0.29	57,57,58,59	0
6	GOL	C	703	6/6	0.95	0.31	46,48,49,50	0
6	GOL	F	703	6/6	0.95	0.28	46,47,48,49	0
5	PO4	A	702	5/5	0.96	0.27	58,60,61,63	0
6	GOL	D	702	6/6	0.96	0.26	52,53,54,55	0
5	PO4	B	703	5/5	0.96	0.22	66,67,68,69	0
5	PO4	B	702	5/5	0.97	0.28	53,53,54,55	0
5	PO4	C	702	5/5	0.97	0.25	68,69,70,71	0
5	PO4	E	702	5/5	0.97	0.24	63,64,65,65	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
5	PO4	E	703	5/5	0.97	0.24	61,61,63,63	0
5	PO4	F	702	5/5	0.97	0.25	61,61,63,63	0
5	PO4	F	701	5/5	0.98	0.32	53,54,56,56	0
5	PO4	E	704	5/5	0.99	0.26	58,59,60,60	0
5	PO4	D	701	5/5	0.99	0.28	58,60,61,61	0

6.5 Other polymers [i](#)

There are no such residues in this entry.