

Full wwPDB X-ray Structure Validation Report (i)

Feb 25, 2024 – 04:19 PM EST

PDB ID	:	6CYD
Title	:	PDE2 in complex with compound 7
Authors	:	Lu, J.
Deposited on		
Resolution	:	1.69 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:


MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as 541 be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.36
buster-report	:	1.1.7(2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY \, DIFFRACTION$

The reported resolution of this entry is 1.69 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
R_{free}	130704	4298 (1.70-1.70)
Clashscore	141614	4695 (1.70-1.70)
Ramachandran outliers	138981	4610 (1.70-1.70)
Sidechain outliers	138945	4610 (1.70-1.70)
RSRZ outliers	127900	4222 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain			
1	А	373	5% 89%	·	8%	
1	В	373	^{2%} 88%	5%	7%	



2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 6044 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	Λ	343	Total	С	Ν	0	\mathbf{S}	0	1	0
	A	040	2767	1765	466	510	26	0	4	
1	В	348	Total	С	Ν	0	S	0	5	0
	D	340	2840	1809	485	520	26	0	5	0

• Molecule 1 is a protein called cGMP-dependent 3',5'-cyclic phosphodiesterase.

Chain	Residue	Modelled	Actual	Comment	Reference
А	547	MET	-	expression tag	UNP 000408
А	548	HIS	-	expression tag	UNP 000408
А	549	HIS	-	expression tag	UNP 000408
А	550	HIS	-	expression tag	UNP 000408
А	551	HIS	-	expression tag	UNP 000408
A	552	HIS	-	expression tag	UNP 000408
А	553	HIS	-	expression tag	UNP 000408
А	554	GLU	-	expression tag	UNP 000408
А	555	ASN	-	expression tag	UNP 000408
А	556	LEU	-	expression tag	UNP 000408
А	557	TYR	-	expression tag	UNP 000408
А	558	PHE	-	expression tag	UNP 000408
А	559	GLN	-	expression tag	UNP 000408
А	560	GLY	-	expression tag	UNP 000408
А	561	GLU	-	expression tag	UNP 000408
А	562	LEU	-	expression tag	UNP 000408
А	563	SER	-	expression tag	UNP 000408
А	564	THR	-	expression tag	UNP 000408
А	565	SER	-	expression tag	UNP 000408
А	566	LEU	-	expression tag	UNP 000408
А	567	TYR	-	expression tag	UNP 000408
А	568	LYS	-	expression tag	UNP 000408
А	569	LYS	-	expression tag	UNP 000408
А	570	ALA	-	expression tag	UNP 000408
А	571	GLY	-	expression tag	UNP 000408

There are 62 discrepancies between the modelled and reference sequences:

Continued on next page...

6CYD	
------	--

	Continued from previous page									
Chain	Residue	Modelled	Actual	Comment	Reference					
A	572	PHE	-	expression tag	UNP 000408					
A	573	ASP	-	expression tag	UNP 000408					
A	574	ASP	-	expression tag	UNP 000408					
A	575	ASP	-	expression tag	UNP 000408					
А	576	ASP	-	expression tag	UNP 000408					
А	577	LYS	-	expression tag	UNP 000408					
В	547	MET	-	expression tag	UNP 000408					
В	548	HIS	-	expression tag	UNP 000408					
В	549	HIS	-	expression tag	UNP 000408					
В	550	HIS	-	expression tag	UNP 000408					
В	551	HIS	-	expression tag	UNP 000408					
В	552	HIS	-	expression tag	UNP 000408					
В	553	HIS	-	expression tag	UNP 000408					
В	554	GLU	-	expression tag	UNP 000408					
В	555	ASN	-	expression tag	UNP 000408					
В	556	LEU	-	expression tag	UNP 000408					
В	557	TYR	-	expression tag	UNP 000408					
В	558	PHE	-	expression tag	UNP 000408					
В	559	GLN	-	expression tag	UNP 000408					
В	560	GLY	-	expression tag	UNP 000408					
В	561	GLU	-	expression tag	UNP 000408					
В	562	LEU	-	expression tag	UNP 000408					
В	563	SER	-	expression tag	UNP 000408					
В	564	THR	-	expression tag	UNP 000408					
В	565	SER	-	expression tag	UNP 000408					
В	566	LEU	-	expression tag	UNP 000408					
В	567	TYR	-	expression tag	UNP 000408					
В	568	LYS	-	expression tag	UNP 000408					
В	569	LYS	-	expression tag	UNP 000408					
В	570	ALA	-	expression tag	UNP 000408					
В	571	GLY	-	expression tag	UNP 000408					
В	572	PHE	-	expression tag	UNP 000408					
В	573	ASP	-	expression tag	UNP 000408					
В	574	ASP	-	expression tag	UNP 000408					
В	575	ASP	-	expression tag	UNP 000408					
В	576	ASP	-	expression tag	UNP 000408					
В	577	LYS	-	expression tag	UNP 000408					

• Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn). Continued on next page...



 $\mathbf{2}$

В

0

0

Continued from previous page										
Mol	Chain	Residues	Atoms	ZeroOcc	AltConf					
Mol	Chain	Residues	Atoms	ZeroOcc	AltConf					

1

• Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Zn


1

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	А	1	Total Mg 1 1	0	0
3	В	1	Total Mg 1 1	0	0

Total

1


• Molecule 4 is 3-(hydroxymethyl)-6-methyl-1-{(1S)-1-[4-(trifluoromethyl)phenyl]eth yl}-1,5-dihydro-4H-pyrazolo[3,4-d]pyrimidin-4-one (three-letter code: FKG) (formula: $C_{16}H_{15}F_{3}N_{4}O_{2}$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	Δ	1	Total	С	F	Ν	Ο	0	0
4	4 A	1	25	16	3	4	2	0	0
4	р	1	Total	С	F	Ν	Ο	0	0
4	D	I	25	16	3	4	2	0	0

• Molecule 5 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: $C_2H_6O_2$).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	А	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 4 & 2 & 2 \end{array}$	0	0
5	А	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 4 & 2 & 2 \end{array}$	0	0
5	В	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 4 & 2 & 2 \end{array}$	0	0

• Molecule 6 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	А	186	Total O 186 186	0	0
6	В	185	Total O 185 185	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: cGMP-dependent 3',5'-cyclic phosphodiesterase





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	71.94Å 91.82Å 102.83Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.91 - 1.69	Depositor
	44.86 - 1.69	EDS
% Data completeness	99.9 (45.91-1.69)	Depositor
(in resolution range)	99.9(44.86 - 1.69)	EDS
R _{merge}	0.08	Depositor
R _{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	2.36 (at 1.69 Å)	Xtriage
Refinement program	BUSTER 2.11.7	Depositor
R, R_{free}	0.182 , 0.210	Depositor
It, Itfree	0.189 , 0.222	DCC
R_{free} test set	3853 reflections $(5.02%)$	wwPDB-VP
Wilson B-factor $(Å^2)$	20.2	Xtriage
Anisotropy	0.545	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.34 , 47.4	EDS
L-test for twinning ²	$ < L >=0.46, < L^2>=0.28$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6044	wwPDB-VP
Average B, all atoms $(Å^2)$	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 19.25% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, MG, ZN, FKG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain		lengths	Bond angles		
		RMSZ	# Z > 5	RMSZ	# Z > 5	
1	А	0.51	0/2833	0.59	0/3833	
1	В	0.50	0/2907	0.58	0/3926	
All	All	0.51	0/5740	0.59	0/7759	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	2767	0	2643	3	0
1	В	2840	0	2743	8	0
2	А	1	0	0	0	0
2	В	1	0	0	0	0
3	А	1	0	0	0	0
3	В	1	0	0	0	0
4	А	25	0	0	0	0
4	В	25	0	0	0	0
5	А	8	0	12	0	0
5	В	4	0	6	0	0
6	А	186	0	0	0	0

Continued on next page...



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	В	185	0	0	0	0
All	All	6044	0	5404	11	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (11) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:845:MET:HG2	1:B:848:MET:HG2	1.64	0.80
1:B:845:MET:HG2	1:B:848:MET:CG	2.35	0.55
1:A:589:GLY:O	1:A:617:MET:HG2	2.09	0.53
1:B:735:ILE:HD12	1:B:758:LEU:HD13	1.96	0.47
1:A:770:LEU:HD11	1:A:866:ILE:HD12	1.98	0.46
1:B:626:MET:HG2	1:B:672:TYR:CD1	2.53	0.43
1:B:819[A]:THR:HG21	1:B:891:ASN:ND2	2.33	0.43
1:B:671:LEU:HD13	1:B:803:LEU:HD22	2.02	0.42
1:A:834:GLY:HA3	1:A:848:MET:O	2.20	0.41
1:B:902:PHE:HE2	1:B:913:LEU:HD22	1.86	0.41
1:B:571:GLY:HA3	1:B:648:LYS:HE3	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	А	343/373~(92%)	333~(97%)	8 (2%)	2(1%)	25 11
1	В	349/373~(94%)	344 (99%)	5 (1%)	0	100 100
All	All	692/746~(93%)	677 (98%)	13 (2%)	2(0%)	41 24

All (2) Ramachandran outliers are listed below:



Mol	Chain	Res	Type
1	А	899	SER
1	А	915	PHE

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles		
1	А	297/337~(88%)	293~(99%)	4 (1%)	69 56		
1	В	308/337~(91%)	303~(98%)	5(2%)	62 48		
All	All	605/674~(90%)	596~(98%)	9~(2%)	65 51		

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	А	714	VAL
1	А	781	GLN
1	А	898	VAL
1	А	902	PHE
1	В	582	TYR
1	В	613	ASP
1	В	718	LEU
1	В	886	GLU
1	В	892	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	А	630	ASN
1	В	794	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.



5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 9 ligands modelled in this entry, 4 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Iol Type Chain I		Res	Link	Bond lengths			Bond angles		
	ol Type Chain	nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2	
5	EDO	А	1005	-	3,3,3	0.55	0	2,2,2	0.44	0
5	EDO	В	1004	-	3,3,3	0.41	0	2,2,2	0.60	0
4	FKG	В	1003	-	25,27,27	0.86	1 (4%)	29,41,41	1.11	2 (6%)
5	EDO	А	1004	-	3,3,3	0.67	0	2,2,2	0.39	0
4	FKG	А	1003	-	25,27,27	0.63	0	29,41,41	1.06	1 (3%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	EDO	А	1005	-	-	0/1/1/1	-
5	EDO	В	1004	-	-	0/1/1/1	-
4	FKG	В	1003	-	-	1/12/16/16	0/3/3/3
5	EDO	А	1004	-	-	0/1/1/1	-
4	FKG	А	1003	-	-	1/12/16/16	0/3/3/3

All (1) bond length outliers are listed below:



Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	В	1003	FKG	C14-N3	3.01	1.37	1.31

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
4	А	1003	FKG	C12-C2-C1	-4.75	122.23	130.34
4	В	1003	FKG	C12-C2-C1	-4.53	122.61	130.34
4	В	1003	FKG	C2-N-N1	3.01	106.43	104.24

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	А	1003	FKG	C11-C3-C4-C5
4	В	1003	FKG	C11-C3-C4-C5


There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	$\langle RSRZ \rangle$	#RSRZ>2	$OWAB(Å^2)$	Q<0.9
1	А	343/373~(91%)	0.10	19 (5%) 25 27	13, 23, 49, 84	0
1	В	348/373~(93%)	0.06	8 (2%) 60 65	14, 23, 45, 73	0
All	All	691/746~(92%)	0.08	27 (3%) 39 44	13, 23, 47, 84	0

All (27) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	А	902	PHE	13.1
1	А	916	LEU	6.7
1	В	581	GLU	6.3
1	А	586	LEU	6.1
1	В	583	THR	6.1
1	В	562	LEU	5.3
1	В	587	HIS	5.1
1	А	588	ASP	5.1
1	А	903	THR	5.1
1	А	904	ILE	4.9
1	А	901	LYS	4.5
1	В	582	TYR	4.1
1	А	915	PHE	3.9
1	А	582	TYR	3.9
1	В	915	PHE	3.7
1	А	910	ASN	3.3
1	В	588	ASP	3.2
1	А	909	SER	3.2
1	А	854	TYR	2.9
1	В	585	LEU	2.7
1	А	589	GLY	2.7
1	А	583	THR	2.6
1	А	587	HIS	2.4
1	А	898	VAL	2.4

Continued on next page...



Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	А	585	LEU	2.3
1	А	900	HIS	2.1
1	А	843	ARG	2.1

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.


6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} extsf{-}\mathbf{B} extsf{-}\mathbf{factors}(\mathbf{A}^2)$	Q<0.9
5	EDO	А	1005	4/4	0.92	0.13	32,36,41,44	0
5	EDO	А	1004	4/4	0.94	0.09	26,28,29,29	0
4	FKG	А	1003	25/25	0.94	0.07	17,19,23,29	0
4	FKG	В	1003	25/25	0.95	0.07	17,20,27,31	0
5	EDO	В	1004	4/4	0.95	0.16	27,29,36,44	0
3	MG	В	1002	1/1	0.98	0.06	18,18,18,18	0
3	MG	А	1002	1/1	0.99	0.04	16,16,16,16	0
2	ZN	В	1001	1/1	1.00	0.03	20,20,20,20	0
2	ZN	А	1001	1/1	1.00	0.03	19,19,19,19	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.









6.5 Other polymers (i)

There are no such residues in this entry.

