



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 7, 2023 – 03:42 PM EDT

PDB ID : 8CYN
Title : [2T7+20bp Linker] Self-Assembled 3D DNA Hexagonal Tensegrity Triangle with 20 bp Sticky-End Linker
Authors : Lu, B.; Vecchioni, S.; Ohayon, Y.P.; Seeman, N.C.; Mao, C.; Sha, R.
Deposited on : 2022-05-24
Resolution : 9.45 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtrriage (Phenix) : 1.13
EDS : 2.35
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 9.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1005 (11.50-3.90)
Clashscore	141614	1070 (11.50-3.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	21	
2	B	7	
3	C	6	
4	D	8	
5	X	20	
6	Y	20	

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 1675 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called DNA (5'-D(*GP*CP*GP*CP*AP*GP*CP*CP*TP*GP*TP*AP*CP*GP*GP*AP*CP*AP*TP*CP*A)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
1	A	21	427	203	82	122	20	0	0	0

- Molecule 2 is a DNA chain called DNA (5'-D(P*CP*CP*GP*TP*AP*CP*A)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
2	B	7	141	67	26	41	7	0	0	0

- Molecule 3 is a DNA chain called DNA (5'-D(P*GP*GP*CP*TP*GP*C)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
3	C	6	124	58	23	37	6	0	0	0

- Molecule 4 is a DNA chain called DNA (5'-D(*TP*CP*TP*GP*AP*TP*GP*T)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
4	D	8	161	79	26	49	7	0	0	0

- Molecule 5 is a DNA chain called DNA (5'-D(P*GP*AP*TP*GP*CP*TP*GP*AP*CP*GP*TP*AP*GP*TP*AP*GP*CP*AP*GP*A)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
5	X	20	417	197	82	118	20	0	0	0

- Molecule 6 is a DNA chain called DNA (5'-D(P*GP*CP*TP*CP*TP*GP*CP*TP*AP*CP*TP*AP*CP*GP*TP*CP*AP*GP*CP*A)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
6	Y	20	405	193	71	121	20	0	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DNA (5'-D(*GP*CP*GP*CP*AP*GP*CP*CP*TP*GP*TP*AP*CP*GP*GP*AP*CP*AP*TP*CP*A)-3')

Chain A:  19% 81%

G101	T111
C102	A112
G103	C113
C104	G114
A105	G115
G106	A116
C107	C117
C108	A118
	T119
	C120
	A121

- Molecule 2: DNA (5'-D(P*CP*CP*GP*TP*AP*CP*A)-3')

Chain B:  57% 43%

C119	A125
C120	
G121	
T122	

- Molecule 3: DNA (5'-D(P*GP*GP*CP*TP*GP*C)-3')

Chain C:  17% 83%


G209	G213
G210	C214
C211	
T212	

- Molecule 4: DNA (5'-D(*TP*CP*TP*GP*AP*TP*GP*T)-3')

Chain D:  75% 25%


T201	T208
T206	
G207	

- Molecule 5: DNA (5'-D(P*GP*AP*TP*GP*CP*TP*GP*AP*CP*GP*TP*AP*GP*TP*AP*GP*CP*AP*GP*A)-3')

Chain X:  90% 10%

G122	A141
T127	
G128	

- Molecule 6: DNA (5'-D(P*GP*CP*TP*CP*TP*GP*CP*TP*AP*CP*TP*AP*CP*GP*TP*CP*AP*GP*CP*A)-3')



4 Data and refinement statistics i

Property	Value	Source
Space group	P 63	Depositor
Cell constants a, b, c, α , β , γ	231.03Å 231.03Å 100.85Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	50.02 – 9.45 90.06 – 9.45	Depositor EDS
% Data completeness (in resolution range)	63.2 (50.02-9.45) 59.5 (90.06-9.45)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$	-	Xtrriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, R_{free}	0.096 , 0.144 0.097 , 0.145	Depositor DCC
R_{free} test set	69 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	(Not available)	Xtrriage
Anisotropy	(Not available)	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	1.24 , 32.1	EDS
L-test for twinning ¹	$\langle L \rangle =$ (Not available), $\langle L^2 \rangle =$ (Not available)	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.62	EDS
Total number of atoms	1675	wwPDB-VP
Average B, all atoms (Å ²)	977.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *(Not available)*

¹Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.66	0/479	0.98	0/737
2	B	0.57	0/157	0.90	0/239
3	C	0.61	0/138	1.03	0/211
4	D	0.56	0/179	1.09	0/275
5	X	0.52	0/469	0.89	0/723
6	Y	0.59	0/452	0.98	0/694
All	All	0.59	0/1874	0.97	0/2879

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	427	0	236	19	0
2	B	141	0	79	2	0
3	C	124	0	68	5	0
4	D	161	0	94	1	0
5	X	417	0	225	1	0
6	Y	405	0	226	5	0
All	All	1675	0	928	32	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (32) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:112:DA:H2'	1:A:113:DC:C6	2.28	0.69
1:A:111:DT:H2''	1:A:112:DA:C8	2.35	0.61
1:A:120:DC:H2'	1:A:121:DA:C8	2.36	0.60
6:Y:113:DA:H2'	6:Y:114:DC:C6	2.40	0.56
2:B:121:DG:H2'	2:B:122:DT:H71	1.87	0.56
1:A:103:DG:H22	3:C:214:DC:H42	1.53	0.55
1:A:104:DC:H1'	1:A:105:DA:C5	2.42	0.55
3:C:209:DG:H2'	3:C:210:DG:C8	2.42	0.54
1:A:113:DC:H2''	1:A:114:DG:C5	2.45	0.51
1:A:115:DG:N3	1:A:116:DA:N7	2.59	0.51
2:B:120:DC:H2''	2:B:121:DG:N7	2.24	0.51
1:A:103:DG:H2''	1:A:104:DC:H2'	1.93	0.51
1:A:118:DA:H2''	1:A:119:DT:H71	1.94	0.50
3:C:211:DC:H2''	3:C:212:DT:O5'	2.14	0.48
6:Y:102:DG:H2''	6:Y:103:DC:OP2	2.12	0.48
6:Y:102:DG:H1'	6:Y:103:DC:H4'	1.95	0.48
1:A:107:DC:H2''	1:A:108:DC:H5'	1.95	0.47
5:X:127:DT:H2''	5:X:128:DG:C8	2.49	0.47
1:A:114:DG:H2''	1:A:115:DG:C8	2.51	0.46
3:C:211:DC:H2'	3:C:212:DT:H72	1.98	0.46
1:A:101:DG:H2'	1:A:102:DC:C6	2.50	0.46
1:A:113:DC:H2''	1:A:114:DG:N7	2.31	0.45
6:Y:108:DC:H2''	6:Y:109:DT:C5	2.50	0.45
3:C:211:DC:H4'	3:C:212:DT:OP1	2.17	0.45
1:A:103:DG:H1'	1:A:104:DC:H5'	1.99	0.44
6:Y:109:DT:H2''	6:Y:110:DA:H5''	1.99	0.44
1:A:112:DA:H2''	1:A:113:DC:O5'	2.18	0.43
4:D:206:DT:H2''	4:D:207:DG:N7	2.33	0.43
1:A:112:DA:H2''	1:A:113:DC:C5'	2.48	0.43
1:A:114:DG:H2''	1:A:115:DG:OP2	2.19	0.43
1:A:115:DG:H1'	1:A:116:DA:C8	2.55	0.41
1:A:111:DT:H2''	1:A:112:DA:H8	1.82	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

There are no protein molecules in this entry.

5.3.2 Protein sidechains [i](#)

There are no protein molecules in this entry.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

6.3 Carbohydrates

Unable to reproduce the depositors R factor - this section is therefore empty.

6.4 Ligands

Unable to reproduce the depositors R factor - this section is therefore empty.

6.5 Other polymers

Unable to reproduce the depositors R factor - this section is therefore empty.