



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 30, 2023 – 05:35 PM EDT

PDB ID : 8D2J
Title : A novel insecticidal protein from ferns IPD113_Cow.
Authors : Maher, M.J.
Deposited on : 2022-05-30
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	512	<div style="display: flex; align-items: center;"> <div style="width: 100%; height: 15px; background: linear-gradient(to right, red 1%, orange 1%, yellow 1%, green 87%, grey 87%);"></div> <div style="margin-left: 10px;"> <p>87% 6% 6%</p> </div> </div>
1	B	512	<div style="display: flex; align-items: center;"> <div style="width: 100%; height: 15px; background: linear-gradient(to right, red 3%, orange 3%, yellow 1%, green 88%, grey 88%);"></div> <div style="margin-left: 10px;"> <p>88% 6% 6%</p> </div> </div>
1	C	512	<div style="display: flex; align-items: center;"> <div style="width: 100%; height: 15px; background: linear-gradient(to right, red 4%, orange 4%, yellow 1%, green 87%, grey 87%);"></div> <div style="margin-left: 10px;"> <p>87% 6% 7%</p> </div> </div>
1	D	512	<div style="display: flex; align-items: center;"> <div style="width: 100%; height: 15px; background: linear-gradient(to right, red 3%, orange 3%, yellow 1%, green 89%, grey 89%);"></div> <div style="margin-left: 10px;"> <p>89% 5% 6%</p> </div> </div>

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 16286 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called IPD113_Cow.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	B	483	Total 3919	C 2483	N 689	O 733	S 14	0	5	0
1	A	479	Total 3862	C 2447	N 682	O 719	S 14	0	1	0
1	C	478	Total 3860	C 2445	N 681	O 720	S 14	0	1	0
1	D	480	Total 3876	C 2455	N 683	O 724	S 14	0	0	0

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
2	A	1	Total 6	C 3	O 3	0	0
2	C	1	Total 6	C 3	O 3	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	C	1	Total	C	O	0	0
			6	3	3		
2	D	1	Total	C	O	0	0
			6	3	3		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	204	Total	O	0	0
			204	204		
3	A	221	Total	O	0	0
			221	221		
3	C	175	Total	O	0	0
			175	175		
3	D	145	Total	O	0	0
			145	145		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: IPD113_Cow




- Molecule 1: IPD113_Cow




- Molecule 1: IPD113_Cow



- Molecule 1: IPD113_Cow





4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	65.64Å 79.39Å 108.55Å 97.92° 104.95° 97.41°	Depositor
Resolution (Å)	47.82 – 2.10 47.82 – 1.98	Depositor EDS
% Data completeness (in resolution range)	97.2 (47.82-2.10) 96.3 (47.82-1.98)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.39 (at 1.98Å)	Xtrriage
Refinement program	REFMAC 5.8.0258	Depositor
R, R_{free}	0.220 , 0.256 0.224 , 0.260	Depositor DCC
R_{free} test set	6955 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	27.3	Xtrriage
Anisotropy	0.434	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 38.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	16286	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.30% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.64	0/3959	0.70	0/5381
1	B	0.64	0/4030	0.70	0/5479
1	C	0.64	0/3956	0.69	0/5376
1	D	0.64	0/3970	0.69	0/5395
All	All	0.64	0/15915	0.70	0/21631

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3862	0	3754	22	0
1	B	3919	0	3815	20	0
1	C	3860	0	3752	19	0
1	D	3876	0	3766	18	0
2	A	6	0	8	0	0
2	C	12	0	16	1	0
2	D	6	0	8	0	0
3	A	221	0	0	1	0
3	B	204	0	0	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	C	175	0	0	2	0
3	D	145	0	0	2	0
All	All	16286	0	15119	78	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (78) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:253:ASN:O	1:B:254:LEU:HB3	1.85	0.77
1:C:104:LEU:O	1:C:110:ILE:HD11	1.89	0.72
1:B:98:VAL:CG1	1:B:102:GLU:HB3	2.29	0.63
1:B:107[B]:GLU:OE2	1:B:107[B]:GLU:HA	1.99	0.61
1:D:422:ARG:O	3:D:701:HOH:O	2.15	0.60
1:D:152:LEU:HD22	3:D:702:HOH:O	2.01	0.59
1:A:104:LEU:HD11	1:A:177:LEU:HB3	1.86	0.58
1:B:205:ASN:HD21	1:B:208:ASN:HD22	1.51	0.57
1:A:116:GLN:HA	1:A:119:PHE:CE2	2.39	0.57
1:A:98:VAL:HG13	1:A:102:GLU:HB3	1.87	0.55
1:A:107[B]:GLU:OE2	1:A:107[B]:GLU:HA	2.07	0.55
1:B:253:ASN:O	1:B:254:LEU:CB	2.56	0.54
1:B:390:PHE:CD2	3:B:720:HOH:O	2.54	0.54
1:C:104:LEU:HD11	1:C:177:LEU:HB3	1.88	0.54
1:D:26:GLU:OE1	1:D:26:GLU:N	2.41	0.53
1:D:152:LEU:O	1:D:156:ILE:HG12	2.08	0.53
1:C:98:VAL:HG13	1:C:102:GLU:HB3	1.91	0.53
1:C:107[B]:GLU:OE2	1:C:107[B]:GLU:HA	2.09	0.53
1:C:427:LEU:CD2	1:C:441:LEU:HD22	2.38	0.53
1:B:417:VAL:HG11	1:B:441:LEU:HD12	1.91	0.53
1:C:101:TRP:O	1:C:104:LEU:HB3	2.08	0.53
1:D:223:ASP:OD1	1:D:227:ASN:ND2	2.42	0.53
1:B:98:VAL:HG13	1:B:102:GLU:HB3	1.90	0.52
1:C:98:VAL:CG1	1:C:102:GLU:HB3	2.40	0.52
1:D:98:VAL:CG1	1:D:102:GLU:HB3	2.40	0.52
1:A:193:ARG:NH2	1:A:212:ASP:OD2	2.42	0.51
1:D:225:ILE:HB	1:D:226:PRO:HD3	1.95	0.49
1:A:101:TRP:O	1:A:104:LEU:HB3	2.13	0.49
1:D:104:LEU:HD11	1:D:177:LEU:HB3	1.95	0.48
1:C:116:GLN:HA	1:C:119:PHE:CE2	2.49	0.48
1:B:257:ARG:NH1	1:A:99:ASP:OD1	2.46	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:311:SER:N	1:B:312:PRO:CD	2.77	0.48
1:A:365:ARG:NH1	3:A:702:HOH:O	2.46	0.48
1:C:341:ARG:NH2	3:C:701:HOH:O	2.46	0.48
1:D:311:SER:N	1:D:312:PRO:CD	2.76	0.47
1:A:426:GLU:HB2	1:A:442:HIS:HB3	1.96	0.47
1:A:219:THR:O	1:A:223:ASP:HB3	2.14	0.46
1:C:225:ILE:HB	1:C:226:PRO:HD3	1.97	0.46
1:A:378:ILE:HD12	1:A:378:ILE:N	2.31	0.46
1:C:311:SER:N	1:C:312:PRO:CD	2.78	0.46
1:A:253:ASN:O	1:A:254:LEU:CB	2.63	0.45
1:C:139:ARG:O	1:C:143:ALA:HB2	2.17	0.45
1:C:219:THR:O	1:C:223:ASP:HB3	2.17	0.45
1:A:311:SER:N	1:A:312:PRO:CD	2.80	0.45
1:A:104:LEU:HD11	1:A:177:LEU:CB	2.46	0.45
1:C:419:VAL:HG11	1:C:424:ILE:HD11	1.97	0.45
1:D:193:ARG:NH2	1:D:212:ASP:OD2	2.49	0.45
1:B:235:TRP:HZ2	1:B:239:GLN:HE21	1.64	0.44
1:A:225:ILE:HB	1:A:226:PRO:HD3	1.98	0.44
1:A:427:LEU:HD22	1:A:441:LEU:HD22	2.00	0.44
1:C:160:ASP:O	1:C:164:ILE:HG12	2.17	0.44
1:B:107[B]:GLU:OE2	1:B:107[B]:GLU:CA	2.66	0.43
1:A:243:SER:O	1:A:263:ALA:HA	2.19	0.43
1:B:66:ARG:HD3	1:B:91:LEU:O	2.19	0.43
2:C:602:GOL:H12	3:C:710:HOH:O	2.17	0.43
1:B:225:ILE:HB	1:B:226:PRO:HD3	2.01	0.42
1:C:109:TYR:HB3	1:C:305:ASP:HB3	1.99	0.42
1:C:360:ARG:HD3	1:C:491:SER:HB2	2.01	0.42
1:B:223:ASP:OD1	1:B:227:ASN:ND2	2.51	0.42
1:D:378:ILE:HD12	1:D:378:ILE:N	2.35	0.42
1:B:99:ASP:OD1	1:A:257:ARG:NH1	2.53	0.42
1:A:427:LEU:CD2	1:A:441:LEU:HD22	2.49	0.42
1:D:378:ILE:HD13	1:D:391:VAL:HG23	2.02	0.42
1:B:219:THR:O	1:B:223:ASP:HB3	2.20	0.42
1:A:98:VAL:CG1	1:A:102:GLU:HB3	2.50	0.42
1:D:104:LEU:HD11	1:D:177:LEU:CB	2.50	0.41
1:A:109:TYR:HB3	1:A:305:ASP:HB3	2.02	0.41
1:C:104:LEU:HD11	1:C:177:LEU:CB	2.50	0.41
1:A:253:ASN:O	1:A:254:LEU:HB3	2.21	0.41
1:D:206:SER:HA	1:D:209:TRP:CE3	2.55	0.41
1:B:116:GLN:HA	1:B:119:PHE:CE2	2.55	0.41
1:D:98:VAL:HG13	1:D:102:GLU:HB3	2.03	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:376:GLU:HB2	1:D:391:VAL:HB	2.03	0.41
1:D:236:ARG:N	1:D:237:PRO:CD	2.85	0.40
1:B:427:LEU:CD2	1:B:441:LEU:HD22	2.51	0.40
1:D:360:ARG:HD3	1:D:491:SER:HB2	2.04	0.40
1:B:419:VAL:HG11	1:B:424:ILE:HD11	2.03	0.40
1:C:115:GLN:HE21	1:C:235:TRP:HZ2	1.70	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	476/512 (93%)	463 (97%)	11 (2%)	2 (0%)	34	32
1	B	486/512 (95%)	473 (97%)	11 (2%)	2 (0%)	34	32
1	C	475/512 (93%)	465 (98%)	9 (2%)	1 (0%)	47	49
1	D	476/512 (93%)	465 (98%)	11 (2%)	0	100	100
All	All	1913/2048 (93%)	1866 (98%)	42 (2%)	5 (0%)	41	41

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	254	LEU
1	A	254	LEU
1	C	79	PRO
1	B	450	ARG
1	A	79	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	414/440 (94%)	413 (100%)	1 (0%)	93	96
1	B	423/440 (96%)	419 (99%)	4 (1%)	78	84
1	C	414/440 (94%)	414 (100%)	0	100	100
1	D	416/440 (94%)	415 (100%)	1 (0%)	93	96
All	All	1667/1760 (95%)	1661 (100%)	6 (0%)	92	94

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	119	PHE
1	B	167	ARG
1	B	393[A]	ASP
1	B	393[B]	ASP
1	A	205	ASN
1	D	87	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	358	ASN
1	D	115	GLN
1	D	128	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GOL	A	601	-	5,5,5	0.08	0	5,5,5	0.27	0
2	GOL	D	601	-	5,5,5	0.09	0	5,5,5	0.29	0
2	GOL	C	602	-	5,5,5	0.08	0	5,5,5	0.26	0
2	GOL	C	601	-	5,5,5	0.07	0	5,5,5	0.24	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	601	-	-	2/4/4/4	-
2	GOL	D	601	-	-	0/4/4/4	-
2	GOL	C	602	-	-	0/4/4/4	-
2	GOL	C	601	-	-	0/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	601	GOL	C1-C2-C3-O3
2	A	601	GOL	O1-C1-C2-C3

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	602	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	479/512 (93%)	0.11	6 (1%) 77 80	21, 30, 45, 57	0
1	B	483/512 (94%)	0.17	13 (2%) 54 60	23, 32, 49, 63	0
1	C	478/512 (93%)	0.31	20 (4%) 36 42	24, 35, 54, 77	0
1	D	480/512 (93%)	0.32	15 (3%) 49 55	26, 36, 52, 73	0
All	All	1920/2048 (93%)	0.23	54 (2%) 53 59	21, 33, 51, 77	0

All (54) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	47	VAL	6.9
1	C	76	SER	6.4
1	D	77	ALA	6.0
1	B	77	ALA	5.8
1	C	256	CYS	5.4
1	C	46	GLU	5.4
1	C	78	VAL	5.2
1	D	256	CYS	4.6
1	C	369	THR	4.3
1	C	45	GLU	4.0
1	C	79	PRO	3.7
1	C	398	ASN	3.7
1	D	42	ALA	3.5
1	B	78	VAL	3.4
1	B	253	ASN	3.4
1	D	231	ARG	3.1
1	B	79	PRO	3.1
1	D	46	GLU	3.0
1	D	251	CYS	3.0
1	D	333	ARG	2.9
1	B	508	SER	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	256	CYS	2.9
1	C	77	ALA	2.9
1	B	251	CYS	2.8
1	A	508	SER	2.8
1	B	249	GLY	2.8
1	A	398	ASN	2.8
1	C	370	LEU	2.7
1	A	253	ASN	2.7
1	C	43	GLU	2.5
1	C	48	PRO	2.5
1	B	76	SER	2.4
1	C	486	ARG	2.4
1	A	204	ASN	2.4
1	B	252	GLY	2.4
1	B	421	ASP	2.3
1	D	124	ARG	2.3
1	A	111	ARG	2.3
1	D	36	GLN	2.3
1	D	224	LEU	2.2
1	D	171	ASN	2.2
1	D	393	ASP	2.2
1	C	111	ARG	2.2
1	A	121	PHE	2.2
1	B	398	ASN	2.2
1	D	47	VAL	2.1
1	C	507	ALA	2.1
1	C	251	CYS	2.1
1	C	247	ARG	2.1
1	D	29	PRO	2.1
1	D	40	ALA	2.1
1	C	323	ASP	2.1
1	B	254	LEU	2.1
1	C	384	HIS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	GOL	A	601	6/6	0.79	0.18	45,46,46,47	0
2	GOL	D	601	6/6	0.80	0.18	39,39,39,41	0
2	GOL	C	601	6/6	0.82	0.21	39,40,40,41	0
2	GOL	C	602	6/6	0.87	0.20	39,41,41,41	0

6.5 Other polymers [i](#)

There are no such residues in this entry.