



# Full wwPDB X-ray Structure Validation Report ⓘ

May 21, 2020 – 07:03 pm BST

PDB ID : 1D48  
Title : STRUCTURE OF THE PURE-SPERMINE FORM OF Z-DNA (MAGNESIUM FREE) AT 1 ANGSTROM RESOLUTION  
Authors : Egli, M.; Williams, L.D.; Gao, Q.; Rich, A.  
Deposited on : 1991-09-11  
Resolution : 1.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtrriage (Phenix) : **NOT EXECUTED**  
EDS : **NOT EXECUTED**  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.00 Å.



Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	1117 (1.06-0.94)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	6	 100%
1	B	6	 100%

## 2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 301 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called DNA (5'-D(\*CP\*GP\*CP\*GP\*CP\*G)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
1	A	6	Total 120	C 57	N 24	O 34	P 5	0	0	0
1	B	6	Total 120	C 57	N 24	O 34	P 5	0	0	0

- Molecule 2 is SPERMINE (three-letter code: SPM) (formula: C<sub>10</sub>H<sub>26</sub>N<sub>4</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	N		
2	A	1	Total 14	C 10	N 4	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
3	A	20	Total 20	O 20	0	0

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
3	B	27	Total	O	0	0
			27	27		


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


Note EDS was not executed.


- Molecule 1: DNA (5'-D(\*CP\*GP\*CP\*GP\*CP\*G)-3')

Chain A:  100%

 G1 G2 G3 G4 G5 G6

- Molecule 1: DNA (5'-D(\*CP\*GP\*CP\*GP\*CP\*G)-3')

Chain B:  100%

 G7 G8 G9 G10 G11 G12

## 4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value
Space group	P 21 21 21
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	18.41Å 30.77Å 43.15Å 90.00° 90.00° 90.00°
Resolution (Å)	(Not available) – 1.00
% Data completeness (in resolution range)	(Not available) ((Not available)-1.00)
$R_{merge}$	(Not available)
$R_{sym}$	(Not available)
Refinement program	PROLSQ (MODIFIED BY G.J.QUIGLEY), MODIFIED BY G.J.QUIGLEY
R, $R_{free}$	0.185 , (Not available)
Estimated twinning fraction	No twinning to report.
Total number of atoms	301
Average B, all atoms (Å <sup>2</sup> )	12.0

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: SPM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	3.63	26/134 (19.4%)	4.02	35/205 (17.1%)
1	B	3.45	16/134 (11.9%)	3.88	31/205 (15.1%)
All	All	3.54	42/268 (15.7%)	3.95	66/410 (16.1%)

All (42) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	6	DG	C6-N1	-9.92	1.32	1.39
1	A	6	DG	C2-N2	-9.35	1.25	1.34
1	A	2	DG	C2-N2	-9.10	1.25	1.34
1	B	10	DG	C2-N2	-8.97	1.25	1.34
1	B	8	DG	C2-N2	-8.84	1.25	1.34
1	B	9	DC	C2'-C1'	8.04	1.60	1.52
1	B	12	DG	C2-N2	-7.86	1.26	1.34
1	A	4	DG	C2'-C1'	7.69	1.60	1.52
1	A	4	DG	C6-N1	-7.68	1.34	1.39
1	B	10	DG	C6-N1	-7.60	1.34	1.39
1	A	4	DG	C2-N2	-7.49	1.27	1.34
1	A	2	DG	C4'-O4'	-7.20	1.37	1.45
1	B	12	DG	C6-N1	-7.03	1.34	1.39
1	B	8	DG	C6-N1	-6.90	1.34	1.39
1	A	3	DC	C2'-C1'	6.85	1.59	1.52
1	A	2	DG	C6-N1	-6.84	1.34	1.39
1	A	3	DC	N3-C4	6.79	1.38	1.33
1	A	2	DG	P-OP2	-6.73	1.37	1.49
1	B	8	DG	C8-N7	6.68	1.34	1.30
1	B	11	DC	N3-C4	6.63	1.38	1.33
1	B	8	DG	N9-C8	-6.52	1.33	1.37
1	B	9	DC	N3-C4	6.42	1.38	1.33
1	B	7	DC	N3-C4	6.42	1.38	1.33
1	A	4	DG	N7-C5	-6.33	1.35	1.39

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	2	DG	C6-O6	6.30	1.29	1.24
1	A	1	DC	N3-C4	6.20	1.38	1.33
1	B	10	DG	P-O5'	6.10	1.65	1.59
1	A	4	DG	N9-C8	-6.01	1.33	1.37
1	A	6	DG	C4'-C3'	5.93	1.59	1.53
1	B	10	DG	C6-O6	5.92	1.29	1.24
1	B	8	DG	C5'-C4'	5.67	1.57	1.51
1	B	12	DG	C4'-C3'	5.67	1.59	1.53
1	A	2	DG	C4'-C3'	5.43	1.58	1.53
1	A	6	DG	C8-N7	5.32	1.34	1.30
1	A	6	DG	N9-C4	5.25	1.42	1.38
1	A	1	DC	C3'-O3'	-5.25	1.37	1.44
1	A	2	DG	C3'-C2'	-5.23	1.46	1.52
1	A	6	DG	C1'-N9	-5.20	1.40	1.47
1	A	6	DG	P-OP2	-5.20	1.40	1.49
1	A	2	DG	C2-N3	5.19	1.36	1.32
1	A	1	DC	C2'-C1'	5.15	1.57	1.52
1	A	3	DC	P-OP1	-5.08	1.40	1.49

All (66) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	5	DC	N3-C4-C5	-13.87	116.35	121.90
1	A	1	DC	N3-C4-C5	-13.34	116.56	121.90
1	A	2	DG	C5-C6-N1	13.18	118.09	111.50
1	A	5	DC	C2-N3-C4	13.09	126.44	119.90
1	A	2	DG	N3-C2-N2	-12.67	111.03	119.90
1	B	10	DG	N3-C2-N2	-12.59	111.09	119.90
1	A	2	DG	C5-C6-O6	-11.64	121.62	128.60
1	B	10	DG	C5-C6-N1	11.39	117.19	111.50
1	B	8	DG	C5-C6-N1	11.31	117.15	111.50
1	A	6	DG	C5-C6-N1	10.83	116.92	111.50
1	B	8	DG	N3-C2-N2	-10.55	112.51	119.90
1	A	1	DC	N1-C2-O2	10.37	125.12	118.90
1	B	10	DG	C5-C6-O6	-10.06	122.56	128.60
1	B	11	DC	N3-C4-C5	-9.97	117.91	121.90
1	B	12	DG	N3-C2-N2	-9.96	112.93	119.90
1	B	9	DC	N1-C2-O2	9.66	124.70	118.90
1	A	2	DG	C6-N1-C2	-9.61	119.33	125.10
1	A	6	DG	N3-C2-N2	-9.58	113.19	119.90
1	B	8	DG	C5-C6-O6	-9.37	122.98	128.60
1	A	1	DC	C2-N3-C4	9.16	124.48	119.90

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	12	DG	C5-C6-O6	-9.10	123.14	128.60
1	A	4	DG	N3-C2-N2	-9.02	113.58	119.90
1	A	6	DG	C4-C5-N7	8.77	114.31	110.80
1	B	7	DC	N3-C4-C5	-8.63	118.45	121.90
1	B	12	DG	C5-C6-N1	8.61	115.80	111.50
1	B	8	DG	C6-N1-C2	-8.53	119.98	125.10
1	B	8	DG	N1-C2-N3	8.52	129.01	123.90
1	A	5	DC	N1-C2-O2	8.47	123.98	118.90
1	B	12	DG	N1-C2-N3	8.29	128.87	123.90
1	B	11	DC	C2-N3-C4	8.27	124.03	119.90
1	A	6	DG	C5-C6-O6	-8.17	123.70	128.60
1	B	9	DC	N3-C4-C5	-8.02	118.69	121.90
1	A	4	DG	C5-C6-N1	7.93	115.46	111.50
1	B	12	DG	C6-N1-C2	-7.61	120.53	125.10
1	B	10	DG	C6-N1-C2	-7.55	120.57	125.10
1	A	2	DG	N1-C2-N3	7.37	128.32	123.90
1	B	11	DC	N1-C2-O2	7.36	123.32	118.90
1	A	1	DC	C5-C4-N4	7.02	125.11	120.20
1	A	6	DG	C6-N1-C2	-6.96	120.92	125.10
1	A	3	DC	N1-C2-O2	6.93	123.06	118.90
1	A	4	DG	C5-C6-O6	-6.89	124.46	128.60
1	A	3	DC	N3-C4-C5	-6.76	119.20	121.90
1	B	11	DC	C5-C4-N4	6.71	124.90	120.20
1	B	8	DG	C2-N3-C4	-6.67	108.57	111.90
1	B	10	DG	N1-C2-N2	6.47	122.02	116.20
1	B	12	DG	C2-N3-C4	-6.38	108.71	111.90
1	B	9	DC	C2-N3-C4	6.20	123.00	119.90
1	A	4	DG	N1-C2-N3	6.15	127.59	123.90
1	A	4	DG	C4-C5-N7	5.66	113.06	110.80
1	A	6	DG	N1-C2-N3	5.63	127.28	123.90
1	A	4	DG	C6-N1-C2	-5.63	121.72	125.10
1	B	11	DC	N1-C2-N3	-5.58	115.30	119.20
1	A	5	DC	N1-C2-N3	-5.55	115.32	119.20
1	A	6	DG	N9-C4-C5	-5.54	103.18	105.40
1	A	3	DC	C2-N3-C4	5.54	122.67	119.90
1	B	7	DC	N1-C2-O2	5.50	122.20	118.90
1	A	1	DC	C5-C6-N1	5.44	123.72	121.00
1	B	11	DC	C6-N1-C2	5.27	122.41	120.30
1	A	2	DG	C4-C5-C6	-5.25	115.65	118.80
1	A	1	DC	P-O3'-C3'	5.24	125.99	119.70
1	B	7	DC	C2-N3-C4	5.18	122.49	119.90
1	A	2	DG	C2-N3-C4	-5.17	109.31	111.90

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	A	5	DC	O4'-C1'-C2'	-5.12	101.81	105.90
1	B	7	DC	P-O3'-C3'	5.11	125.83	119.70
1	A	4	DG	N9-C4-C5	-5.06	103.38	105.40
1	B	12	DG	C4-C5-N7	5.01	112.80	110.80

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	120	0	62	0	0
1	B	120	0	62	0	0
2	A	14	0	26	0	0
3	A	20	0	0	0	0
3	B	27	0	0	0	0
All	All	301	0	150	0	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 0.

There are no clashes within the asymmetric unit.

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

There are no protein molecules in this entry.

### 5.3.2 Protein sidechains [i](#)

There are no protein molecules in this entry.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	SPM	A	13	-	13,13,13	1.69	3 (23%)	12,12,12	1.41	2 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SPM	A	13	-	-	2/11/11/11	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	13	SPM	C9-N10	3.76	1.59	1.46
2	A	13	SPM	C11-N10	2.28	1.54	1.46
2	A	13	SPM	C6-N5	2.15	1.54	1.46

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	13	SPM	C8-C9-N10	-2.21	106.19	112.14
2	A	13	SPM	C11-N10-C9	-2.10	103.53	113.45

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	13	SPM	N10-C11-C12-C13
2	A	13	SPM	C11-C12-C13-N14

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

### 6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

### 6.4 Ligands

EDS was not executed - this section is therefore empty.

### 6.5 Other polymers

EDS was not executed - this section is therefore empty.