

wwPDB X-ray Structure Validation Summary Report (i)

May 25, 2020 – 05:55 pm BST

PDB ID : 5D56

Title: In meso in situ serial X-ray crystallography structure of diacylglycerol kinase,

DgkA, at 100 K

Authors: Huang, C.-Y.; Howe, N.; Olieric, V.; Warshamanage, R.; Diederichs, K.; Wang,

M.; Caffrey, M.

Deposited on : 2015-08-10

Resolution : 2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at $\frac{\text{https://www.wwpdb.org/validation/2017/XrayValidationReportHelp}}{\text{with specific help available everywhere you see the (i) symbol.}$

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4.02b-467

Mogul: 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.11

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac: 5.8.0158

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.11

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \textbf{Whole archive} \\ (\#\text{Entries}) \end{array}$	$\begin{array}{c} {\rm Similar \; resolution} \\ (\#{\rm Entries, \; resolution \; range(\AA)}) \end{array}$
R_{free}	130704	3140 (2.80-2.80)
Clashscore	141614	3569 (2.80-2.80)
Ramachandran outliers	138981	3498 (2.80-2.80)
Sidechain outliers	138945	3500 (2.80-2.80)
RSRZ outliers	127900	3078 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain					
1	A	130	76%			13%	11%	
1	В	130	63%		12%	25%)	
1	С	130	65%			31%		
1	D	130	72%			10% •	17%	
1	Е	130	56%	11%	•	32%		
1	F	130	^{2%} 55%	9% •		35%		



The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	78M	A	202	-	-	-	X
2	78M	A	204	-	-	-	X
2	78M	В	201	-	-	-	X
2	78M	В	203	_	-	-	X



2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 4717 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Diacylglycerol kinase.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
1	A	116	Total	С	N	О	S	0	0	0
1	A	110	887	580	149	155	3	0	U	U
1	В	97	Total	С	N	О	S	0	1	0
1	Ъ	91	747	489	121	134	3	0	1	U
1	С	90	Total	С	N	О	S	0	0	0
1		90	681	443	112	123	3			
1	D	108	Total	С	N	О	S	0	0	0
1	ש	100	824	538	137	146	3	0	U	
1	Е	89	Total	С	N	О	S	0	0	0
1	L	09	670	437	108	122	3	0	U	U
1	F	85	Total	С	N	О	S	0	0	0
	Г	00	635	415	101	116	3	U	U	0

There are 96 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-8	GLY	-	expression tag	UNP P0ABN1
A	-7	HIS	=	expression tag	UNP P0ABN1
A	-6	HIS	=	expression tag	UNP P0ABN1
A	-5	HIS	_	expression tag	UNP P0ABN1
A	-4	HIS	=	expression tag	UNP P0ABN1
A	-3	HIS	_	expression tag	UNP P0ABN1
A	-2	HIS	=	expression tag	UNP P0ABN1
A	-1	GLU	=	expression tag	UNP P0ABN1
A	0	LEU	=	expression tag	UNP P0ABN1
A	41	CYS	ALA	engineered mutation	UNP P0ABN1
A	46	ALA	CYS	engineered mutation	UNP P0ABN1
A	53	VAL	ILE	engineered mutation	UNP P0ABN1
A	70	LEU	ILE	engineered mutation	UNP P0ABN1
A	96	LEU	MET	engineered mutation	UNP P0ABN1
A	107	ASP	VAL	engineered mutation	UNP P0ABN1
A	113	ALA	CYS	engineered mutation	UNP P0ABN1
В	-8	GLY	-	expression tag	UNP P0ABN1



Continued from previous page...

Chain	Residue	$oxed{f Modelled}$	Actual	Comment	Reference
В	-7	HIS	-	expression tag	UNP P0ABN1
В	-6	HIS	-	expression tag	UNP P0ABN1
В	-5	HIS	-	expression tag	UNP P0ABN1
В	-4	HIS	-	expression tag	UNP P0ABN1
В	-3	HIS	-	expression tag	UNP P0ABN1
В	-2	HIS	-	expression tag	UNP P0ABN1
В	-1	GLU	-	expression tag	UNP P0ABN1
В	0	LEU	-	expression tag	UNP P0ABN1
В	41	CYS	ALA	engineered mutation	UNP P0ABN1
В	46	ALA	CYS	engineered mutation	UNP P0ABN1
В	53	VAL	ILE	engineered mutation	UNP P0ABN1
В	70	LEU	ILE	engineered mutation	UNP P0ABN1
В	96	LEU	MET	engineered mutation	UNP P0ABN1
В	107	ASP	VAL	engineered mutation	UNP P0ABN1
В	113	ALA	CYS	engineered mutation	UNP P0ABN1
С	-8	GLY	-	expression tag	UNP P0ABN1
С	-7	HIS	_	expression tag	UNP P0ABN1
С	-6	HIS	-	expression tag	UNP P0ABN1
С	-5	HIS	-	expression tag	UNP P0ABN1
С	-4	HIS	_	expression tag	UNP P0ABN1
С	-3	HIS	-	expression tag	UNP P0ABN1
С	-2	HIS	-	expression tag	UNP P0ABN1
С	-1	GLU	-	expression tag	UNP P0ABN1
С	0	LEU	-	expression tag	UNP P0ABN1
С	41	CYS	ALA	engineered mutation	UNP P0ABN1
С	46	ALA	CYS	engineered mutation	UNP P0ABN1
С	53	VAL	ILE	engineered mutation	UNP P0ABN1
С	70	LEU	ILE	engineered mutation	UNP P0ABN1
С	96	LEU	MET	engineered mutation	UNP P0ABN1
С	107	ASP	VAL	engineered mutation	UNP P0ABN1
С	113	ALA	CYS	engineered mutation	UNP P0ABN1
D	-8	GLY	-	expression tag	UNP P0ABN1
D	-7	HIS	-	expression tag	UNP P0ABN1
D	-6	HIS	-	expression tag	UNP P0ABN1
D	-5	HIS	-	expression tag	UNP P0ABN1
D	-4	HIS	-	expression tag	UNP P0ABN1
D	-3	HIS	-	expression tag	UNP P0ABN1
D	-2	HIS	-	expression tag	UNP P0ABN1
D	-1	GLU	-	expression tag	UNP P0ABN1
D	0	LEU	-	expression tag	UNP P0ABN1
D	41	CYS	ALA	engineered mutation	UNP P0ABN1
D	46	ALA	CYS	engineered mutation	UNP P0ABN1




 $Continued\ from\ previous\ page...$

Chain	Residue	Modelled	Actual	Comment	Reference
D	53	VAL	ILE	engineered mutation	UNP P0ABN1
D	70	LEU	ILE	engineered mutation	UNP P0ABN1
D	96	LEU	MET	engineered mutation	UNP P0ABN1
D	107	ASP	VAL	engineered mutation	UNP P0ABN1
D	113	ALA	CYS	engineered mutation	UNP P0ABN1
Е	-8	GLY	-	expression tag	UNP P0ABN1
Е	-7	HIS	-	expression tag	UNP P0ABN1
Е	-6	HIS	-	expression tag	UNP P0ABN1
Е	-5	HIS	-	expression tag	UNP P0ABN1
Е	-4	HIS	-	expression tag	UNP P0ABN1
Е	-3	HIS	_	expression tag	UNP P0ABN1
Е	-2	HIS	-	expression tag	UNP P0ABN1
Е	-1	GLU	_	expression tag	UNP P0ABN1
Е	0	LEU	_	expression tag	UNP P0ABN1
Е	41	CYS	ALA	engineered mutation	UNP P0ABN1
Е	46	ALA	CYS	engineered mutation	UNP P0ABN1
Е	53	VAL	ILE	engineered mutation	UNP P0ABN1
Е	70	LEU	ILE	engineered mutation	UNP P0ABN1
Е	96	LEU	MET	engineered mutation	UNP P0ABN1
Е	107	ASP	VAL	engineered mutation	UNP P0ABN1
Е	113	ALA	CYS	engineered mutation	UNP P0ABN1
F	-8	GLY	_	expression tag	UNP P0ABN1
F	-7	HIS	_	expression tag	UNP P0ABN1
F	-6	HIS	-	expression tag	UNP P0ABN1
F	-5	HIS	_	expression tag	UNP P0ABN1
F	-4	HIS	_	expression tag	UNP P0ABN1
F	-3	HIS	_	expression tag	UNP P0ABN1
F	-2	HIS	_	expression tag	UNP P0ABN1
F	-1	GLU	_	expression tag	UNP P0ABN1
F	0	LEU	_	expression tag	UNP P0ABN1
F	41	CYS	ALA	engineered mutation	UNP P0ABN1
F	46	ALA	CYS	engineered mutation	UNP P0ABN1
F	53	VAL	ILE	engineered mutation	UNP P0ABN1
F	70	LEU	ILE	engineered mutation	UNP P0ABN1
F	96	LEU	MET	engineered mutation	UNP P0ABN1
F	107	ASP	VAL	engineered mutation	UNP P0ABN1
F	113	ALA	CYS	engineered mutation	UNP P0ABN1

• Molecule 2 is (2S)-2,3-DIHYDROXYPROPYL(7Z)-PENTADEC-7-ENOATE (three-letter code: 78M) (formula: $C_{18}H_{34}O_4$).






Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 22 18 4	0	0
2	A	1	Total C O 22 18 4	0	0
2	A	1	Total C O 22 18 4	0	0
2	A	1	Total C O 22 18 4	0	0
2	В	1	Total C O 22 18 4	0	0
2	В	1	Total C O 22 18 4	0	0
2	В	1	Total C O 22 18 4	0	0
2	С	1	Total C O 22 18 4	0	0
2	D	1	Total C O 22 18 4	0	0
2	D	1	Total C O 22 18 4	0	0
2	D	1	Total C O 22 18 4	0	0

 \bullet Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	D	1	Total Zn 1 1	0	0




• Molecule 4 is CITRATE ANION (three-letter code: FLC) (formula: C₆H₅O₇).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf	
4	D	1	Total 13	C 6	O 7	0	0

 \bullet Molecule 5 is ACETATE ION (three-letter code: ACT) (formula: $\mathrm{C_2H_3O_2}).$



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	D	1	Total C O 4 2 2	0	0

• Molecule 6 is water.



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	3	Total O 3 3	0	0
6	В	1	Total O 1 1	0	0
6	D	5	Total O 5 5	0	0
6	E	4	Total O 4 4	0	0




3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Diacylglycerol kinase Chain A: 11% • Molecule 1: Diacylglycerol kinase Chain B: • Molecule 1: Diacylglycerol kinase Chain C: 65% 31% HISSHIPS
HIS • Molecule 1: Diacylglycerol kinase Chain D: 72% 10% 17% • Molecule 1: Diacylglycerol kinase Chain E: 32%



5D56





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	74.80Å 93.04Å 143.05Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.52 - 2.80	Depositor
resolution (11)	46.52 - 2.77	EDS
% Data completeness	97.0 (46.52-2.80)	Depositor
(in resolution range)	97.0 (46.52-2.77)	EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.74 (at 2.77Å)	Xtriage
Refinement program	PHENIX 1.9_1692	Depositor
R, R_{free}	0.243 , 0.283	Depositor
10, 10 free	0.244 , 0.284	DCC
R_{free} test set	1260 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	87.5	Xtriage
Anisotropy	0.071	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.29, 67.3	EDS
L-test for twinning ²	$ < L > = 0.47, < L^2> = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4717	wwPDB-VP
Average B, all atoms $(Å^2)$	94.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 9.70% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $< L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: 78M, ZN, FLC, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	Bond lengths		angles
MIOI	Chain	RMSZ	# Z >5	RMSZ	# Z > 5
1	A	0.22	0/903	0.37	0/1229
1	В	0.23	0/762	0.36	0/1042
1	С	0.22	0/690	0.36	0/941
1	D	0.22	0/839	0.36	0/1144
1	E	0.22	0/679	0.39	0/927
1	F	0.28	0/642	0.43	0/879
All	All	0.23	0/4515	0.38	0/6162

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	887	0	922	17	0
1	В	747	0	776	11	0
1	С	681	0	716	10	0
1	D	824	0	850	11	0
1	Ε	670	0	703	12	0
1	F	635	0	676	9	0
2	A	88	0	136	15	0



~ · · · · · · · · · · · · · · · · · · ·	e		
Continued	trom	nremous	naae
-	110116	picolous	puyc

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	В	66	0	102	6	0
2	С	22	0	34	0	0
2	D	66	0	102	3	0
3	D	1	0	0	0	0
4	D	13	0	5	0	0
5	D	4	0	3	0	0
6	A	3	0	0	0	0
6	В	1	0	0	0	0
6	D	5	0	0	1	0
6	Ε	4	0	0	0	0
All	All	4717	0	5025	66	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 66 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$egin{aligned} ext{Interatomic} \ ext{distance} \ (ext{Å}) \end{aligned}$	$egin{aligned} ext{Clash} \ ext{overlap } (ext{Å}) \end{aligned}$
1:F:54:THR:HA	1:F:57:LEU:HB2	1.71	0.72
1:A:88:GLU:OE2	1:A:88:GLU:N	2.26	0.68
1:A:54:THR:HG23	1:A:115:LEU:HD13	1.79	0.62
1:F:54:THR:HB	1:F:115:LEU:HD13	1.81	0.62
1:F:34:GLU:N	1:F:34:GLU:OE1	2.33	0.62

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	$_{ m ntiles}$
1	A	114/130 (88%)	114 (100%)	0	0	100	100
1	В	96/130 (74%)	96 (100%)	0	0	100	100



Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	С	88/130 (68%)	88 (100%)	0	0	100 100
1	D	$106/130 \; (82\%)$	106 (100%)	0	0	100 100
1	E	87/130 (67%)	85 (98%)	1 (1%)	1 (1%)	14 41
1	F	83/130 (64%)	83 (100%)	0	0	100 100
All	All	574/780 (74%)	572 (100%)	1 (0%)	1 (0%)	47 78

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	E	50	VAL

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Perce	${ m ntiles}$
1	A	90/102 (88%)	89 (99%)	1 (1%)	73	92
1	В	79/102 (78%)	79 (100%)	0	100	100
1	С	73/102 (72%)	72 (99%)	1 (1%)	67	90
1	D	84/102 (82%)	83 (99%)	1 (1%)	71	92
1	E	72/102 (71%)	71 (99%)	1 (1%)	67	90
1	F	69/102 (68%)	67 (97%)	2 (3%)	42	76
All	All	$467/612 \ (76\%)$	461 (99%)	6 (1%)	69	91

5 of 6 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	22	ARG
1	F	81	ARG
1	Ε	48	LEU
1	С	32	ARG
1	F	48	LEU



Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	${f Res}$	Type
1	D	87	HIS
1	D	119	HIS

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

5.6 Ligand geometry (i)

Of 14 ligands modelled in this entry, 1 is monoatomic - leaving 13 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Tune	Chain	Res	Link	Во	Bond lengths			Bond angles		
MIOI	Type	Chain	nes	Link	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2	
2	78M	A	204	-	21,21,21	0.88	1 (4%)	22,22,22	0.81	1 (4%)	
2	78M	В	202	-	21,21,21	0.91	1 (4%)	22,22,22	0.80	1 (4%)	
2	78M	В	203	-	21,21,21	0.90	1 (4%)	22,22,22	0.76	1 (4%)	
2	78M	С	201	-	21,21,21	0.87	1 (4%)	22,22,22	0.79	1 (4%)	
2	78M	A	202	-	21,21,21	0.88	1 (4%)	22,22,22	0.85	1 (4%)	
5	ACT	D	203	_	1,3,3	1.27	0	0,3,3	0.00	-	
2	78M	D	206	-	21,21,21	0.89	1 (4%)	22,22,22	0.78	1 (4%)	
4	FLC	D	202	-	3,12,12	1.29	0	3,17,17	2.36	1 (33%)	



Mol	Type	Chain	Res	es Link	Во	Bond lengths			Bond angles		
WIGI			res		Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2	
2	78M	D	204	-	21,21,21	0.87	1 (4%)	22,22,22	0.81	1 (4%)	
2	78M	В	201	-	21,21,21	0.90	1 (4%)	22,22,22	0.84	1 (4%)	
2	78M	D	205	-	21,21,21	0.87	1 (4%)	22,22,22	0.79	1 (4%)	
2	78M	A	203	-	21,21,21	0.86	1 (4%)	22,22,22	0.82	1 (4%)	
2	78M	A	201	-	21,21,21	0.86	1 (4%)	22,22,22	0.79	1 (4%)	

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	78M	A	204	_	-	11/21/21/21	-
2	78M	В	202	-	-	10/21/21/21	-
2	78M	В	203	-	-	10/21/21/21	-
2	78M	С	201	-	-	10/21/21/21	-
2	78M	A	202	-	-	7/21/21/21	-
2	78M	D	206	-	-	12/21/21/21	-
4	FLC	D	202	-	-	6/6/16/16	-
2	78M	D	204	_	-	6/21/21/21	-
2	78M	В	201	-	-	10/21/21/21	-
2	78M	D	205	-	-	7/21/21/21	-
2	78M	A	203	_	-	6/21/21/21	-
2	78M	A	201	_	-	9/21/21/21	-

The worst 5 of 11 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(\mathbf{\mathring{A}})$	$\operatorname{Ideal}(\operatorname{\AA})$
2	В	203	78M	O2-C1	2.85	1.41	1.33
2	В	202	78M	O2-C1	2.81	1.41	1.33
2	A	202	78M	O2-C1	2.78	1.41	1.33
2	В	201	78M	O2-C1	2.74	1.41	1.33
2	A	204	78M	O2-C1	2.74	1.41	1.33

The worst 5 of 12 bond angle outliers are listed below:

\mathbf{Mol}	Chain	${ m Res}$	Type	${f Atoms}$	Z	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^o)$
4	D	202	FLC	CB-CA-CAC	3.93	121.28	114.98



Continued from previous page...

Mol	Chain	Res	Type	${f Atoms}$	\mathbf{Z}	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^o)$
2	A	202	78M	O2-C1-C2	2.98	121.25	111.91
2	В	201	78M	O2-C1-C2	2.87	120.90	111.91
2	D	204	78M	O2-C1-C2	2.84	120.83	111.91
2	В	202	78M	O2-C1-C2	2.77	120.59	111.91

There are no chirality outliers.

5 of 104 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	С	201	78M	O2-C17-C18-C20
2	С	201	78M	O2-C17-C18-O19
2	D	206	78M	C17-C18-C20-O21
2	D	206	78M	O2-C17-C18-C20
2	D	206	78M	O2-C17-C18-O19


There are no ring outliers.

9 monomers are involved in 23 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	204	78M	7	0
2	В	202	78M	2	0
2	В	203	78M	1	0
2	A	202	78M	6	0
2	D	206	78M	2	0
2	В	201	78M	3	0
2	D	205	78M	1	0
2	A	203	78M	3	0
2	A	201	78M	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.














5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	$\#\mathrm{RSRZ}{>}2$	$\mathbf{OWAB}(\mathrm{\AA}^2)$	Q < 0.9
1	A	$116/130 \ (89\%)$	-0.14	3 (2%) 56 46	61, 81, 134, 159	0
1	В	97/130 (74%)	-0.11	4 (4%) 37 27	58, 77, 131, 142	0
1	С	90/130 (69%)	-0.17	3 (3%) 46 36	61, 81, 124, 144	0
1	D	108/130 (83%)	-0.29	0 100 100	65, 83, 117, 140	0
1	E	89/130 (68%)	0.13	10 (11%) 5 3	66, 98, 144, 175	3 (3%)
1	F	85/130 (65%)	0.04	3 (3%) 44 34	70, 97, 150, 169	0
All	All	585/780 (75%)	-0.10	23 (3%) 39 29	58, 85, 138, 175	3 (0%)

The worst 5 of 23 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	33	GLN	8.4
1	С	86	TYR	5.1
1	F	36	VAL	5.0
1	E	48	LEU	4.9
1	В	27	ASN	4.6

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no carbohydrates in this entry.



6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\operatorname{\textbf{B-factors}}(\mathring{\mathbf{A}}^2)$	Q < 0.9
4	FLC	D	202	13/13	0.58	0.25	161,180,191,191	0
2	78M	В	201	22/22	0.62	0.42	$120,\!135,\!145,\!156$	0
2	78M	С	201	22/22	0.69	0.29	64,101,134,144	0
2	78M	A	202	22/22	0.71	0.59	$75,\!110,\!129,\!132$	0
2	78M	A	204	22/22	0.75	0.73	76,90,107,113	22
2	78M	D	205	22/22	0.75	0.37	107,131,144,155	0
2	78M	В	203	22/22	0.79	0.41	73,97,127,135	22
2	78M	D	206	22/22	0.80	0.35	74,98,117,122	0
2	78M	В	202	22/22	0.81	0.49	81,113,126,128	0
2	78M	D	204	22/22	0.82	0.34	102,121,127,128	0
2	78M	A	203	22/22	0.82	0.41	81,99,129,142	0
2	78M	A	201	22/22	0.91	0.49	66,93,115,119	0
5	ACT	D	203	4/4	0.94	0.23	113,133,135,138	0
3	ZN	D	201	1/1	0.94	0.04	$120,\!120,\!120,\!120$	1


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around 78M B 201:

 $2 \mathrm{mF}_o\text{-DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around 78M C 201:

 $2 \mathrm{mF}_o\text{-DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)












Electron density around 78M A 202:

 $2 \mathrm{mF}_o\text{-DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around 78M A 204:

 $2 \text{mF}_o\text{-DF}_c$ (at 0.7 rmsd) in gray $\text{mF}_o\text{-DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)












Electron density around 78M D 205: 2mF_o-DF_c (at 0.7 rmsd) in gray mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)

Electron density around 78M B 203: 2mF_o-DF_c (at 0.7 rmsd) in gray mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)

















Electron density around 78M A 203:


 $2 {
m mF}_o {
m -DF}_c$ (at 0.7 rmsd) in gray ${
m mF}_o {
m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)











6.5 Other polymers (i)

There are no such residues in this entry.

