



wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 2, 2023 – 08:41 AM EDT

PDB ID : 1DEQ
Title : THE CRYSTAL STRUCTURE OF MODIFIED BOVINE FIBRINOGEN (AT 4 ANGSTROM RESOLUTION)
Authors : Brown, J.H.; Volkmann, N.; Jun, G.; Henschen-Edman, A.H.; Cohen, C.
Deposited on : 1999-11-15
Resolution : 3.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.34
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.34


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1659 (3.60-3.40)
Clashscore	141614	1036 (3.58-3.42)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	390	
1	D	390	
1	N	390	
1	Q	390	
2	B	408	
2	E	408	
2	O	408	
2	R	408	
3	C	411	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	F	411	 89% 10%
3	P	411	 90% 10%
3	S	411	 90% 10%
4	M	90	 87% 13%
4	Z	90	 87% 13%

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 3900 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called FIBRINOGEN (ALPHA CHAIN).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
1	A	180	Total C 180 180	0	0	180
1	D	180	Total C 180 180	0	0	180
1	N	180	Total C 180 180	0	0	180
1	Q	180	Total C 180 180	0	0	180

- Molecule 2 is a protein called FIBRINOGEN (BETA CHAIN).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
2	B	380	Total C 380 380	0	0	380
2	E	380	Total C 380 380	0	0	380
2	O	380	Total C 380 380	0	0	380
2	R	380	Total C 380 380	0	0	380

- Molecule 3 is a protein called FIBRINOGEN (GAMMA CHAIN).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
3	C	370	Total C 370 370	0	0	370
3	F	370	Total C 370 370	0	0	370
3	P	370	Total C 370 370	0	0	370
3	S	370	Total C 370 370	0	0	370

- Molecule 4 is a protein called FIBRINOGEN.



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
4	M	90	Total C 90 90	0	0	90
4	Z	90	Total C 90 90	0	0	90





● Molecule 3: FIBRINOGEN (GAMMA CHAIN)




● Molecule 3: FIBRINOGEN (GAMMA CHAIN)





● Molecule 3: FIBRINOGEN (GAMMA CHAIN)



● Molecule 4: FIBRINOGEN



● Molecule 4: FIBRINOGEN



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	176.01Å 94.94Å 209.81Å 90.00° 94.41° 90.00°	Depositor
Resolution (Å)	10.00 – 3.50 209.18 – 3.34	Depositor EDS
% Data completeness (in resolution range)	86.9 (10.00-3.50) 78.8 (209.18-3.34)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.89 (at 3.33Å)	Xtrriage
Refinement program	X-PLOR 3.851	Depositor
R, R_{free}	0.257 , 0.370 0.479 , 0.485	Depositor DCC
R_{free} test set	3953 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å ²)	91.6	Xtrriage
Anisotropy	0.887	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 152.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.29$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.72	EDS
Total number of atoms	3900	wwPDB-VP
Average B, all atoms (Å ²)	225.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 10.82% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

There are no protein, RNA or DNA chains available to summarize Z scores of covalent bonds and angles.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	180	0	0	3	0
1	D	180	0	0	5	0
1	N	180	0	0	1	0
1	Q	180	0	0	4	0
2	B	380	0	0	1	0
2	E	380	0	0	1	0
2	O	380	0	0	0	0
2	R	380	0	0	1	0
3	C	370	0	0	0	0
3	F	370	0	0	3	0
3	P	370	0	0	0	0
3	S	370	0	0	1	0
4	M	90	0	0	8	0
4	Z	90	0	0	6	0
All	All	3900	0	0	32	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 32 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:E:142:ASN:CA	2:E:143:GLU:CA	2.09	1.29
4:M:134:UNK:CA	4:M:135:UNK:CA	2.16	1.22
4:M:43:UNK:CA	4:M:44:UNK:CA	2.18	1.21
1:Q:82:ASN:CA	1:Q:83:SER:CA	2.22	1.17
4:M:124:UNK:CA	4:M:125:UNK:CA	2.27	1.12

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

There are no protein backbone outliers to report in this entry.

5.3.2 Protein sidechains [i](#)

There are no protein residues with a non-rotameric sidechain to report in this entry.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.3 Carbohydrates [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.4 Ligands [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.5 Other polymers [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.