



wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 15, 2023 – 11:23 AM EDT

PDB ID : 8DH1
Title : T7 RNA polymerase elongation complex with unnatural base dDs-PaTP pair
Authors : Oh, J.; Wang, D.
Deposited on : 2022-06-24
Resolution : 2.65 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<https://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1332 (2.68-2.64)
Clashscore	141614	1374 (2.68-2.64)
Ramachandran outliers	138981	1349 (2.68-2.64)
Sidechain outliers	138945	1349 (2.68-2.64)
RSRZ outliers	127900	1318 (2.68-2.64)
RNA backbone	3102	1010 (2.96-2.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	B	883	 5% 83% 14% ••
1	E	883	 10% 72% 16% • 11%
1	I	883	 10% 72% 19% • 7%
1	L	883	 14% 66% 12% • 20%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	A	18	
2	F	18	
2	J	18	
2	M	18	
3	C	12	
3	G	12	
3	K	12	
3	N	12	
4	D	9	
4	H	9	
4	O	9	

2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 27087 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called T7 RNA polymerase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	B	855	Total 6719	C 4280	N 1169	O 1235	S 35	0	0	0
1	E	785	Total 6125	C 3908	N 1060	O 1125	S 32	0	0	0
1	I	825	Total 6478	C 4133	N 1118	O 1192	S 35	0	0	0
1	L	702	Total 5358	C 3402	N 936	O 990	S 30	0	0	0

- Molecule 2 is a DNA chain called Template strand DNA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	N	O	P				S
2	A	16	Total 332	C 160	N 62	O 93	P 16	S 1	0	0	0
2	F	17	Total 351	C 170	N 67	O 97	P 16	S 1	0	0	0
2	J	12	Total 248	C 120	N 45	O 70	P 12	S 1	0	0	0
2	M	14	Total 289	C 140	N 52	O 82	P 14	S 1	0	0	0


- Molecule 3 is a RNA chain called RNA.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace	
			Total	C	N	O				P
3	C	8	Total 174	C 77	N 33	O 56	P 8	0	0	0
3	G	8	Total 174	C 77	N 33	O 56	P 8	0	0	0
3	K	8	Total 174	C 77	N 33	O 56	P 8	0	0	0
3	N	8	Total 171	C 77	N 33	O 54	P 7	0	0	0

- Molecule 4 is a DNA chain called Non-template strand DNA.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
4	D	6	Total 121	C 58	N 20	O 37	P 6	0	0	0
4	H	8	Total 160	C 77	N 25	O 50	P 8	0	0	0
4	O	5	Total 102	C 49	N 17	O 31	P 5	0	0	0

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	C O		
5	B	1	Total 6	C O 3 3	0	0
5	C	1	Total 6	C O 3 3	0	0
5	I	1	Total 6	C O 3 3	0	0

- Molecule 6 is 1-{5-O-[(S)-hydroxy{[(R)-hydroxy(phosphonoxy)phosphoryl]oxy}phosphoryl]-beta-D-ribofuranosyl}-1H-pyrrole-2-carbaldehyde (three-letter code: S8L) (formula: C₁₀H₁₆NO₁₄P₃) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
6	B	1	28	10	1	14	3	0	0

- Molecule 7 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Mg		
7	B	1	1	1	0	0
7	C	1	1	1	0	0

- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
8	B	38	38	38	0	0
8	C	3	3	3	0	0
8	E	14	14	14	0	0
8	F	1	1	1	0	0
8	I	4	4	4	0	0
8	L	2	2	2	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	O	1	Total	O	0	0
			1	1		


3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: T7 RNA polymerase




- Molecule 1: T7 RNA polymerase







• Molecule 2: Template strand DNA




• Molecule 2: Template strand DNA



• Molecule 2: Template strand DNA



- Molecule 2: Template strand DNA




- Molecule 3: RNA




- Molecule 3: RNA



- Molecule 3: RNA




- Molecule 3: RNA





- Molecule 4: Non-template strand DNA



- Molecule 4: Non-template strand DNA



- Molecule 4: Non-template strand DNA



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	78.87Å 86.16Å 201.68Å 89.85° 85.09° 69.56°	Depositor
Resolution (Å)	39.34 – 2.65 39.34 – 2.65	Depositor EDS
% Data completeness (in resolution range)	89.6 (39.34-2.65) 89.6 (39.34-2.65)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.26 (at 2.65Å)	Xtrriage
Refinement program	PHENIX 1.19	Depositor
R, R_{free}	0.230 , 0.272 0.227 , 0.269	Depositor DCC
R_{free} test set	6387 reflections (4.95%)	wwPDB-VP
Wilson B-factor (Å ²)	65.0	Xtrriage
Anisotropy	0.365	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 60.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	27087	wwPDB-VP
Average B, all atoms (Å ²)	100.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.92% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 91N, MG, S8L, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	B	0.29	0/6869	0.51	1/9291 (0.0%)
1	E	0.29	0/6262	0.52	2/8472 (0.0%)
1	I	0.30	0/6623	0.52	0/8957
1	L	0.28	0/5474	0.52	1/7411 (0.0%)
2	A	0.61	0/343	0.82	0/524
2	F	0.56	0/365	0.82	0/559
2	J	0.63	0/248	0.83	0/377
2	M	0.59	0/294	0.87	0/448
3	C	0.34	0/194	0.82	0/301
3	G	0.24	0/194	0.72	0/301
3	K	0.21	0/194	0.70	0/301
3	N	0.20	0/191	0.65	0/297
4	D	0.48	0/134	0.96	0/204
4	H	0.49	0/177	0.97	0/270
4	O	0.56	0/113	1.06	0/172
All	All	0.31	0/27675	0.56	4/37885 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	I	0	1

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	787	ASP	CB-CG-OD1	5.70	123.43	118.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	L	812	ASP	CB-CG-OD1	5.67	123.40	118.30
1	E	731	ASP	CB-CG-OD2	5.36	123.12	118.30
1	B	77	LEU	CB-CG-CD2	-5.04	102.44	111.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	I	31	ARG	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	6719	0	6670	60	0
1	E	6125	0	6043	66	0
1	I	6478	0	6406	99	1
1	L	5358	0	5131	57	0
2	A	332	0	168	4	0
2	F	351	0	180	4	0
2	J	248	0	123	3	0
2	M	289	0	146	2	0
3	C	174	0	88	0	0
3	G	174	0	88	1	0
3	K	174	0	88	1	0
3	N	171	0	89	0	0
4	D	121	0	69	2	0
4	H	160	0	92	2	0
4	O	102	0	58	2	0
5	B	6	0	8	0	0
5	C	6	0	8	0	0
5	I	6	0	8	0	0
6	B	28	0	0	1	0
7	B	1	0	0	0	0
7	C	1	0	0	0	0
8	B	38	0	0	2	0
8	C	3	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
8	E	14	0	0	0	0
8	F	1	0	0	0	0
8	I	4	0	0	0	0
8	L	2	0	0	0	0
8	O	1	0	0	0	0
All	All	27087	0	25463	293	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 293 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:B:902:S8L:C19	6:B:902:S8L:O27	1.66	1.23
1:I:155:ARG:HA	1:I:163:LYS:HG3	1.63	0.81
1:B:6:ILE:HD12	1:B:291:ARG:HH22	1.50	0.76
1:B:632:ARG:NH1	2:A:10:91N:S8	2.62	0.73
1:I:163:LYS:HA	1:I:166:VAL:HG12	1.70	0.72

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:I:31:ARG:NH2	1:I:194:GLY:O[1_655]	2.16	0.04

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	B	847/883 (96%)	830 (98%)	17 (2%)	0	100 100
1	E	773/883 (88%)	757 (98%)	16 (2%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	I	811/883 (92%)	791 (98%)	20 (2%)	0	100	100
1	L	684/883 (78%)	666 (97%)	18 (3%)	0	100	100
All	All	3115/3532 (88%)	3044 (98%)	71 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	698/729 (96%)	665 (95%)	33 (5%)	26	40
1	E	628/729 (86%)	581 (92%)	47 (8%)	13	21
1	I	673/729 (92%)	622 (92%)	51 (8%)	13	21
1	L	528/729 (72%)	483 (92%)	45 (8%)	10	15
All	All	2527/2916 (87%)	2351 (93%)	176 (7%)	15	23

5 of 176 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	I	600	GLU
1	L	257	ARG
1	I	666	MET
1	I	879	ASP
1	L	423	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 11 such sidechains are listed below:

Mol	Chain	Res	Type
1	I	523	HIS
1	L	486	HIS
1	L	649	GLN
1	L	588	ASN
1	E	869	ASN

5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
3	C	7/12 (58%)	0	0
3	G	7/12 (58%)	0	0
3	K	7/12 (58%)	0	0
3	N	7/12 (58%)	0	0
All	All	28/48 (58%)	0	0

There are no RNA backbone outliers to report.

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	S8L	B	902	7	23,29,29	5.70	8 (34%)	29,45,45	1.43	7 (24%)
5	GOL	B	901	-	5,5,5	1.07	0	5,5,5	0.98	0
5	GOL	C	101	-	5,5,5	0.94	0	5,5,5	1.02	0
5	GOL	I	901	-	5,5,5	1.02	0	5,5,5	0.92	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	S8L	B	902	7	-	7/18/40/40	0/2/2/2
5	GOL	B	901	-	-	0/4/4/4	-
5	GOL	C	101	-	-	2/4/4/4	-
5	GOL	I	901	-	-	0/4/4/4	-

The worst 5 of 8 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	B	902	S8L	O27-C19	18.48	1.66	1.41
6	B	902	S8L	C17-C19	-15.83	1.29	1.53
6	B	902	S8L	O27-C15	-7.84	1.27	1.45
6	B	902	S8L	P09-O10	5.03	1.66	1.50
6	B	902	S8L	O28-C16	-4.66	1.32	1.43

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	B	902	S8L	C16-C17-C19	3.50	106.25	100.98
6	B	902	S8L	P05-O04-P01	-3.05	122.36	132.83
6	B	902	S8L	P05-O08-P09	-2.58	123.98	132.83
6	B	902	S8L	O23-C22-C21	-2.12	121.17	124.17
6	B	902	S8L	O12-P09-O08	2.08	111.61	104.64

There are no chirality outliers.

5 of 9 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	C	101	GOL	C1-C2-C3-O3
6	B	902	S8L	C15-C14-O13-P01
6	B	902	S8L	C14-O13-P01-O04
6	B	902	S8L	O13-C14-C15-C16
5	C	101	GOL	O2-C2-C3-O3


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	B	902	S8L	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths,

bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	B	855/883 (96%)	0.37	42 (4%) 29 26	39, 68, 134, 215	0
1	E	785/883 (88%)	0.62	85 (10%) 5 4	45, 76, 169, 252	0
1	I	825/883 (93%)	0.75	85 (10%) 6 4	59, 104, 167, 218	0
1	L	702/883 (79%)	0.96	126 (17%) 1 1	69, 114, 171, 267	0
2	A	15/18 (83%)	0.95	4 (26%) 0 0	52, 71, 237, 239	0
2	F	16/18 (88%)	0.71	3 (18%) 1 1	62, 110, 248, 259	0
2	J	11/18 (61%)	0.73	1 (9%) 9 7	70, 79, 176, 202	0
2	M	13/18 (72%)	0.68	1 (7%) 13 10	90, 107, 176, 182	0
3	C	8/12 (66%)	0.48	0 100 100	52, 63, 72, 83	0
3	G	8/12 (66%)	-0.02	0 100 100	59, 77, 119, 140	0
3	K	8/12 (66%)	0.22	0 100 100	77, 94, 120, 141	0
3	N	8/12 (66%)	0.56	1 (12%) 3 2	88, 107, 161, 177	0
4	D	6/9 (66%)	3.24	5 (83%) 0 0	211, 225, 229, 241	0
4	H	8/9 (88%)	2.44	5 (62%) 0 0	182, 197, 224, 234	0
4	O	5/9 (55%)	1.39	1 (20%) 1 1	204, 210, 212, 223	0
All	All	3273/3679 (88%)	0.67	359 (10%) 5 4	39, 93, 168, 267	0

The worst 5 of 359 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	L	17	ALA	9.0
1	L	113	ALA	6.8
1	L	21	PHE	6.0
1	L	16	LEU	5.5
1	L	744	GLN	5.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	GOL	I	901	6/6	0.82	0.21	84,85,98,105	0
6	S8L	B	902	28/28	0.90	0.20	68,104,128,150	0
5	GOL	C	101	6/6	0.91	0.20	60,69,73,77	0
7	MG	B	903	1/1	0.92	0.37	97,97,97,97	0
5	GOL	B	901	6/6	0.95	0.24	51,60,74,75	0
7	MG	C	102	1/1	0.95	0.44	143,143,143,143	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.