



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 15, 2023 – 04:43 AM EDT

PDB ID : 8DHV  
Title : Treponema lecithinolyticum beta-glucuronidase  
Authors : Lietzan, A.D.; Redinbo, M.R.  
Deposited on : 2022-06-28  
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3398 (1.60-1.60)
Clashscore	141614	3665 (1.60-1.60)
Ramachandran outliers	138981	3564 (1.60-1.60)
Sidechain outliers	138945	3563 (1.60-1.60)
RSRZ outliers	127900	3321 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	631	 89% 5% 5%
1	B	631	 2% 90% 5%

## 2 Entry composition i

There are 5 unique types of molecules in this entry. The entry contains 10723 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glycosyl hydrolase family 2, TIM barrel domain protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	598	4842	3103	798	914	27	0	9	0
1	B	597	4810	3086	791	905	28	0	10	0

There are 68 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-6	MET	-	initiating methionine	UNP U2KI81
A	-5	HIS	-	expression tag	UNP U2KI81
A	-4	HIS	-	expression tag	UNP U2KI81
A	-3	HIS	-	expression tag	UNP U2KI81
A	-2	HIS	-	expression tag	UNP U2KI81
A	-1	HIS	-	expression tag	UNP U2KI81
A	0	HIS	-	expression tag	UNP U2KI81
A	1	SER	-	expression tag	UNP U2KI81
A	2	SER	-	expression tag	UNP U2KI81
A	3	GLY	-	expression tag	UNP U2KI81
A	4	VAL	-	expression tag	UNP U2KI81
A	5	ASP	-	expression tag	UNP U2KI81
A	6	LEU	-	expression tag	UNP U2KI81
A	7	GLY	-	expression tag	UNP U2KI81
A	8	THR	-	expression tag	UNP U2KI81
A	9	GLU	-	expression tag	UNP U2KI81
A	10	ASN	-	expression tag	UNP U2KI81
A	11	LEU	-	expression tag	UNP U2KI81
A	12	TYR	-	expression tag	UNP U2KI81
A	13	PHE	-	expression tag	UNP U2KI81
A	14	GLN	-	expression tag	UNP U2KI81
A	15	SER	-	expression tag	UNP U2KI81
A	16	ASN	-	expression tag	UNP U2KI81
A	17	ALA	-	expression tag	UNP U2KI81
A	18	MET	-	expression tag	UNP U2KI81

*Continued on next page...*

*Continued from previous page...*


Chain	Residue	Modelled	Actual	Comment	Reference
A	19	LEU	-	expression tag	UNP U2KI81
A	20	TYR	-	expression tag	UNP U2KI81
A	21	PRO	-	expression tag	UNP U2KI81
A	22	ILE	-	expression tag	UNP U2KI81
A	23	LEU	-	expression tag	UNP U2KI81
A	24	THR	-	expression tag	UNP U2KI81
A	25	GLN	-	expression tag	UNP U2KI81
A	26	SER	-	expression tag	UNP U2KI81
A	27	ARG	-	expression tag	UNP U2KI81
B	-6	MET	-	initiating methionine	UNP U2KI81
B	-5	HIS	-	expression tag	UNP U2KI81
B	-4	HIS	-	expression tag	UNP U2KI81
B	-3	HIS	-	expression tag	UNP U2KI81
B	-2	HIS	-	expression tag	UNP U2KI81
B	-1	HIS	-	expression tag	UNP U2KI81
B	0	HIS	-	expression tag	UNP U2KI81
B	1	SER	-	expression tag	UNP U2KI81
B	2	SER	-	expression tag	UNP U2KI81
B	3	GLY	-	expression tag	UNP U2KI81
B	4	VAL	-	expression tag	UNP U2KI81
B	5	ASP	-	expression tag	UNP U2KI81
B	6	LEU	-	expression tag	UNP U2KI81
B	7	GLY	-	expression tag	UNP U2KI81
B	8	THR	-	expression tag	UNP U2KI81
B	9	GLU	-	expression tag	UNP U2KI81
B	10	ASN	-	expression tag	UNP U2KI81
B	11	LEU	-	expression tag	UNP U2KI81
B	12	TYR	-	expression tag	UNP U2KI81
B	13	PHE	-	expression tag	UNP U2KI81
B	14	GLN	-	expression tag	UNP U2KI81
B	15	SER	-	expression tag	UNP U2KI81
B	16	ASN	-	expression tag	UNP U2KI81
B	17	ALA	-	expression tag	UNP U2KI81
B	18	MET	-	expression tag	UNP U2KI81
B	19	LEU	-	expression tag	UNP U2KI81
B	20	TYR	-	expression tag	UNP U2KI81
B	21	PRO	-	expression tag	UNP U2KI81
B	22	ILE	-	expression tag	UNP U2KI81
B	23	LEU	-	expression tag	UNP U2KI81
B	24	THR	-	expression tag	UNP U2KI81
B	25	GLN	-	expression tag	UNP U2KI81
B	26	SER	-	expression tag	UNP U2KI81

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	27	ARG	-	expression tag	UNP U2KI81

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
2	A	1	5	4	1	0	0
2	A	1	5	4	1	0	0
2	A	1	5	4	1	0	0
2	A	1	5	4	1	0	0
2	A	1	5	4	1	0	0
2	B	1	5	4	1	0	0
2	B	1	5	4	1	0	0
2	B	1	5	4	1	0	0
2	B	1	5	4	1	0	0

- Molecule 3 is (4S)-2-METHYL-2,4-PENTANEDIOL (three-letter code: MPD) (formula: C<sub>6</sub>H<sub>14</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 8 6 2	0	0
3	A	1	Total C O 8 6 2	0	0
3	A	1	Total C O 8 6 2	0	0
3	B	1	Total C O 8 6 2	0	0
3	B	1	Total C O 8 6 2	0	0

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	2	Total Cl 2 2	0	0
4	B	2	Total Cl 2 2	0	0


- Molecule 5 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	529	Total O 529 529	0	0
5	B	453	Total O 453 453	0	0

### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Glycosyl hydrolase family 2, TIM barrel domain protein

Chain A: 



- Molecule 1: Glycosyl hydrolase family 2, TIM barrel domain protein

Chain B: 



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	94.30Å 94.30Å 287.95Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.15 – 1.60 49.15 – 1.60	Depositor EDS
% Data completeness (in resolution range)	99.5 (49.15-1.60) 99.5 (49.15-1.60)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.35 (at 1.60Å)	Xtrriage
Refinement program	PHENIX 1.19.2_4158	Depositor
R, $R_{free}$	0.167 , 0.196 0.166 , 0.196	Depositor DCC
$R_{free}$ test set	1997 reflections (1.17%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	23.7	Xtrriage
Anisotropy	0.524	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 45.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.98	EDS
Total number of atoms	10723	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	29.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.56% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MPD, CL, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.37	0/4971	0.62	0/6743
1	B	0.36	0/4942	0.61	0/6708
All	All	0.37	0/9913	0.62	0/13451

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
All	All	0	2

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	317	TYR	Peptide
1	B	317	TYR	Peptide

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4842	0	4640	23	0
1	B	4810	0	4572	22	0
2	A	25	0	0	1	0
2	B	20	0	0	0	0
3	A	24	0	42	0	0
3	B	16	0	28	0	0
4	A	2	0	0	0	0
4	B	2	0	0	0	0
5	A	529	0	0	2	1
5	B	453	0	0	1	0
All	All	10723	0	9282	39	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (39) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:271:GLU:OE1	5:A:801:HOH:O	1.97	0.83
1:A:16:ASN:N	5:A:803:HOH:O	2.29	0.65
1:A:95:LEU:HD11	1:B:28[B]:MET:HE1	1.79	0.65
1:A:40:LEU:HD21	1:A:76[A]:LEU:HD13	1.78	0.64
1:B:242:VAL:HG11	1:B:264:ILE:HD11	1.85	0.58
1:A:30:ILE:HD11	1:B:28[B]:MET:HE2	1.86	0.57
1:B:51:TYR:HD1	1:B:117:GLY:HA2	1.70	0.57
1:A:94:TYR:HE2	1:B:28[A]:MET:HE1	1.71	0.56
1:B:169:LEU:HG	1:B:587:ILE:HD11	1.89	0.53
1:B:133:GLU:OE1	5:B:801:HOH:O	2.19	0.53
1:B:317:TYR:CE2	1:B:342:MET:HG2	2.46	0.51
1:B:40:LEU:HD21	1:B:76[B]:LEU:HD13	1.91	0.51
1:A:94:TYR:OH	1:B:28[A]:MET:HE2	2.11	0.51
1:A:169:LEU:HD22	1:A:587:ILE:HD11	1.91	0.51
1:A:477:ASN:ND2	1:A:480:GLU:OE2	2.30	0.50
1:A:253:THR:HG23	1:A:256[B]:THR:HG21	1.95	0.48
1:B:339[A]:ILE:HD12	1:B:370:ILE:HG13	1.95	0.48
1:A:238:THR:O	1:A:256[B]:THR:HG23	2.13	0.48
1:A:60:LEU:HD11	1:B:367:GLU:HB3	1.97	0.46
1:A:353:SER:HA	1:A:354:HIS:HA	1.81	0.45
1:B:229:LYS:HG3	1:B:258:THR:HG22	1.98	0.45
1:B:317:TYR:HE2	1:B:342:MET:HG2	1.81	0.45
1:A:271:GLU:HB2	1:A:274:LYS:HB2	1.98	0.44
1:A:498:TRP:CZ2	1:A:534:ASP:HB2	2.52	0.44

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:498:TRP:CZ2	1:B:534:ASP:HB2	2.53	0.44
1:A:108:THR:HA	1:A:109:HIS:HA	1.76	0.43
1:B:108:THR:HA	1:B:109:HIS:HA	1.81	0.43
1:B:137:ASN:ND2	1:B:137:ASN:H	2.16	0.43
1:B:534:ASP:O	1:B:547:PHE:HA	2.20	0.42
1:A:253:THR:HG23	1:A:256[B]:THR:CG2	2.50	0.41
1:B:102:LEU:HG	1:B:134:ILE:HD11	2.01	0.41
1:A:158:LEU:HD22	1:A:378:ALA:HA	2.02	0.41
1:A:168:VAL:HG11	1:A:385:PHE:HD2	1.84	0.41
1:A:338:ASP:O	1:A:342:MET:HG3	2.19	0.41
1:A:28:MET:HE1	1:B:95:LEU:HD21	2.03	0.41
1:A:419:LYS:HG2	1:A:461:LEU:HD11	2.03	0.40
1:A:124:LYS:NZ	2:A:702:SO4:O2	2.54	0.40
1:B:588:ARG:HD3	1:B:588:ARG:HA	1.88	0.40
1:B:95:LEU:HD23	1:B:95:LEU:HA	1.90	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:824:HOH:O	5:A:827:HOH:O[7_555]	2.14	0.06

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	603/631 (96%)	580 (96%)	23 (4%)	0	100	100
1	B	603/631 (96%)	581 (96%)	22 (4%)	0	100	100
All	All	1206/1262 (96%)	1161 (96%)	45 (4%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	514/550 (94%)	509 (99%)	5 (1%)	76	61
1	B	503/550 (92%)	498 (99%)	5 (1%)	76	61
All	All	1017/1100 (92%)	1007 (99%)	10 (1%)	78	61

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	94	TYR
1	A	296	ARG
1	A	317	TYR
1	A	353	SER
1	A	531	TYR
1	B	76[A]	LEU
1	B	76[B]	LEU
1	B	296	ARG
1	B	317	TYR
1	B	531	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	182	GLN
1	B	137	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 18 ligands modelled in this entry, 4 are monoatomic - leaving 14 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	SO4	B	706	-	4,4,4	0.16	0	6,6,6	0.14	0
2	SO4	B	703	-	4,4,4	0.14	0	6,6,6	0.13	0
3	MPD	A	708	-	7,7,7	0.25	0	9,10,10	0.27	0
3	MPD	B	701	-	7,7,7	0.36	0	9,10,10	0.33	0
2	SO4	A	704	-	4,4,4	0.18	0	6,6,6	0.22	0
3	MPD	A	705	-	7,7,7	0.31	0	9,10,10	0.44	0
2	SO4	B	704	-	4,4,4	0.17	0	6,6,6	0.19	0
2	SO4	B	705	-	4,4,4	0.11	0	6,6,6	0.06	0
2	SO4	A	702	-	4,4,4	0.13	0	6,6,6	0.08	0
2	SO4	A	706	-	4,4,4	0.14	0	6,6,6	0.06	0
2	SO4	A	703	-	4,4,4	0.13	0	6,6,6	0.06	0
2	SO4	A	701	-	4,4,4	0.12	0	6,6,6	0.08	0
3	MPD	A	707	-	7,7,7	0.27	0	9,10,10	0.50	0
3	MPD	B	702	-	7,7,7	0.26	0	9,10,10	0.26	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	MPD	A	708	-	-	1/5/5/5	-
3	MPD	B	701	-	-	2/5/5/5	-
3	MPD	A	705	-	-	2/5/5/5	-
3	MPD	A	707	-	-	0/5/5/5	-
3	MPD	B	702	-	-	0/5/5/5	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	705	MPD	C2-C3-C4-C5
3	A	708	MPD	C2-C3-C4-C5
3	B	701	MPD	C2-C3-C4-C5
3	A	705	MPD	C2-C3-C4-O4
3	B	701	MPD	C2-C3-C4-O4

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	702	SO4	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	598/631 (94%)	-0.52	3 (0%) 91 90	18, 25, 40, 70	0
1	B	597/631 (94%)	-0.22	14 (2%) 60 59	20, 29, 49, 80	0
All	All	1195/1262 (94%)	-0.37	17 (1%) 75 75	18, 27, 44, 80	0

All (17) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	170	GLY	5.5
1	A	170	GLY	4.6
1	B	219	LEU	4.2
1	B	258	THR	3.6
1	B	166	SER	3.4
1	A	16	ASN	3.3
1	B	222	ASN	3.3
1	B	223	THR	3.3
1	B	221	ASN	3.0
1	B	262	LEU	2.6
1	B	264	ILE	2.4
1	B	266	ASN	2.3
1	B	137	ASN	2.3
1	B	263	THR	2.1
1	A	168	VAL	2.1
1	B	224	ALA	2.1
1	B	397	VAL	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	SO4	A	703	5/5	0.85	0.16	45,46,56,58	5
2	SO4	A	702	5/5	0.86	0.17	41,44,50,50	5
3	MPD	A	707	8/8	0.86	0.16	34,39,43,46	0
2	SO4	A	706	5/5	0.89	0.14	48,51,56,58	5
3	MPD	B	702	8/8	0.90	0.14	43,48,49,49	0
3	MPD	A	705	8/8	0.92	0.11	32,34,40,49	0
2	SO4	A	701	5/5	0.92	0.14	44,45,53,56	5
2	SO4	B	703	5/5	0.92	0.14	39,51,53,55	5
2	SO4	A	704	5/5	0.93	0.13	42,42,45,51	5
3	MPD	B	701	8/8	0.94	0.09	35,38,43,51	0
2	SO4	B	705	5/5	0.94	0.22	37,50,53,55	5
2	SO4	B	704	5/5	0.95	0.11	42,45,55,57	5
3	MPD	A	708	8/8	0.95	0.16	38,42,45,46	0
2	SO4	B	706	5/5	0.98	0.11	39,39,43,43	5
4	CL	B	708	1/1	0.99	0.05	29,29,29,29	0
4	CL	A	710	1/1	1.00	0.02	28,28,28,28	0
4	CL	B	707	1/1	1.00	0.02	28,28,28,28	0
4	CL	A	709	1/1	1.00	0.03	25,25,25,25	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.