



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 5, 2023 – 03:47 PM EDT

PDB ID : 4DIF
Title : Structure of A1-type ketoreductase
Authors : Zheng, J.; Keatinge-Clay, A.T.
Deposited on : 2012-01-30
Resolution : 1.52 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.52 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4009 (1.54-1.50)
Clashscore	141614	4249 (1.54-1.50)
Ramachandran outliers	138981	4148 (1.54-1.50)
Sidechain outliers	138945	4146 (1.54-1.50)
RSRZ outliers	127900	3943 (1.54-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	496	 4% 88% 8%
1	B	496	 4% 87% 9%

2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 7653 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called AmphB.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	476	3469	2166	636	658	9	0	0	0
1	B	476	3469	2166	636	658	9	0	0	0

There are 46 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-20	MET	-	expression tag	UNP Q93NW7
A	-19	GLY	-	expression tag	UNP Q93NW7
A	-18	SER	-	expression tag	UNP Q93NW7
A	-17	SER	-	expression tag	UNP Q93NW7
A	-16	HIS	-	expression tag	UNP Q93NW7
A	-15	HIS	-	expression tag	UNP Q93NW7
A	-14	HIS	-	expression tag	UNP Q93NW7
A	-13	HIS	-	expression tag	UNP Q93NW7
A	-12	HIS	-	expression tag	UNP Q93NW7
A	-11	HIS	-	expression tag	UNP Q93NW7
A	-10	SER	-	expression tag	UNP Q93NW7
A	-9	SER	-	expression tag	UNP Q93NW7
A	-8	GLY	-	expression tag	UNP Q93NW7
A	-7	LEU	-	expression tag	UNP Q93NW7
A	-6	VAL	-	expression tag	UNP Q93NW7
A	-5	PRO	-	expression tag	UNP Q93NW7
A	-4	ARG	-	expression tag	UNP Q93NW7
A	-3	GLY	-	expression tag	UNP Q93NW7
A	-2	SER	-	expression tag	UNP Q93NW7
A	-1	HIS	-	expression tag	UNP Q93NW7
A	0	MET	-	expression tag	UNP Q93NW7
A	355	THR	GLY	engineered mutation	UNP Q93NW7
A	364	HIS	GLN	engineered mutation	UNP Q93NW7
B	-20	MET	-	expression tag	UNP Q93NW7
B	-19	GLY	-	expression tag	UNP Q93NW7

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	-18	SER	-	expression tag	UNP Q93NW7
B	-17	SER	-	expression tag	UNP Q93NW7
B	-16	HIS	-	expression tag	UNP Q93NW7
B	-15	HIS	-	expression tag	UNP Q93NW7
B	-14	HIS	-	expression tag	UNP Q93NW7
B	-13	HIS	-	expression tag	UNP Q93NW7
B	-12	HIS	-	expression tag	UNP Q93NW7
B	-11	HIS	-	expression tag	UNP Q93NW7
B	-10	SER	-	expression tag	UNP Q93NW7
B	-9	SER	-	expression tag	UNP Q93NW7
B	-8	GLY	-	expression tag	UNP Q93NW7
B	-7	LEU	-	expression tag	UNP Q93NW7
B	-6	VAL	-	expression tag	UNP Q93NW7
B	-5	PRO	-	expression tag	UNP Q93NW7
B	-4	ARG	-	expression tag	UNP Q93NW7
B	-3	GLY	-	expression tag	UNP Q93NW7
B	-2	SER	-	expression tag	UNP Q93NW7
B	-1	HIS	-	expression tag	UNP Q93NW7
B	0	MET	-	expression tag	UNP Q93NW7
B	355	THR	GLY	engineered mutation	UNP Q93NW7
B	364	HIS	GLN	engineered mutation	UNP Q93NW7

- Molecule 2 is NADPH DIHYDRO-NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NDP) (formula: $C_{21}H_{30}N_7O_{17}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	48	21	7	17	3	0	0
2	B	1	48	21	7	17	3	0	0

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
3	B	1	6	3	3	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
4	A	309	309	309	0	0
4	B	304	304	304	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: AmphB



- Molecule 1: AmphB



4 Data and refinement statistics i

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	61.55Å 63.72Å 71.98Å 72.92° 67.22° 89.74°	Depositor
Resolution (Å)	50.00 – 1.52 26.61 – 1.52	Depositor EDS
% Data completeness (in resolution range)	86.9 (50.00-1.52) 87.0 (26.61-1.52)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	6.98 (at 1.52Å)	Xtrriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.198 , 0.224 0.196 , 0.221	Depositor DCC
R_{free} test set	6435 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	12.7	Xtrriage
Anisotropy	0.043	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.41 , 53.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.064 for h,-k,h-l	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	7653	wwPDB-VP
Average B, all atoms (Å ²)	17.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.25% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, NDP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.54	1/3540 (0.0%)	0.50	0/4837
1	B	0.53	0/3540	0.51	0/4837
All	All	0.53	1/7080 (0.0%)	0.51	0/9674

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	B	0	2
All	All	0	4

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	396	TRP	NE1-CE2	-5.98	1.29	1.37

There are no bond angle outliers.

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	124	ARG	Sidechain
1	A	401	MET	Mainchain
1	B	124	ARG	Sidechain
1	B	405	PRO	Mainchain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3469	0	3429	39	0
1	B	3469	0	3429	49	0
2	A	48	0	26	1	0
2	B	48	0	26	1	0
3	B	6	0	8	1	0
4	A	309	0	0	5	0
4	B	304	0	0	6	0
All	All	7653	0	6918	88	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (88) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:407:VAL:O	1:B:411:LEU:HD13	1.55	1.07
1:A:124:ARG:HE	1:A:166:GLU:HA	1.15	1.04
1:A:90:HIS:HD2	1:A:92:ASP:H	1.08	0.98
1:A:124:ARG:NE	1:A:166:GLU:HA	1.79	0.97
1:B:19:ARG:NH2	1:B:45:LEU:O	1.99	0.95
1:A:341:THR:HA	1:A:344:LEU:HD13	1.48	0.95
1:B:208:VAL:HG23	1:B:211:SER:OG	1.69	0.92
1:B:41:VAL:CG2	1:B:164:LEU:HD11	2.01	0.88
1:B:411:LEU:HD12	1:B:411:LEU:N	1.96	0.79
1:B:407:VAL:O	1:B:411:LEU:CD1	2.29	0.78
1:B:213:LYS:HE3	1:B:435:ASP:CB	2.15	0.77
1:B:59:ARG:HB2	1:B:105:LEU:CD1	2.15	0.76
1:B:59:ARG:HB2	1:B:105:LEU:HD12	1.68	0.76
1:B:404:ASP:C	1:B:406:GLU:H	1.87	0.75
1:A:139:VAL:HG22	4:A:859:HOH:O	1.87	0.74
1:A:90:HIS:CD2	1:A:92:ASP:H	2.01	0.70
1:A:203:HIS:HD2	4:A:740:HOH:O	1.75	0.69
1:B:139:VAL:HG22	4:B:685:HOH:O	1.91	0.69
1:B:213:LYS:HE3	1:B:435:ASP:HB2	1.72	0.69
1:A:203:HIS:HE1	1:A:467:GLU:OE2	1.76	0.68

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:411:LEU:N	1:B:411:LEU:CD1	2.57	0.67
1:A:364:HIS:HD2	4:A:685:HOH:O	1.78	0.67
1:B:245:HIS:HE1	1:B:274:THR:OG1	1.80	0.64
1:A:166:GLU:H	1:A:166:GLU:CD	1.98	0.64
1:A:402:ALA:O	1:A:408:HIS:HB2	1.98	0.63
1:A:245:HIS:HE1	1:A:274:THR:OG1	1.81	0.62
1:A:341:THR:CA	1:A:344:LEU:HD13	2.26	0.61
1:B:23:ARG:NH1	1:B:73:GLU:OE1	2.34	0.61
1:B:213:LYS:HE3	1:B:435:ASP:HB3	1.83	0.60
1:B:41:VAL:HG21	1:B:164:LEU:HD11	1.81	0.60
1:B:87:GLU:OE1	1:B:139:VAL:HG21	2.02	0.60
1:A:124:ARG:CZ	1:A:166:GLU:HA	2.32	0.59
1:B:402:ALA:O	1:B:408:HIS:HB2	2.03	0.59
1:B:41:VAL:CG2	1:B:164:LEU:CD1	2.79	0.58
1:B:124:ARG:HE	1:B:166:GLU:HA	1.66	0.58
1:B:208:VAL:HG23	1:B:211:SER:HG	1.68	0.58
1:B:411:LEU:CD1	1:B:411:LEU:H	2.17	0.57
1:A:41:VAL:CG2	1:A:164:LEU:HD11	2.35	0.57
1:B:364:HIS:HD2	4:B:661:HOH:O	1.88	0.57
1:A:77:HIS:HE1	1:A:180:LEU:O	1.88	0.57
1:A:391:VAL:HG21	1:A:428:LEU:HD13	1.87	0.57
1:B:380:HIS:HE1	4:B:789:HOH:O	1.87	0.56
1:B:391:VAL:HG21	1:B:428:LEU:HD13	1.87	0.56
1:A:41:VAL:HG21	1:A:164:LEU:HD11	1.87	0.56
1:B:404:ASP:C	1:B:406:GLU:N	2.59	0.55
1:B:87:GLU:OE1	1:B:139:VAL:CG2	2.55	0.54
1:B:404:ASP:O	1:B:406:GLU:N	2.39	0.53
1:B:37:TRP:HE3	1:B:164:LEU:HD12	1.74	0.53
1:B:237:ARG:HD2	1:B:425:LEU:HD13	1.91	0.53
1:B:77:HIS:HE1	1:B:180:LEU:O	1.92	0.53
1:B:278:CYS:SG	4:B:866:HOH:O	2.53	0.52
1:A:344:LEU:HD12	1:A:344:LEU:N	2.25	0.52
1:B:237:ARG:CD	1:B:425:LEU:HD13	2.40	0.51
1:A:450:ALA:HB3	1:A:451:PRO:HD3	1.93	0.51
1:A:335:ARG:HD2	4:A:778:HOH:O	2.09	0.51
1:A:224:THR:OG1	1:A:304:HIS:CD2	2.64	0.50
1:B:135:LEU:O	1:B:380:HIS:HD2	1.93	0.50
1:A:124:ARG:NH1	4:A:735:HOH:O	2.45	0.50
1:B:70:ARG:HH11	1:B:70:ARG:HG3	1.77	0.50
1:B:37:TRP:CE3	1:B:164:LEU:HD12	2.47	0.49
1:A:135:LEU:O	1:A:380:HIS:HD2	1.95	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:57:THR:OG1	1:A:90:HIS:HE1	1.96	0.48
1:A:224:THR:OG1	1:A:304:HIS:HD2	1.97	0.47
1:B:125:GLY:HA2	4:B:733:HOH:O	2.15	0.47
1:B:208:VAL:HG23	1:B:211:SER:CB	2.44	0.47
1:B:41:VAL:HG22	1:B:164:LEU:HD11	1.91	0.46
1:B:164:LEU:HD13	1:B:168:VAL:HG21	1.98	0.45
1:A:41:VAL:CG2	1:A:164:LEU:CD1	2.95	0.45
1:B:429:ASP:OD1	3:B:502:GOL:H12	2.17	0.44
1:A:330:LYS:HD3	1:A:374:LEU:HD11	1.98	0.44
1:A:341:THR:HA	1:A:344:LEU:CD1	2.33	0.44
1:A:220:SER:OG	1:A:245:HIS:HD2	2.02	0.42
1:B:404:ASP:OD1	1:B:406:GLU:HB3	2.20	0.42
1:A:393:TRP:HB2	2:A:501:NDP:C5N	2.49	0.42
1:B:19:ARG:HG2	1:B:177:THR:HB	2.01	0.42
1:A:124:ARG:NE	1:A:166:GLU:CA	2.68	0.42
1:A:37:TRP:CE3	1:A:164:LEU:HD12	2.54	0.42
1:A:364:HIS:N	1:A:365:PRO:CD	2.83	0.42
1:B:292:GLU:HG2	4:B:811:HOH:O	2.19	0.42
1:A:304:HIS:CE1	1:A:329:ALA:O	2.73	0.41
1:A:344:LEU:N	1:A:344:LEU:CD1	2.84	0.41
1:A:411:LEU:O	1:A:414:GLN:HG2	2.20	0.41
1:A:89:LEU:HD11	1:A:335:ARG:HH22	1.86	0.41
1:B:393:TRP:HB2	2:B:501:NDP:C5N	2.50	0.41
1:B:450:ALA:HB3	1:B:451:PRO:HD3	2.03	0.41
1:B:213:LYS:CE	1:B:435:ASP:HB3	2.50	0.40
1:A:304:HIS:HE1	1:A:329:ALA:O	2.04	0.40
1:B:41:VAL:HG11	1:B:82:LEU:HD21	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	474/496 (96%)	464 (98%)	10 (2%)	0	100	100
1	B	474/496 (96%)	464 (98%)	9 (2%)	1 (0%)	47	23
All	All	948/992 (96%)	928 (98%)	19 (2%)	1 (0%)	51	25

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	405	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	335/352 (95%)	334 (100%)	1 (0%)	92	84
1	B	335/352 (95%)	335 (100%)	0	100	100
All	All	670/704 (95%)	669 (100%)	1 (0%)	93	86

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	311	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (20) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	6	HIS
1	A	10	ASN
1	A	77	HIS
1	A	90	HIS
1	A	102	GLN
1	A	203	HIS
1	A	245	HIS
1	A	304	HIS
1	A	336	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	364	HIS
1	A	380	HIS
1	B	32	HIS
1	B	77	HIS
1	B	102	GLN
1	B	241	GLN
1	B	245	HIS
1	B	336	HIS
1	B	364	HIS
1	B	380	HIS
1	B	423	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	GOL	B	502	-	5,5,5	0.22	0	5,5,5	0.33	0
2	NDP	A	501	-	45,52,52	1.36	4 (8%)	53,80,80	1.28	2 (3%)
2	NDP	B	501	-	45,52,52	1.28	3 (6%)	53,80,80	1.35	4 (7%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	B	502	-	-	0/4/4/4	-
2	NDP	A	501	-	-	7/30/77/77	0/5/5/5
2	NDP	B	501	-	-	7/30/77/77	0/5/5/5

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	501	NDP	C4N-C3N	-5.40	1.39	1.49
2	B	501	NDP	C4N-C3N	-5.31	1.39	1.49
2	A	501	NDP	C4N-C5N	-3.97	1.38	1.48
2	B	501	NDP	C4N-C5N	-3.77	1.39	1.48
2	A	501	NDP	C6N-C5N	3.21	1.39	1.33
2	B	501	NDP	C6N-C5N	2.76	1.38	1.33
2	A	501	NDP	P2B-O2B	2.38	1.63	1.59

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	NDP	N3A-C2A-N1A	-5.46	120.14	128.68
2	B	501	NDP	N3A-C2A-N1A	-5.34	120.34	128.68
2	B	501	NDP	C4A-C5A-N7A	-2.40	106.90	109.40
2	A	501	NDP	C3N-C2N-N1N	-2.25	119.89	123.10
2	B	501	NDP	O2B-P2B-O1X	-2.19	100.95	109.39
2	B	501	NDP	O3X-P2B-O2X	2.09	115.61	107.64

There are no chirality outliers.

All (14) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	501	NDP	C5D-O5D-PN-O2N
2	B	501	NDP	C5D-O5D-PN-O2N
2	A	501	NDP	O4D-C1D-N1N-C6N
2	B	501	NDP	O4D-C1D-N1N-C6N
2	A	501	NDP	C5D-O5D-PN-O3
2	A	501	NDP	C5D-O5D-PN-O1N
2	B	501	NDP	C5D-O5D-PN-O1N
2	B	501	NDP	C5D-O5D-PN-O3

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
2	A	501	NDP	O4B-C4B-C5B-O5B
2	B	501	NDP	O4B-C4B-C5B-O5B
2	A	501	NDP	PA-O3-PN-O2N
2	B	501	NDP	PA-O3-PN-O2N
2	A	501	NDP	C2N-C3N-C7N-N7N
2	B	501	NDP	C2N-C3N-C7N-N7N


There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	502	GOL	1	0
2	A	501	NDP	1	0
2	B	501	NDP	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	476/496 (95%)	0.37	21 (4%) 34 38	8, 14, 32, 79	0
1	B	476/496 (95%)	0.38	20 (4%) 36 40	8, 14, 32, 66	0
All	All	952/992 (95%)	0.38	41 (4%) 35 39	8, 14, 32, 79	0

All (41) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	405	PRO	7.6
1	A	407	VAL	7.3
1	A	406	GLU	5.5
1	A	412	VAL	5.2
1	A	411	LEU	5.1
1	A	410	ARG	4.8
1	A	405	PRO	4.7
1	A	208	VAL	4.6
1	A	413	ARG	4.2
1	B	411	LEU	3.8
1	A	131	ARG	3.7
1	A	21	ALA	3.7
1	B	413	ARG	3.4
1	B	131	ARG	3.2
1	A	213	LYS	3.1
1	B	406	GLU	3.1
1	B	410	ARG	3.0
1	B	208	VAL	2.9
1	A	212	GLY	2.9
1	A	295	GLU	2.9
1	A	404	ASP	2.8
1	A	209	PRO	2.8
1	B	407	VAL	2.8
1	A	-1	HIS	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	302	VAL	2.6
1	B	351	LEU	2.5
1	A	409	ASP	2.5
1	A	408	HIS	2.5
1	B	132	GLY	2.4
1	A	342	ALA	2.4
1	A	352	PHE	2.3
1	B	260	ALA	2.3
1	B	-1	HIS	2.2
1	B	71	GLU	2.2
1	B	473	SER	2.2
1	B	412	VAL	2.1
1	B	343	ASP	2.1
1	B	21	ALA	2.1
1	B	211	SER	2.1
1	B	74	GLY	2.1
1	A	166	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	GOL	B	502	6/6	0.92	0.11	21,21,23,23	0
2	NDP	B	501	48/48	0.97	0.07	10,11,13,14	0
2	NDP	A	501	48/48	0.98	0.07	11,12,15,18	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers

as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.