



# Full wwPDB X-ray Structure Validation Report i

Jan 30, 2021 – 02:33 PM EST

PDB ID : 3DLS  
Title : Crystal structure of human PAS kinase bound to ADP  
Authors : Antonysamy, S.; Bonanno, J.B.; Romero, R.; Russell, M.; Iizuka, M.; Gheyi, T.; Wasserman, S.R.; Rutter, J.; Sauder, J.M.; Burley, S.K.; New York SGX Research Center for Structural Genomics (NYSGXRC)  
Deposited on : 2008-06-29  
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>  
with specific help available everywhere you see the i symbol.

---

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.16  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.16















## 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	85.78Å 85.84Å 94.15Å 77.28° 77.50° 60.09°	Depositor
Resolution (Å)	(Not available) – 2.30 31.20 – 2.30	Depositor EDS
% Data completeness (in resolution range)	(Not available) ((Not available)-2.30) 95.9 (31.20-2.30)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) >$ <sup>1</sup>	1.42 (at 2.29Å)	Xtriage
Refinement program	REFMAC	Depositor
$R$ , $R_{free}$	0.241 , 0.297 0.243 , (Not available)	Depositor DCC
$R_{free}$ test set	4846 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	40.0	Xtriage
Anisotropy	0.092	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 23.9	EDS
L-test for twinning <sup>2</sup>	$<  L  > = 0.40$ , $< L^2 > = 0.22$	Xtriage
Estimated twinning fraction	0.216 for -k,-h,-l	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	13724	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.46% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $< |L| >$ ,  $< L^2 >$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.















*Continued from previous page...*

Mol	Chain	Res	Type
1	A	1153	LEU
1	A	1197	VAL
1	A	1206	LEU
1	A	1209	THR
1	A	1225	MET
1	A	1237	GLU
1	A	1247	THR
1	B	985	LEU
1	B	1032	LYS
1	B	1048	LYS
1	B	1070	ILE
1	B	1078	GLN
1	B	1122	LYS
1	B	1125	ILE
1	B	1153	LEU
1	B	1155	ARG
1	B	1157	LYS
1	C	1032	LYS
1	C	1043	ASP
1	C	1048	LYS
1	C	1051	LEU
1	C	1078	GLN
1	C	1079	LEU
1	C	1122	LYS
1	C	1125	ILE
1	C	1153	LEU
1	C	1209	THR
1	C	1245	LEU
1	D	985	LEU
1	D	1022	ASN
1	D	1032	LYS
1	D	1048	LYS
1	D	1070	ILE
1	D	1078	GLN
1	D	1079	LEU
1	D	1100	LEU
1	D	1122	LYS
1	D	1125	ILE
1	D	1153	LEU
1	D	1157	LYS
1	D	1209	THR
1	D	1219	LEU


*Continued on next page...*

















## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data i

### 6.1 Protein, DNA and RNA chains i

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å <sup>2</sup> )	Q<0.9
1	A	285/335 (85%)	-0.14	1 (0%)	92	95	14, 29, 46, 59
1	B	280/335 (83%)	0.01	7 (2%)	57	64	17, 33, 53, 72
1	C	281/335 (83%)	-0.05	6 (2%)	63	70	17, 32, 54, 73
1	D	284/335 (84%)	-0.01	2 (0%)	87	91	22, 38, 56, 70
1	E	282/335 (84%)	0.12	5 (1%)	68	74	22, 42, 58, 73
1	F	281/335 (83%)	0.22	10 (3%)	42	49	26, 46, 67, 76
All	All	1693/2010 (84%)	0.02	31 (1%)	68	74	14, 37, 59, 76

All (31) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	1204	CYS	4.8
1	C	1046	LEU	4.6
1	F	1121	LEU	4.0
1	F	1041	ILE	3.6
1	C	1048	LYS	3.5
1	B	1155	ARG	3.5
1	B	1040	TRP	3.4
1	B	1007	SER	3.3
1	F	1257	LEU	3.1
1	B	1041	ILE	2.9
1	F	1017	VAL	2.8
1	F	1235	VAL	2.8
1	C	1041	ILE	2.7
1	A	1007	SER	2.7
1	F	990	ALA	2.7
1	E	1096	ARG	2.7
1	E	1022	ASN	2.5
1	F	1070	ILE	2.5
1	C	1044	PRO	2.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	1097	HIS	2.4
1	E	1007	SER	2.3
1	B	1046	LEU	2.3
1	C	1049	VAL	2.3
1	B	1204	CYS	2.2
1	F	1209	THR	2.2
1	D	1074	GLN	2.2
1	D	1010	PHE	2.2
1	F	1009	ALA	2.1
1	F	991	CYS	2.1
1	E	1035	VAL	2.0
1	E	1156	GLY	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	MG	A	20	1/1	0.89	0.18	24,24,24,24	0
2	MG	B	19	1/1	0.90	0.20	33,33,33,33	0
2	MG	C	12	1/1	0.93	0.30	35,35,35,35	0
2	MG	A	8	1/1	0.93	0.27	28,28,28,28	0
2	MG	E	16	1/1	0.94	0.25	36,36,36,36	0
3	ADP	E	5	27/27	0.94	0.14	27,36,52,53	0
2	MG	C	22	1/1	0.94	0.09	35,35,35,35	0
2	MG	F	18	1/1	0.94	0.25	31,31,31,31	0
2	MG	D	14	1/1	0.95	0.29	33,33,33,33	0
2	MG	E	15	1/1	0.95	0.21	24,24,24,24	0
2	MG	B	10	1/1	0.95	0.18	23,23,23,23	0


*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	MG	F	17	1/1	0.96	0.24	25,25,25,25	0
2	MG	A	25	1/1	0.96	0.23	22,22,22,22	0
3	ADP	F	6	27/27	0.96	0.10	34,40,45,46	0
3	ADP	D	4	27/27	0.96	0.13	22,32,37,43	0
2	MG	D	23	1/1	0.96	0.12	31,31,31,31	0
2	MG	A	7	1/1	0.96	0.14	10,10,10,10	0
3	ADP	B	2	27/27	0.97	0.13	12,23,34,41	0
3	ADP	C	3	27/27	0.98	0.12	17,21,30,36	0
2	MG	B	24	1/1	0.98	0.21	26,26,26,26	0
2	MG	B	9	1/1	0.98	0.16	15,15,15,15	0
2	MG	D	13	1/1	0.98	0.20	14,14,14,14	0
3	ADP	A	1	27/27	0.98	0.11	13,20,33,36	0
2	MG	C	21	1/1	0.98	0.27	30,30,30,30	0
2	MG	C	11	1/1	0.99	0.20	13,13,13,13	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.









**Electron density around ADP B 2:**

2mF<sub>o</sub>-DF<sub>c</sub> (at 0.7 rmsd) in gray  
mF<sub>o</sub>-DF<sub>c</sub> (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around ADP C 3:**

2mF<sub>o</sub>-DF<sub>c</sub> (at 0.7 rmsd) in gray  
mF<sub>o</sub>-DF<sub>c</sub> (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.