



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 23, 2023 – 02:03 PM EST

PDB ID : 3DMB  
Title : Crystal structure of a putative general stress family protein (xcc2264) from xanthomonas campestris pv. campestris at 2.30 Å resolution  
Authors : Joint Center for Structural Genomics (JCSG)  
Deposited on : 2008-06-30  
Resolution : 2.30 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.31.2  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.31.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	147	
1	B	147	
1	C	147	

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 3303 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative General Stress Protein 26 with a PNP-Oxidase like Fold.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	Se			
1	A	146	1117	712	192	207	6	0	3	0
1	B	145	1100	700	188	208	4	0	0	0
1	C	134	1022	657	174	187	4	0	1	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	GLY	-	expression tag	UNP Q8P8H7
B	0	GLY	-	expression tag	UNP Q8P8H7
C	0	GLY	-	expression tag	UNP Q8P8H7


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	26	Total 26	O 26	0	0
2	B	18	Total 18	O 18	0	0
2	C	20	Total 20	O 20	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Putative General Stress Protein 26 with a PNP-Oxidase like Fold



- Molecule 1: Putative General Stress Protein 26 with a PNP-Oxidase like Fold



- Molecule 1: Putative General Stress Protein 26 with a PNP-Oxidase like Fold



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	105.32Å 105.32Å 68.34Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	28.66 – 2.30 28.66 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.7 (28.66-2.30) 99.7 (28.66-2.30)	Depositor EDS
$R_{merge}$	0.05	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.49 (at 2.29Å)	Xtrriage
Refinement program	REFMAC 5.2.0019, PHENIX	Depositor
R, $R_{free}$	0.210 , 0.264 0.214 , 0.261	Depositor DCC
$R_{free}$ test set	1006 reflections (5.10%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	48.8	Xtrriage
Anisotropy	0.290	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 65.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.035 for -h,-k,l	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	3303	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	67.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.42% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.70	0/1147	0.86	2/1547 (0.1%)
1	B	0.63	0/1122	0.81	0/1516
1	C	0.62	0/1045	0.81	0/1408
All	All	0.65	0/3314	0.83	2/4471 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	88	ARG	NE-CZ-NH1	7.05	123.82	120.30
1	A	67	ARG	NE-CZ-NH1	5.23	122.92	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1117	0	1100	10	0
1	B	1100	0	1072	12	0
1	C	1022	0	1013	3	0
2	A	26	0	0	0	0
2	B	18	0	0	0	0
2	C	20	0	0	0	0
All	All	3303	0	3185	21	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (21) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:25:LEU:HD22	1:A:61[A]:MSE:HE3	1.45	0.97
1:B:25:LEU:HD23	1:B:68:VAL:HG12	1.61	0.81
1:B:28:VAL:HG12	1:B:29:GLU:HG2	1.68	0.75
1:B:25:LEU:HD12	1:B:58:LEU:HD22	1.75	0.67
1:B:25:LEU:CD2	1:B:68:VAL:HG12	2.28	0.64
1:A:137:LEU:HD22	1:B:106:ALA:HB2	1.83	0.59
1:B:39:GLN:HE22	1:B:104:VAL:HG22	1.69	0.58
1:A:25:LEU:HD22	1:A:61[A]:MSE:CE	2.29	0.54
1:B:25:LEU:CD1	1:B:58:LEU:HD22	2.38	0.53
1:A:1:MSE:HE2	1:C:103:TYR:CD2	2.44	0.52
1:B:25:LEU:HD11	1:B:58:LEU:HD13	1.91	0.52
1:B:25:LEU:HD23	1:B:68:VAL:CG1	2.36	0.50
1:A:25:LEU:HD12	1:A:68:VAL:HG12	1.94	0.49
1:B:95:VAL:HG22	1:B:98:ARG:HH21	1.78	0.48
1:A:58:LEU:HA	1:A:61[A]:MSE:HE2	1.94	0.48
1:B:25:LEU:CD2	1:B:68:VAL:CG1	2.91	0.47
1:A:136:LEU:HD21	1:C:35:PRO:HG2	1.96	0.46
1:A:2:ALA:HB2	1:C:103:TYR:CE2	2.50	0.46
1:B:25:LEU:HD22	1:B:66:ARG:HD2	1.97	0.45
1:A:22[B]:MSE:HE3	1:A:35:PRO:HB3	1.98	0.44
1:A:25:LEU:HD13	1:A:58:LEU:HD22	2.01	0.42

There are no symmetry-related clashes.

## 5.3 Torsion angles [\(i\)](#)

### 5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	147/147 (100%)	141 (96%)	6 (4%)	0	100 100

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	143/147 (97%)	140 (98%)	3 (2%)	0	100	100
1	C	131/147 (89%)	127 (97%)	4 (3%)	0	100	100
All	All	421/441 (96%)	408 (97%)	13 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	113/111 (102%)	112 (99%)	1 (1%)	78	89
1	B	111/111 (100%)	111 (100%)	0	100	100
1	C	103/111 (93%)	98 (95%)	5 (5%)	25	35
All	All	327/333 (98%)	321 (98%)	6 (2%)	65	75

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	MSE
1	C	5	LYS
1	C	22[A]	MSE
1	C	22[B]	MSE
1	C	34	ARG
1	C	125	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	39	GLN



### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	142/147 (96%)	1.42	40 (28%) <b>0</b> <b>0</b>	57, 66, 80, 87	0
1	B	141/147 (95%)	0.98	17 (12%) <b>4</b> <b>6</b>	58, 67, 80, 91	0
1	C	131/147 (89%)	1.37	34 (25%) <b>0</b> <b>0</b>	57, 67, 78, 81	0
All	All	414/441 (93%)	1.25	91 (21%) <b>0</b> <b>1</b>	57, 67, 79, 91	0

All (91) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	2	ALA	7.5
1	A	110	GLY	5.6
1	B	109	GLU	5.5
1	B	110	GLY	5.4
1	A	144	LEU	5.4
1	C	6	GLU	5.2
1	A	29	GLU	5.0
1	A	0	GLY	4.8
1	C	33	ALA	4.8
1	A	107	TRP	4.7
1	C	126	HIS	4.5
1	A	115	PRO	4.4
1	C	8	GLN	4.3
1	A	108	TYR	4.3
1	C	56	ASN	4.2
1	C	5	LYS	4.2
1	C	58	LEU	4.1
1	B	108	TYR	4.1
1	A	21	VAL	4.1
1	C	59	ILE	4.0
1	C	67	ARG	3.9
1	B	113	ASP	3.9
1	A	142	VAL	3.9

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
1	A	106	ALA	3.9
1	B	3	ASP	3.8
1	A	105	ALA	3.8
1	B	4	PRO	3.6
1	A	23	LEU	3.6
1	B	21	VAL	3.6
1	A	109	GLU	3.6
1	C	21	VAL	3.5
1	A	120	LEU	3.5
1	C	63	GLY	3.5
1	A	4	PRO	3.4
1	C	7	LEU	3.4
1	A	83	ILE	3.3
1	A	145	GLY	3.3
1	C	125	ASP	3.3
1	C	53	SER	3.2
1	C	34	ARG	3.2
1	C	115	PRO	3.1
1	C	96	VAL	3.1
1	A	70	GLY	3.1
1	C	38	ALA	3.0
1	C	62	LEU	3.0
1	A	38	ALA	3.0
1	A	87	LEU	2.9
1	B	98	ARG	2.9
1	A	68	VAL	2.9
1	C	81	ALA	2.9
1	A	94	ALA	2.8
1	A	60	ALA	2.8
1	A	71	ALA	2.8
1	A	111	GLY	2.8
1	A	5	LYS	2.8
1	B	115	PRO	2.8
1	B	97	ASP	2.7
1	A	119	LEU	2.7
1	C	116	LYS	2.7
1	A	113	ASP	2.7
1	C	111	GLY	2.7
1	C	69	ILE	2.7
1	C	100	TRP	2.7
1	A	122	LEU	2.5
1	B	23	LEU	2.5

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	C	55	ASP	2.5
1	B	81	ALA	2.5
1	B	117	LEU	2.5
1	A	138	ALA	2.5
1	A	78	ASP	2.5
1	C	110	GLY	2.4
1	C	64	GLN	2.4
1	A	69	ILE	2.4
1	A	116	LYS	2.3
1	C	97	ASP	2.3
1	C	60	ALA	2.3
1	A	48	ILE	2.2
1	A	58	LEU	2.2
1	A	117	LEU	2.2
1	A	50	PHE	2.2
1	C	65	GLY	2.2
1	C	19	ARG	2.2
1	B	107	TRP	2.1
1	C	40	ILE	2.1
1	C	20	THR	2.1
1	A	15	LEU	2.1
1	A	37	THR	2.1
1	C	105	ALA	2.1
1	A	104	VAL	2.1
1	B	50	PHE	2.0
1	B	62	LEU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [\(i\)](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.