



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 12, 2024 – 10:12 PM EDT

PDB ID : 1DN2
Title : FC FRAGMENT OF HUMAN IGG1 IN COMPLEX WITH AN ENGINEERED 13 RESIDUE PEPTIDE DCAWHLGELVWCT-NH2
Authors : DeLano, W.L.; Ultsch, M.H.; de Vos, A.M.; Wells, J.A.
Deposited on : 1999-12-15
Resolution : 2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	3122 (2.70-2.70)
Ramachandran outliers	138981	3069 (2.70-2.70)
Sidechain outliers	138945	3069 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	207	
1	B	207	
2	E	14	
2	F	14	
3	C	8	
4	D	8	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	FUC	D	8	X	-	-	-

2 Entry composition i

There are 5 unique types of molecules in this entry. The entry contains 3779 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called IMMUNOGLOBULIN LAMBDA HEAVY CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	207	Total 1659	C 1056	N 280	O 316	S 7	19	0	0
1	B	207	Total 1659	C 1056	N 280	O 316	S 7	11	0	0


There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	270	ASN	ASP	conflict	GB 2765425
B	270	ASN	ASP	conflict	GB 2765425

- Molecule 2 is a protein called ENGINEERED PEPTIDE.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	E	14	Total 107	C 69	N 18	O 18	S 2	0	0	1
2	F	14	Total 107	C 69	N 18	O 18	S 2	0	0	1

- Molecule 3 is an oligosaccharide called beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[beta-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	C	8	Total 96	C 54	N 3	O 39	0	0	0

- Molecule 4 is an oligosaccharide called beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
4	D	8	96	54	3	39	0	0	0

- Molecule 5 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
5	A	33	33	33	0	0
5	B	22	22	22	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


Note EDS was not executed.

- Molecule 1: IMMUNOGLOBULIN LAMBDA HEAVY CHAIN

Chain A:  84% 16%



- Molecule 1: IMMUNOGLOBULIN LAMBDA HEAVY CHAIN

Chain B:  85% 14%



- Molecule 2: ENGINEERED PEPTIDE

Chain E:  93% 7%




- Molecule 2: ENGINEERED PEPTIDE

Chain F:  93% 7%



- Molecule 3: beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[beta-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose

Chain C:  62% 12% 25%



- Molecule 4: beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose

Chain D:  50% 50%

MAG1	MAG2	BRG3	MAN4	MAG5	GAL6	MAN7	FUC8
------	------	------	------	------	------	------	------

4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	67.54Å 60.83Å 68.17Å 90.00° 103.87° 90.00°	Depositor
Resolution (Å)	20.00 – 2.70	Depositor
% Data completeness (in resolution range)	96.1 (20.00-2.70)	Depositor
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	X-PLOR 3.851	Depositor
R, R_{free}	0.194 , 0.252	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	3779	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MAN, FUL, GAL, NAG, FUC, BMA, NH2

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.39	0/1705	0.61	0/2322
1	B	0.39	0/1705	0.61	0/2322
2	E	0.46	0/110	0.61	0/151
2	F	0.46	0/110	0.60	0/151
All	All	0.40	0/3630	0.61	0/4946

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1659	0	1627	21	0
1	B	1659	0	1627	21	0
2	E	107	0	93	1	0
2	F	107	0	93	1	0
3	C	96	0	82	2	0
4	D	96	0	82	0	0
5	A	33	0	0	0	0
5	B	22	0	0	0	0
All	All	3779	0	3604	42	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (42) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:238:PRO:HD2	1:A:328:LEU:HD13	1.56	0.88
1:B:242:LEU:HG	1:B:336:ILE:HG12	1.59	0.84
1:A:242:LEU:HG	1:A:336:ILE:HG12	1.62	0.79
1:B:346:PRO:HB3	1:B:372:PHE:HB3	1.65	0.79
1:A:346:PRO:HB3	1:A:372:PHE:HB3	1.64	0.78
1:B:238:PRO:HD2	1:B:328:LEU:HD13	1.73	0.70
1:A:434:ASN:OD1	2:E:13:THR:HG23	1.92	0.70
1:A:350:THR:HB	1:A:441:LEU:HD13	1.75	0.68
1:B:434:ASN:OD1	2:F:13:THR:HG23	1.99	0.63
1:B:443:LEU:HD12	1:B:443:LEU:N	2.15	0.61
1:B:350:THR:HB	1:B:441:LEU:HD13	1.85	0.58
1:B:351:LEU:HB2	1:B:366:THR:HB	1.90	0.54
1:B:365:LEU:HD12	1:B:410:LEU:HD23	1.89	0.54
1:B:429:HIS:O	1:B:435:HIS:HA	2.09	0.52
1:A:365:LEU:HD12	1:A:410:LEU:HD23	1.91	0.52
1:A:374:PRO:O	1:A:429:HIS:HE1	1.93	0.50
1:A:351:LEU:HB2	1:A:366:THR:HB	1.93	0.50
1:A:377:ILE:HG12	1:A:378:ALA:H	1.78	0.49
1:B:429:HIS:CD2	1:B:431:ALA:H	2.29	0.49
3:C:1:NAG:O6	3:C:8:FUL:H63	2.12	0.48
1:A:406:LEU:C	1:A:406:LEU:HD12	2.35	0.47
1:A:318:GLU:HA	1:A:337:SER:HB3	1.98	0.46
1:A:238:PRO:HD2	1:A:328:LEU:CD1	2.39	0.45
1:A:266:VAL:HB	1:A:300:TYR:HB2	1.98	0.45
1:A:429:HIS:CD2	1:A:431:ALA:H	2.35	0.45
1:B:374:PRO:O	1:B:429:HIS:HE1	1.98	0.45
1:A:377:ILE:HG12	1:A:378:ALA:N	2.31	0.45
1:A:354:SER:HB2	1:B:349:TYR:HB3	1.99	0.45
1:B:430:GLU:HA	1:B:435:HIS:CD2	2.51	0.45
1:A:367:CYS:HB2	1:A:381:TRP:CZ2	2.51	0.44
1:B:377:ILE:HG12	1:B:378:ALA:H	1.82	0.44
1:A:429:HIS:O	1:A:435:HIS:HA	2.18	0.43
1:B:377:ILE:HG12	1:B:378:ALA:N	2.33	0.43
1:B:238:PRO:HD2	1:B:328:LEU:CD1	2.45	0.43
1:B:406:LEU:HD12	1:B:406:LEU:C	2.39	0.42
1:B:276:ASN:HB2	1:B:322:LYS:HB3	2.02	0.42
1:B:266:VAL:HB	1:B:300:TYR:HB2	2.02	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:328:LEU:HD21	1:B:332:ILE:HD12	2.02	0.41
1:B:256:THR:HA	1:B:257:PRO:HD2	1.89	0.41
1:A:261:CYS:HB2	1:A:277:TRP:CH2	2.56	0.40
1:A:246:LYS:HG2	3:C:6:GAL:O4	2.22	0.40
1:A:276:ASN:HB2	1:A:322:LYS:HB3	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	205/207 (99%)	203 (99%)	2 (1%)	0	100	100
1	B	205/207 (99%)	199 (97%)	5 (2%)	1 (0%)	29	54
2	E	12/14 (86%)	12 (100%)	0	0	100	100
2	F	12/14 (86%)	11 (92%)	1 (8%)	0	100	100
All	All	434/442 (98%)	425 (98%)	8 (2%)	1 (0%)	47	73

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	358	MET

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	193/193 (100%)	192 (100%)	1 (0%)	88	96
1	B	193/193 (100%)	190 (98%)	3 (2%)	62	85
2	E	11/11 (100%)	11 (100%)	0	100	100
2	F	11/11 (100%)	11 (100%)	0	100	100
All	All	408/408 (100%)	404 (99%)	4 (1%)	76	91

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	355	ARG
1	B	350	THR
1	B	439	LYS
1	B	443	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	429	HIS
1	A	438	GLN
1	B	361	ASN
1	B	429	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

16 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAG	C	1	3,1	14,14,15	0.48	0	17,19,21	0.73	0
3	NAG	C	2	3	14,14,15	0.46	0	17,19,21	0.73	0
3	BMA	C	3	3	11,11,12	0.64	0	15,15,17	0.59	0
3	MAN	C	4	3	11,11,12	0.52	0	15,15,17	0.57	0
3	NAG	C	5	3	14,14,15	0.50	0	17,19,21	0.76	0
3	GAL	C	6	3	11,11,12	0.76	0	15,15,17	0.76	1 (6%)
3	MAN	C	7	3	11,11,12	0.62	0	15,15,17	0.80	0
3	FUL	C	8	3	10,10,11	0.75	0	14,14,16	1.14	2 (14%)
4	NAG	D	1	1,4	14,14,15	0.68	0	17,19,21	0.87	0
4	NAG	D	2	4	14,14,15	0.59	0	17,19,21	0.78	1 (5%)
4	BMA	D	3	4	11,11,12	0.81	1 (9%)	15,15,17	0.50	0
4	MAN	D	4	4	11,11,12	0.57	0	15,15,17	0.85	1 (6%)
4	NAG	D	5	4	14,14,15	0.49	0	17,19,21	0.79	0
4	GAL	D	6	4	11,11,12	0.56	0	15,15,17	0.51	0
4	MAN	D	7	4	11,11,12	0.54	0	15,15,17	0.59	0
4	FUC	D	8	4	10,10,11	0.76	0	14,14,16	1.04	1 (7%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	C	1	3,1	-	0/6/23/26	0/1/1/1
3	NAG	C	2	3	-	2/6/23/26	0/1/1/1
3	BMA	C	3	3	-	0/2/19/22	0/1/1/1
3	MAN	C	4	3	-	2/2/19/22	0/1/1/1
3	NAG	C	5	3	-	0/6/23/26	0/1/1/1
3	GAL	C	6	3	-	1/2/19/22	0/1/1/1
3	MAN	C	7	3	-	2/2/19/22	0/1/1/1
3	FUL	C	8	3	-	-	0/1/1/1
4	NAG	D	1	1,4	-	0/6/23/26	0/1/1/1
4	NAG	D	2	4	-	2/6/23/26	0/1/1/1
4	BMA	D	3	4	-	0/2/19/22	0/1/1/1
4	MAN	D	4	4	-	2/2/19/22	0/1/1/1
4	NAG	D	5	4	-	0/6/23/26	0/1/1/1
4	GAL	D	6	4	-	0/2/19/22	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	MAN	D	7	4	-	0/2/19/22	0/1/1/1
4	FUC	D	8	4	1/1/4/5	-	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	D	3	BMA	C2-C3	2.01	1.55	1.52

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	D	4	MAN	C1-O5-C5	2.59	115.71	112.19
4	D	8	FUC	C1-C2-C3	2.59	112.84	109.67
3	C	8	FUL	C3-C4-C5	2.47	113.62	109.77
3	C	8	FUL	C1-C2-C3	2.41	112.63	109.67
3	C	6	GAL	C1-C2-C3	2.30	112.50	109.67
4	D	2	NAG	C2-N2-C7	-2.10	119.92	122.90

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
4	D	8	FUC	C1

All (11) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	D	4	MAN	O5-C5-C6-O6
3	C	2	NAG	O5-C5-C6-O6
3	C	2	NAG	C4-C5-C6-O6
4	D	2	NAG	C4-C5-C6-O6
4	D	4	MAN	C4-C5-C6-O6
4	D	2	NAG	O5-C5-C6-O6
3	C	7	MAN	C4-C5-C6-O6
3	C	7	MAN	O5-C5-C6-O6
3	C	6	GAL	O5-C5-C6-O6
3	C	4	MAN	C4-C5-C6-O6
3	C	4	MAN	O5-C5-C6-O6


There are no ring outliers.

3 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	8	FUL	1	0
3	C	1	NAG	1	0
3	C	6	GAL	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.