



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 20, 2020 – 02:01 PM BST

PDB ID : 5DN5
Title : Structure of a C-terminally truncated glycoside hydrolase domain from *Salmonella typhimurium* FlgJ
Authors : Zaloba, P.; Bailey-Elkin, B.A.; Mark, B.L.
Deposited on : 2015-09-09
Resolution : 2.15 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.13.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1479 (2.16-2.16)
Clashscore	141614	1585 (2.16-2.16)
Ramachandran outliers	138981	1560 (2.16-2.16)
Sidechain outliers	138945	1559 (2.16-2.16)
RSRZ outliers	127900	1456 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	160	 6% 86% 8% • 6%
1	B	160	 13% 83% 11% • 6%
1	C	160	 18% 76% 12% 12%

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 6522 atoms, of which 3080 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Peptidoglycan hydrolase FlgJ.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	A	151	2188	707	1071	189	219	2	0	0	0
1	B	151	2178	706	1059	192	219	2	0	0	0
1	C	141	1964	645	950	174	194	1	0	0	0

There are 27 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	150	MET	-	initiating methionine	UNP P15931
A	302	GLY	-	expression tag	UNP P15931
A	303	SER	-	expression tag	UNP P15931
A	304	HIS	-	expression tag	UNP P15931
A	305	HIS	-	expression tag	UNP P15931
A	306	HIS	-	expression tag	UNP P15931
A	307	HIS	-	expression tag	UNP P15931
A	308	HIS	-	expression tag	UNP P15931
A	309	HIS	-	expression tag	UNP P15931
B	150	MET	-	initiating methionine	UNP P15931
B	302	GLY	-	expression tag	UNP P15931
B	303	SER	-	expression tag	UNP P15931
B	304	HIS	-	expression tag	UNP P15931
B	305	HIS	-	expression tag	UNP P15931
B	306	HIS	-	expression tag	UNP P15931
B	307	HIS	-	expression tag	UNP P15931
B	308	HIS	-	expression tag	UNP P15931
B	309	HIS	-	expression tag	UNP P15931
C	150	MET	-	initiating methionine	UNP P15931
C	302	GLY	-	expression tag	UNP P15931
C	303	SER	-	expression tag	UNP P15931
C	304	HIS	-	expression tag	UNP P15931
C	305	HIS	-	expression tag	UNP P15931

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	306	HIS	-	expression tag	UNP P15931
C	307	HIS	-	expression tag	UNP P15931
C	308	HIS	-	expression tag	UNP P15931
C	309	HIS	-	expression tag	UNP P15931

- Molecule 2 is IODIDE ION (three-letter code: IOD) (formula: I).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	3	Total I 3 3	0	0
2	A	2	Total I 2 2	0	0
2	C	5	Total I 5 5	0	0

- Molecule 3 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Na 1 1	0	0

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	C	2	Total Cl 2 2	0	0



- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	65	Total O 65 65	0	0
5	B	66	Total O 66 66	0	0
5	C	48	Total O 48 48	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: Peptidoglycan hydrolase FlgJ



- Molecule 1: Peptidoglycan hydrolase FlgJ



- Molecule 1: Peptidoglycan hydrolase FlgJ



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	105.70Å 61.12Å 65.22Å 90.00° 106.70° 90.00°	Depositor
Resolution (Å)	29.76 – 2.15 29.76 – 2.15	Depositor EDS
% Data completeness (in resolution range)	99.6 (29.76-2.15) 99.6 (29.76-2.15)	Depositor EDS
R_{merge}	0.15	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.38 (at 2.16Å)	Xtrriage
Refinement program	PHENIX 1.9_1692	Depositor
R, R_{free}	0.175 , 0.252 0.179 , 0.255	Depositor DCC
R_{free} test set	1096 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	24.6	Xtrriage
Anisotropy	0.172	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 64.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	6522	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.74% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NA, IOD, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.66	0/1139	0.66	0/1554
1	B	0.71	0/1141	0.72	1/1559 (0.1%)
1	C	0.62	0/1032	0.67	0/1408
All	All	0.67	0/3312	0.69	1/4521 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	194	ARG	NE-CZ-NH2	-6.99	116.81	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1117	1071	1071	7	0
1	B	1119	1059	1059	13	0
1	C	1014	950	949	11	0
2	A	2	0	0	0	0
2	B	3	0	0	0	0
2	C	5	0	0	0	0
3	A	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	C	2	0	0	0	0
5	A	65	0	0	1	0
5	B	66	0	0	1	0
5	C	48	0	0	1	0
All	All	3442	3080	3079	28	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (28) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:195:GLU:OE1	5:C:501:HOH:O	2.00	0.79
1:C:190:ARG:NH2	1:C:203:VAL:O	2.22	0.72
1:C:290:LEU:O	1:C:293:MET:N	2.25	0.68
1:C:269:GLU:N	1:C:269:GLU:OE2	2.26	0.68
1:B:155:PHE:CZ	1:B:159:LEU:HD12	2.40	0.56
1:A:289:LYS:NZ	5:A:501:HOH:O	2.27	0.56
1:B:194:ARG:HD3	1:B:238:VAL:HG21	1.89	0.54
1:B:285:ASN:OD1	1:B:286:TYR:N	2.41	0.54
1:B:194:ARG:HH22	1:C:198:GLU:CD	2.13	0.52
1:B:283:ASP:N	1:B:283:ASP:OD1	2.44	0.51
1:A:247:SER:HB2	1:B:161:LEU:HD13	1.91	0.51
1:B:255:ARG:NH2	1:C:154:ASP:OD2	2.38	0.51
1:A:261:ALA:HB3	1:A:274:ALA:HB1	1.94	0.49
1:B:271:GLY:HA2	1:B:274:ALA:HB3	1.95	0.49
1:B:156:LEU:HD12	1:B:297:LEU:HD22	1.97	0.47
1:A:151:ASP:O	1:A:153:LYS:N	2.48	0.47
1:C:296:GLN:O	1:C:300:MET:HG3	2.15	0.47
1:B:176:LEU:HD21	1:B:275:LEU:HD11	2.00	0.44
1:B:278:ALA:HA	5:B:522:HOH:O	2.18	0.44
1:B:184:GLU:OE2	1:B:286:TYR:OH	2.27	0.43
1:A:194:ARG:HA	1:A:217:THR:OG1	2.19	0.43
1:C:277:ASN:O	1:C:278:ALA:HB2	2.18	0.42
1:A:155:PHE:CE1	1:A:159:LEU:HD22	2.55	0.42
1:A:178:LEU:HB3	1:A:297:LEU:CD1	2.49	0.42
1:C:221:THR:OG1	1:C:232:VAL:HG13	2.19	0.41
1:B:267:THR:O	1:B:270:GLN:O	2.38	0.41
1:C:290:LEU:O	1:C:291:THR:C	2.59	0.41
1:C:254:THR:O	1:C:255:ARG:CB	2.69	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	149/160 (93%)	144 (97%)	3 (2%)	2 (1%)	12	6
1	B	149/160 (93%)	144 (97%)	5 (3%)	0	100	100
1	C	137/160 (86%)	129 (94%)	8 (6%)	0	100	100
All	All	435/480 (91%)	417 (96%)	16 (4%)	2 (0%)	29	22

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	152	SER
1	A	278	ALA

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	109/127 (86%)	108 (99%)	1 (1%)	78	83
1	B	107/127 (84%)	107 (100%)	0	100	100
1	C	91/127 (72%)	89 (98%)	2 (2%)	52	55
All	All	307/381 (81%)	304 (99%)	3 (1%)	76	81

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	247	SER
1	C	175	HIS
1	C	247	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	189	GLN
1	B	175	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 13 ligands modelled in this entry, 13 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	151/160 (94%)	0.48	10 (6%) 18 24	14, 31, 57, 67	0
1	B	151/160 (94%)	0.75	20 (13%) 3 4	15, 29, 59, 74	0
1	C	141/160 (88%)	0.94	28 (19%) 1 1	13, 38, 59, 69	0
All	All	443/480 (92%)	0.72	58 (13%) 3 4	13, 32, 59, 74	0

All (58) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	279	GLY	6.1
1	C	300	MET	5.6
1	A	284	PRO	5.2
1	C	254	THR	4.9
1	C	297	LEU	4.9
1	C	293	MET	4.8
1	B	300	MET	4.6
1	B	282	THR	4.5
1	B	299	ALA	4.5
1	C	299	ALA	4.3
1	B	284	PRO	4.1
1	B	297	LEU	4.1
1	B	301	SER	3.6
1	A	226	ASN	3.5
1	B	281	ALA	3.4
1	B	286	TYR	3.3
1	C	301	SER	3.3
1	B	285	ASN	3.3
1	B	243	LEU	3.2
1	C	290	LEU	3.2
1	A	282	THR	3.2
1	A	152	SER	3.1
1	A	210	ALA	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	260	ALA	3.1
1	C	278	ALA	3.1
1	C	255	ARG	3.1
1	C	273	VAL	2.9
1	C	259	TYR	2.9
1	B	283	ASP	2.9
1	B	156	LEU	2.8
1	B	278	ALA	2.8
1	B	280	TYR	2.7
1	C	242	TYR	2.6
1	A	286	TYR	2.6
1	B	153	LYS	2.6
1	C	243	LEU	2.6
1	A	228	GLU	2.5
1	C	274	ALA	2.5
1	C	271	GLY	2.5
1	C	267	THR	2.4
1	A	224	TYR	2.4
1	C	252	LEU	2.4
1	B	298	LYS	2.4
1	B	277	ASN	2.4
1	C	209	THR	2.4
1	C	159	LEU	2.4
1	A	285	ASN	2.4
1	A	301	SER	2.4
1	B	247	SER	2.3
1	C	266	ALA	2.3
1	C	156	LEU	2.3
1	C	275	LEU	2.3
1	C	268	ALA	2.2
1	C	241	SER	2.2
1	C	296	GLN	2.1
1	C	295	GLN	2.1
1	B	241	SER	2.1
1	C	201	TYR	2.1

6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	CL	C	406	1/1	0.88	0.26	37,37,37,37	0
4	CL	C	407	1/1	0.93	0.15	44,44,44,44	0
3	NA	A	403	1/1	0.95	0.24	33,33,33,33	0
2	IOD	C	401	1/1	0.98	0.06	47,47,47,47	1
2	IOD	C	402	1/1	0.99	0.06	40,40,40,40	1
2	IOD	C	403	1/1	0.99	0.09	36,36,36,36	1
2	IOD	B	401	1/1	0.99	0.10	39,39,39,39	1
2	IOD	A	401	1/1	1.00	0.14	20,20,20,20	1
2	IOD	B	403	1/1	1.00	0.12	22,22,22,22	0
2	IOD	A	402	1/1	1.00	0.10	26,26,26,26	0
2	IOD	B	402	1/1	1.00	0.10	25,25,25,25	0
2	IOD	C	404	1/1	1.00	0.11	23,23,23,23	0
2	IOD	C	405	1/1	1.00	0.13	22,22,22,22	0

6.5 Other polymers [i](#)

There are no such residues in this entry.