



# Full wwPDB X-ray Structure Validation Report ⓘ

Nov 20, 2023 – 09:40 PM JST

PDB ID : 7DNR  
Title : Crystal Structure of Zn-bound SIS Domain of Glucosamine-6-P Synthase from E. coli  
Authors : Gao, C.; Xiao, J.  
Deposited on : 2020-12-10  
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4298 (1.70-1.70)
Clashscore	141614	4695 (1.70-1.70)
Ramachandran outliers	138981	4610 (1.70-1.70)
Sidechain outliers	138945	4610 (1.70-1.70)
RSRZ outliers	127900	4222 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	368	 6% 86% 11% .
1	B	368	 6% 83% 12% . .

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 5976 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glutamine--fructose-6-phosphate aminotransferase [isomerizing].

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	357	2753	1741	472	528	12	0	0	0
1	B	354	2731	1728	466	525	12	0	0	0

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total 2	Zn 2	0	0
2	B	1	Total 1	Zn 1	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	228	Total 228	O 228	0	0
3	B	261	Total 261	O 261	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Glutamine--fructose-6-phosphate aminotransferase [isomerizing]



- Molecule 1: Glutamine--fructose-6-phosphate aminotransferase [isomerizing]



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	70.10Å 67.89Å 73.95Å 90.00° 111.44° 90.00°	Depositor
Resolution (Å)	23.52 – 1.70 32.10 – 1.60	Depositor EDS
% Data completeness (in resolution range)	100.0 (23.52-1.70) 96.2 (32.10-1.60)	Depositor EDS
$R_{merge}$	0.05	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.91 (at 1.60Å)	Xtrriage
Refinement program	PHENIX 1.18.2_3874	Depositor
R, $R_{free}$	0.214 , 0.235 0.211 , 0.228	Depositor DCC
$R_{free}$ test set	2003 reflections (2.35%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	23.2	Xtrriage
Anisotropy	0.261	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 34.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	5976	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	27.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.08% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section:  
ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.43	0/2800	0.63	3/3790 (0.1%)
1	B	0.48	0/2777	0.64	3/3759 (0.1%)
All	All	0.45	0/5577	0.64	6/7549 (0.1%)

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	564	GLU	CA-CB-CG	7.76	130.47	113.40
1	B	250	ARG	NE-CZ-NH2	-6.63	116.98	120.30
1	A	452	ASP	C-N-CA	-5.66	107.56	121.70
1	A	289	LEU	CB-CG-CD2	-5.61	101.46	111.00
1	A	292	LYS	CD-CE-NZ	-5.11	99.95	111.70
1	B	486	LYS	CB-CG-CD	-5.01	98.58	111.60

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2753	0	2773	29	0
1	B	2731	0	2750	39	0
2	A	2	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	1	0	0	0	0
3	A	228	0	0	13	0
3	B	261	0	0	16	0
All	All	5976	0	5523	65	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (65) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:393:ASN:ND2	3:A:802:HOH:O	2.15	0.79
1:B:259:GLU:OE1	3:B:804:HOH:O	2.02	0.77
1:B:512:ASP:HA	3:B:842:HOH:O	1.87	0.74
1:A:259:GLU:OE1	3:A:801:HOH:O	2.06	0.73
1:A:469:LEU:HD11	1:A:498:TYR:HD1	1.52	0.73
1:A:559:ASN:ND2	3:A:805:HOH:O	2.22	0.72
1:B:597:ASP:OD2	3:B:805:HOH:O	2.09	0.71
1:A:494:HIS:HE1	3:A:992:HOH:O	1.74	0.71
1:B:260:GLN:NE2	1:B:582:GLN:OE1	2.22	0.69
1:B:489:GLU:OE2	3:B:806:HOH:O	2.10	0.69
1:A:469:LEU:HD11	1:A:498:TYR:CD1	2.28	0.68
1:B:488:LYS:CE	3:B:802:HOH:O	2.38	0.66
1:B:439:GLN:OE1	3:B:807:HOH:O	2.14	0.65
1:B:439:GLN:NE2	3:B:803:HOH:O	1.98	0.65
1:B:564:GLU:OE2	3:B:808:HOH:O	2.15	0.64
1:A:402:SER:HB2	1:A:405:ALA:HB3	1.81	0.62
1:B:561:HIS:HE1	3:B:811:HOH:O	1.83	0.62
1:B:481:LEU:HD23	1:B:497:ALA:HB3	1.83	0.59
1:B:578:THR:HG23	3:B:905:HOH:O	2.03	0.58
1:A:559:ASN:CG	3:A:805:HOH:O	2.43	0.57
1:B:257:ILE:HA	1:B:260:GLN:HE21	1.70	0.57
1:A:456:GLU:OE1	3:A:803:HOH:O	2.17	0.57
1:B:488:LYS:HE3	3:B:802:HOH:O	2.05	0.56
1:A:347:LEU:HD23	1:A:374:ILE:HB	1.88	0.55
1:A:341:ASN:HA	1:A:369:LEU:HG	1.88	0.55
1:B:257:ILE:O	1:B:260:GLN:HG3	2.07	0.54
1:B:559:ASN:ND2	3:B:818:HOH:O	2.41	0.54
1:B:347:LEU:HD23	1:B:374:ILE:HB	1.90	0.53
1:A:304:SER:OG	1:A:349:GLN:OE1	2.27	0.52
1:B:294:GLU:OE1	1:B:340:ARG:NH2	2.42	0.52

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:285:ASN:ND2	3:B:812:HOH:O	2.30	0.52
1:A:509:ALA:O	1:A:540:ARG:NH2	2.44	0.51
1:A:444:ARG:NH2	1:A:569:GLU:OE2	2.41	0.51
1:B:502:GLU:C	1:B:504:LYS:H	2.16	0.49
1:A:591:ILE:HD11	3:A:825:HOH:O	2.13	0.49
1:B:568:VAL:HG21	1:B:576:PHE:CD2	2.47	0.49
1:A:339:ARG:HD3	3:A:859:HOH:O	2.13	0.48
1:A:498:TYR:CZ	1:A:507:PRO:HG3	2.49	0.48
1:B:561:HIS:CE1	3:B:811:HOH:O	2.63	0.48
1:A:507:PRO:O	1:A:511:ILE:HG13	2.15	0.47
1:A:490:ILE:HG22	1:A:589:ALA:HB2	1.96	0.47
1:B:260:GLN:N	1:B:261:PRO:CD	2.77	0.47
1:B:282:LEU:HD21	1:B:390:LEU:HG	1.96	0.47
1:B:441:LEU:HD13	1:B:572:ILE:HD12	1.96	0.47
1:B:594:THR:HG22	1:B:595:ASP:O	2.14	0.47
1:A:600:ARG:HB3	3:A:913:HOH:O	2.14	0.47
1:A:414:LEU:HG	1:A:434:ILE:HG23	1.98	0.45
1:B:439:GLN:HG2	3:B:803:HOH:O	2.15	0.45
1:B:488:LYS:NZ	3:B:802:HOH:O	1.86	0.45
1:A:594:THR:HG22	3:A:821:HOH:O	2.16	0.45
1:B:486:LYS:HD3	1:B:486:LYS:HA	1.63	0.44
1:A:284:PRO:O	3:A:804:HOH:O	2.21	0.43
1:A:543:GLN:NE2	3:A:818:HOH:O	2.51	0.43
1:A:510:LEU:HD11	1:B:488:LYS:HG3	2.01	0.42
1:B:282:LEU:HD22	1:B:388:LEU:HB3	2.02	0.41
1:B:348:SER:O	1:B:375:CYS:HA	2.20	0.41
1:A:333:TYR:CZ	1:B:529:LYS:HD3	2.55	0.41
1:B:305:TYR:CE2	1:B:309:MET:SD	3.14	0.41
1:B:414:LEU:HG	1:B:434:ILE:HG23	2.03	0.41
1:A:488:LYS:HD2	1:B:506:GLY:HA3	2.02	0.41
1:B:292:LYS:HB2	1:B:369:LEU:HD21	2.03	0.41
1:A:490:ILE:HG22	1:A:589:ALA:CB	2.51	0.41
1:B:488:LYS:HE3	1:B:488:LYS:HB2	1.87	0.40
1:B:498:TYR:CZ	1:B:507:PRO:HG3	2.56	0.40
1:A:439:GLN:NE2	3:A:808:HOH:O	2.36	0.40

There are no symmetry-related clashes.



## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	355/368 (96%)	350 (99%)	5 (1%)	0	100	100
1	B	352/368 (96%)	344 (98%)	8 (2%)	0	100	100
All	All	707/736 (96%)	694 (98%)	13 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	297/306 (97%)	295 (99%)	2 (1%)	84	77
1	B	295/306 (96%)	294 (100%)	1 (0%)	92	89
All	All	592/612 (97%)	589 (100%)	3 (0%)	88	83

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	285	ASN
1	A	558	ASP
1	B	402	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	447	GLN
1	A	494	HIS
1	A	550	GLN
1	B	561	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 3 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	357/368 (97%)	0.39	21 (5%) <span style="border: 1px solid red; padding: 2px;">22</span> <span style="border: 1px solid red; padding: 2px;">24</span>	13, 27, 49, 88	0
1	B	354/368 (96%)	0.38	23 (6%) <span style="border: 1px solid red; padding: 2px;">18</span> <span style="border: 1px solid red; padding: 2px;">21</span>	13, 22, 45, 73	0
All	All	711/736 (96%)	0.38	44 (6%) <span style="border: 1px solid red; padding: 2px;">20</span> <span style="border: 1px solid red; padding: 2px;">23</span>	13, 25, 47, 88	0

All (44) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	244	GLY	5.9
1	A	285	ASN	4.9
1	A	506	GLY	4.8
1	A	600	ARG	4.4
1	B	398	ILE	4.2
1	B	400	VAL	4.0
1	B	245	ASP	3.9
1	B	402	SER	3.8
1	B	288	GLU	3.8
1	B	525	GLU	3.6
1	A	498	TYR	3.5
1	A	555	VAL	3.3
1	B	432	HIS	3.2
1	B	309	MET	3.2
1	A	539	ALA	3.1
1	A	362	LEU	3.0
1	B	514	ASP	3.0
1	B	513	ALA	2.9
1	B	504	LYS	2.8
1	B	543	GLN	2.8
1	A	540	ARG	2.8
1	A	513	ALA	2.8
1	A	576	PHE	2.7
1	B	550	GLN	2.7

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	252	TYR	2.7
1	B	498	TYR	2.6
1	B	246	LYS	2.6
1	B	492	TYR	2.6
1	B	558	ASP	2.5
1	A	559	ASN	2.4
1	A	492	TYR	2.4
1	B	384	ARG	2.4
1	A	508	LEU	2.4
1	B	349	GLN	2.4
1	A	309	MET	2.4
1	A	557	SER	2.3
1	A	349	GLN	2.3
1	A	558	ASP	2.2
1	A	365	GLU	2.2
1	B	564	GLU	2.1
1	B	396	THR	2.1
1	B	260	GLN	2.1
1	A	427	ASP	2.1
1	A	477	TYR	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	ZN	A	701	1/1	0.97	0.28	20,20,20,20	0
2	ZN	B	701	1/1	0.97	0.32	20,20,20,20	0
2	ZN	A	702	1/1	0.99	0.17	16,16,16,16	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




**Electron density around ZN B 701:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)







## 6.5 Other polymers ⓘ

There are no such residues in this entry.