

Full wwPDB EM Validation Report (i)

Nov 21, 2022 – 11:55 PM JST

PDB ID : 7DRT

EMDB ID : EMD-30827

Title : Human Wntless in complex with Wnt3a

Authors: Zhong, Q.; Zhao, Y.; Ye, F.; Xiao, Z.; Huang, G.; Zhang, Y.; Lu, P.; Xu, W.;

Zhou, Q.; Ma, D.

Deposited on : 2020-12-29

Resolution : 2.20 Å(reported)

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/EMValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev43

Mogul : 1.8.5 (274361), CSD as541be (2020)

MolProbity : 4.02b-467 buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

 $MapQ \quad : \quad 1.9.9$

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.31.3

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $ELECTRON\ MICROSCOPY$

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	${ m EM\ structures} \ (\#{ m Entries})$
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion <40%). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	A	352	16% 82%	11% • 5%
2	В	541	85%	7% 8%
3	Н	2	50% 50%	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	NAG	Н	1	-	-	X	-



2 Entry composition (i)

There are 8 unique types of molecules in this entry. The entry contains 6849 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Protein Wnt-3a.

\mathbf{Mol}	Chain	Residues		At	oms			AltConf	Trace
1	A	334	Total 2618	C 1620	N 490	O 479	S 29	0	0

• Molecule 2 is a protein called Protein wntless homolog.

Mol	Chain	Residues		At	oms			AltConf	Trace
2	В	496	Total 4037	C 2675	N 658	O 667	S 37	0	0


• Molecule 3 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-a cetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	F	A ton	ns		AltConf	Trace
3	Н	2	Total 28	C 16	N 2	O 10	0	0


• Molecule 4 is PALMITOLEIC ACID (three-letter code: PAM) (formula: C₁₆H₃₀O₂).





Mol	Chain	Residues	Atoms	AltConf
4	A	1	Total C O 17 16 1	0


 \bullet Molecule 5 is 1,2-DIOLEOYL-SN-GLYCERO-3-PHOSPHOCHOLINE (three-letter code: PCW) (formula: $\rm C_{44}H_{85}NO_8P).$



\mathbf{M}	ol	Chain	Residues	Atoms	AltConf
63	5	A	1	Total C O P 44 35 8 1	0
63	5	В	1	Total C N O P 41 31 1 8 1	0




 \bullet Molecule 6 is CHOLESTEROL (three-letter code: CLR) (formula: $\mathrm{C_{27}H_{46}O}).$



Mol	Chain	Residues	Atoms	AltConf
6	В	1	Total C O 28 27 1	0

• Molecule 7 is 1-O-OCTADECYL-SN-GLYCERO-3-PHOSPHOCHOLINE (three-letter code: LPE) (formula: $C_{26}H_{57}NO_6P$).



Mol	Chain	Residues		Atoms			AltConf	
7	D	1	Total	С	N	О	Р	0
'	D	1	26	18	1	6	1	U



• Molecule 8 is water.


Mol	Chain	Residues	Atoms	AltConf
8	A	5	Total O 5 5	0
8	В	5	Total O 5 5	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Protein Wnt-3a



• Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-gluc opyranose

Chain H: 50% 50%



4 Experimental information (i)

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	8482332	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE	Depositor
	CORRECTION	
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose $(e^-/\text{Å}^2)$	50	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	Not provided	
Image detector	GATAN K3 BIOQUANTUM (6k x 4k)	Depositor
Maximum map value	9.254	Depositor
Minimum map value	-4.234	Depositor
Average map value	-0.000	Depositor
Map value standard deviation	0.057	Depositor
Recommended contour level	0.2	Depositor
Map size (Å)	278.272, 278.272, 278.272	wwPDB
Map dimensions	512, 512, 512	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	0.5435, 0.5435, 0.5435	Depositor



5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: PCW, CLR, LPE, PAM, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
MIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	A	0.72	0/2686	0.66	0/3629	
2	В	0.64	0/4161	0.60	0/5640	
All	All	0.67	0/6847	0.63	0/9269	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2618	0	2475	41	0
2	В	4037	0	4054	28	0
3	Н	28	0	25	10	0
4	A	17	0	29	5	0
5	A	44	0	60	2	0
5	В	41	0	52	3	0
6	В	28	0	46	0	0
7	В	26	0	37	5	0
8	A	5	0	0	3	0
8	В	5	0	0	7	0
All	All	6849	0	6778	74	0



The all-atom clash score is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clash score for this structure is 6.

All (74) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic	Clash
		distance (Å)	overlap (Å)
1:A:87:ASN:HD21	3:H:1:NAG:C1	1.33	1.39
1:A:87:ASN:ND2	3:H:1:NAG:C1	2.06	1.16
2:B:75:ARG:HB3	8:B:701:HOH:O	1.59	0.99
1:A:87:ASN:HB2	3:H:1:NAG:H82	1.48	0.95
1:A:142:HIS:O	1:A:143:GLN:HG2	1.69	0.93
1:A:19:SER:HA	1:A:102:LEU:HB2	1.51	0.93
1:A:87:ASN:HB2	3:H:1:NAG:C8	2.05	0.85
1:A:87:ASN:CB	3:H:1:NAG:H82	2.06	0.84
1:A:48:LEU:HA	1:A:52:GLN:OE1	1.81	0.81
1:A:143:GLN:HG3	1:A:149:GLY:HA3	1.61	0.79
1:A:87:ASN:CG	3:H:1:NAG:H82	2.04	0.78
2:B:122:LEU:HG	8:B:704:HOH:O	1.83	0.77
2:B:272:ALA:HB2	7:B:602:LPE:H162	1.68	0.74
2:B:120:PHE:CE2	8:B:704:HOH:O	2.39	0.74
2:B:272:ALA:HB1	7:B:602:LPE:H182	1.70	0.74
1:A:295:ARG:HG2	1:A:310:LEU:HD21	1.68	0.74
1:A:142:HIS:O	1:A:143:GLN:CG	2.37	0.72
5:A:502:PCW:H32	2:B:232:LYS:HB3	1.70	0.72
1:A:193:GLN:HG3	1:A:268:GLU:HB3	1.72	0.70
2:B:451:ILE:O	2:B:455:VAL:HG22	1.96	0.66
2:B:478:TYR:CD1	5:B:603:PCW:H2	2.32	0.64
1:A:192:ARG:HG2	8:A:603:HOH:O	1.96	0.64
4:A:501:PAM:H112	4:A:501:PAM:H152	1.80	0.62
4:A:501:PAM:H112	4:A:501:PAM:C15	2.30	0.61
2:B:272:ALA:CB	7:B:602:LPE:H162	2.32	0.59
4:A:501:PAM:H152	4:A:501:PAM:C11	2.33	0.59
2:B:122:LEU:CD2	8:B:704:HOH:O	2.50	0.59
1:A:87:ASN:ND2	3:H:1:NAG:C2	2.65	0.58
1:A:26:LEU:HD11	1:A:45:ILE:HG23	1.86	0.57
2:B:280:ILE:HD11	2:B:307:PHE:HB2	1.86	0.57
1:A:89:THR:HG21	1:A:290:PHE:CD2	2.40	0.56
2:B:360:GLN:NE2	2:B:366:TYR:O	2.38	0.56
1:A:48:LEU:HG	1:A:52:GLN:CB	2.36	0.55
1:A:45:ILE:HB	1:A:48:LEU:HD22	1.89	0.54
2:B:122:LEU:CG	8:B:704:HOH:O	2.50	0.54
1:A:196:ALA:HB3	8:A:604:HOH:O	2.08	0.54
1:A:87:ASN:ND2	3:H:1:NAG:O5	2.28	0.53

Continued on next page...



 $Continued\ from\ previous\ page...$

Continued from prec		Interatomic	Clash
Atom-1	Atom-2	distance (Å)	$\text{overlap } (\mathring{\mathbf{A}})$
1:A:211:SER:HB3	8:A:605:HOH:O	2.11	0.51
1:A:48:LEU:HG	1:A:52:GLN:HB3	1.93	0.50
1:A:84:ARG:NH1	1:A:232:LYS:HE2	2.27	0.50
1:A:84:ARG:HH12	1:A:232:LYS:HE2	1.77	0.50
1:A:87:ASN:ND2	3:H:1:NAG:H82	2.27	0.49
1:A:330:VAL:HG21	2:B:205:GLU:CD	2.33	0.48
2:B:267:GLU:HG2	8:B:702:HOH:O	2.12	0.48
1:A:87:ASN:HB2	3:H:1:NAG:H81	1.91	0.48
1:A:193:GLN:HG3	1:A:268:GLU:CB	2.43	0.48
1:A:23:TRP:HA	1:A:46:PRO:HD2	1.95	0.47
1:A:20:TYR:H	1:A:21:PRO:CD	2.28	0.47
1:A:48:LEU:HG	1:A:52:GLN:HB2	1.97	0.47
2:B:478:TYR:HD1	5:B:603:PCW:H2	1.76	0.46
2:B:242:THR:HB	2:B:243:PRO:HD3	1.98	0.46
4:A:501:PAM:H72	2:B:302:ILE:HG23	1.97	0.46
1:A:30:PRO:O	1:A:31:GLN:C	2.53	0.46
2:B:448:ILE:O	2:B:452:VAL:HG23	2.16	0.46
1:A:142:HIS:O	1:A:143:GLN:NE2	2.49	0.45
1:A:207:GLY:HA3	1:A:213:GLU:HG2	1.98	0.45
1:A:49:VAL:HB	1:A:50:PRO:HD2	1.98	0.44
5:A:502:PCW:H161	5:A:502:PCW:H19	1.73	0.44
1:A:215:LYS:HB3	1:A:215:LYS:HE2	1.47	0.44
1:A:297:CYS:SG	1:A:298:ASN:N	2.90	0.44
2:B:284:VAL:HG21	2:B:299:PHE:HE2	1.83	0.44
2:B:414:LEU:N	2:B:415:PRO:CD	2.80	0.43
5:B:603:PCW:H41	5:B:603:PCW:H63	1.60	0.43
7:B:602:LPE:H2N3	7:B:602:LPE:H311	1.77	0.43
2:B:87:ALA:O	2:B:88:ASN:HB2	2.18	0.43
1:A:84:ARG:O	1:A:85:ARG:HB2	2.19	0.42
2:B:120:PHE:HE2	8:B:704:HOH:O	1.91	0.42
2:B:286:TRP:CD1	2:B:289:ILE:HD11	2.54	0.42
4:A:501:PAM:H52	2:B:305:GLY:HA3	2.02	0.42
2:B:275:ILE:HD12	7:B:602:LPE:H161	2.01	0.42
2:B:81:ILE:HB	2:B:82:PRO:HD3	2.02	0.41
2:B:278:THR:O	2:B:282:ILE:HG13	2.21	0.41
1:A:90:THR:HG22	1:A:97:ILE:HG12	2.02	0.41
1:A:258:PRO:HB2	1:A:260:TYR:CE2	2.56	0.40

There are no symmetry-related clashes.



5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed Favoured Allowed		Allowed	Outliers	Percentiles
1	A	$332/352 \ (94\%)$	299 (90%)	30 (9%)	3 (1%)	17 16
2	В	$494/541 \ (91\%)$	478 (97%)	15 (3%)	1 (0%)	47 55
All	All	826/893 (92%)	777 (94%)	45 (5%)	4 (0%)	32 31

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	20	TYR
1	A	143	GLN
1	A	266	PRO
2	В	39	PRO

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles		
1	A	280/294~(95%)	273 (98%)	7 (2%)	47 60		
2	В	435/473 (92%)	430 (99%)	5 (1%)	73 85		
All	All	715/767 (93%)	703 (98%)	12 (2%)	62 74		

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	45	ILE
1	A	88	CYS

Continued on next page...



Continued from previous page...

Mol	Chain	Res	Type
1	A	215	LYS
1	A	232	LYS
1	A	269	ARG
1	A	295	ARG
1	A	326	LYS
2	В	277	MET
2	В	279	PHE
2	В	280	ILE
2	В	297	LEU
2	В	395	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	87	ASN
1	A	186	HIS
2	В	108	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

2 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

	Mol	Type	Chain	Pog	Link	Bo	ond leng	$ ag{ths}$	В	ond ang	les
		туре	Chain			Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
	3	NAG	Н	1	3	14,14,15	0.65	0	17,19,21	2.64	4 (23%)



Mol	Type	Chain	Res	Link	Bo	ond leng	$ ag{ths}$	В	ond ang	les
MIOI	туре	Chain	am nes	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAG	Н	2	3	14,14,15	0.29	0	17,19,21	0.69	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	Н	1	3	-	3/6/23/26	0/1/1/1
3	NAG	Н	2	3	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^{o})$
3	Н	1	NAG	O5-C1-C2	-8.48	97.89	111.29
3	Н	1	NAG	C1-O5-C5	-4.59	105.97	112.19
3	Н	1	NAG	C2-N2-C7	-2.98	118.65	122.90
3	Н	1	NAG	O4-C4-C3	-2.60	104.33	110.35

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	Н	1	NAG	C8-C7-N2-C2
3	Н	1	NAG	O7-C7-N2-C2
3	Н	1	NAG	C4-C5-C6-O6


There are no ring outliers.

1 monomer is involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	Н	1	NAG	10	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry (i)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Tune	Chain	Res	Link	Bond lengths			Bond angles		
MIOI	Type	Chain	nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	CLR	В	601	-	31,31,31	0.66	0	48,48,48	0.92	1 (2%)
7	LPE	В	602	-	25,25,33	0.55	0	29,31,39	0.71	1 (3%)
4	PAM	A	501	-	16,16,17	0.85	1 (6%)	15,15,17	0.34	0



Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
MIOI	туре	Chain	nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z >2
5	PCW	A	502	-	43,43,53	1.04	2 (4%)	47,48,61	1.13	4 (8%)
5	PCW	В	603	-	40,40,53	1.09	2 (5%)	45,48,61	1.12	3 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	CLR	В	601	-	-	2/10/68/68	0/4/4/4
7	LPE	В	602	-	-	7/26/26/34	-
4	PAM	A	501	-	-	12/14/14/15	-
5	PCW	A	502	-	-	11/45/45/57	-
5	PCW	В	603	-	-	5/44/44/57	-

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\text{\AA})$	Ideal(Å)
5	A	502	PCW	O3-C11	4.43	1.46	1.33
5	A	502	PCW	O2-C31	3.95	1.45	1.34
5	В	603	PCW	O3-C11	3.82	1.44	1.33
5	В	603	PCW	O2-C31	3.72	1.44	1.34
4	A	501	PAM	O1-C1	-3.28	1.25	1.42

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
5	В	603	PCW	O2-C31-C32	3.95	120.02	111.50
5	A	502	PCW	O2-C31-C32	3.80	119.68	111.50
5	A	502	PCW	O3-C11-C12	2.60	120.07	111.91
5	В	603	PCW	O3-C11-C12	2.34	119.25	111.91
5	A	502	PCW	O1P-P-O2P	2.25	119.50	110.68
6	В	601	CLR	C7-C8-C9	2.24	112.43	109.71
5	В	603	PCW	C34-C33-C32	-2.18	105.36	113.19
7	В	602	LPE	C31-C32-N	-2.13	108.66	115.78
5	A	502	PCW	C13-C12-C11	-2.08	106.06	113.62

There are no chirality outliers.

All (37) torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
5	В	603	PCW	C4-O4P-P-O1P
7	В	602	LPE	C3-O3-P-O31
5	A	502	PCW	O11-C11-O3-C3
5	A	502	PCW	C12-C11-O3-C3
4	A	501	PAM	C11-C12-C13-C14
6	В	601	CLR	C17-C20-C22-C23
6	В	601	CLR	C21-C20-C22-C23
5	В	603	PCW	C4-O4P-P-O3P
7	В	602	LPE	C3-O3-P-O33
4	A	501	PAM	O1-C1-C2-C3
4	A	501	PAM	C2-C3-C4-C5
4	A	501	PAM	C5-C6-C7-C8
5	A	502	PCW	C22-C23-C24-C25
4	A	501	PAM	C6-C7-C8-C9
5	A	502	PCW	C33-C34-C35-C36
5	A	502	PCW	O3P-C1-C2-O2
4	A	501	PAM	C1-C2-C3-C4
5	В	603	PCW	C39-C40-C41-C42
5	A	502	PCW	C1-O3P-P-O4P
5	В	603	PCW	C32-C33-C34-C35
7	В	602	LPE	C3-O3-P-O32
7	В	602	LPE	C31-O33-P-O32
5	A	502	PCW	O3P-C1-C2-C3
5	A	502	PCW	C34-C35-C36-C37
7	В	602	LPE	O33-C31-C32-N
4	A	501	PAM	C3-C4-C5-C6
4	A	501	PAM	C11-C10-C9-C8
4	A	501	PAM	C10-C11-C12-C13
5	В	603	PCW	C11-C12-C13-C14
4	A	501	PAM	C4-C5-C6-C7
5	A	502	PCW	C1-O3P-P-O1P
5	A	502	PCW	C17-C18-C19-C20
4	A	501	PAM	C9-C10-C11-C12
5	A	502	PCW	C39-C40-C41-C42
7	В	602	LPE	C31-O33-P-O31
4	A	501	PAM	C13-C14-C15-C16
7	В	602	LPE	O1-C1-C2-O2H

There are no ring outliers.

4 monomers are involved in 15 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	В	602	LPE	5	0


Continued on next page...




Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	501	PAM	5	0
5	A	502	PCW	2	0
5	В	603	PCW	3	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.
















5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.




6 Map visualisation (i)

This section contains visualisations of the EMDB entry EMD-30827. These allow visual inspection of the internal detail of the map and identification of artifacts.

No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections (i)


6.1.1 Primary map



The images above show the map projected in three orthogonal directions.

6.2 Central slices (i)


6.2.1 Primary map







Y Index: 256



Z Index: 256



The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices (i)


6.3.1 Primary map



The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal surface views (i)

6.4.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.2. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.



6.5 Mask visualisation (i)


This section was not generated. No masks/segmentation were deposited.



7 Map analysis (i)

This section contains the results of statistical analysis of the map.


7.1 Map-value distribution (i)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.



7.2 Volume estimate (i)




The volume at the recommended contour level is 52 nm^3 ; this corresponds to an approximate mass of 47 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.



7.3 Rotationally averaged power spectrum (i)



*Reported resolution corresponds to spatial frequency of 0.455 $\rm \mathring{A}^{-1}$



8 Fourier-Shell correlation (i)


This section was not generated. No FSC curve or half-maps provided.



9 Map-model fit (i)

This section contains information regarding the fit between EMDB map EMD-30827 and PDB model 7DRT. Per-residue inclusion information can be found in section 3 on page 7.


9.1 Map-model overlay (i)



The images above show the 3D surface view of the map at the recommended contour level 0.2 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.




9.2 Q-score mapped to coordinate model (i)



The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.


9.3 Atom inclusion mapped to coordinate model (i)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.2).



9.4 Atom inclusion (i)



At the recommended contour level, 80% of all backbone atoms, 80% of all non-hydrogen atoms, are inside the map.



9.5 Map-model fit summary (i)

The table lists the average atom inclusion at the recommended contour level (0.2) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.8014	0.5300
A	0.7095	0.4610
В	0.8658	0.5760
Н	0.6786	0.4520

