



# wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 21, 2020 – 07:46 AM BST

PDB ID : 5DX0  
Title : Crystal structure of CARM1, sinefungin, and H3 peptide (R17)  
Authors : Boriack-Sjodin, P.A.  
Deposited on : 2015-09-23  
Resolution : 2.05 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.13.1  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.13.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1692 (2.04-2.04)
Clashscore	141614	1773 (2.04-2.04)
Ramachandran outliers	138981	1752 (2.04-2.04)
Sidechain outliers	138945	1752 (2.04-2.04)
RSRZ outliers	127900	1672 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	349	 90% 8%
1	B	349	 89% 9%
1	C	349	 90% 8%
1	D	349	 88% 10%
2	F	20	 10% 30% 15% 55%
2	G	20	 40% 10% 50%

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
2	H	20	
2	I	20	

## 2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 11892 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Histone-arginine methyltransferase CARM1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	343	Total 2756	C 1779	N 454	O 509	S 14	0	1	0
1	B	343	Total 2761	C 1783	N 454	O 509	S 15	0	2	0
1	C	343	Total 2751	C 1776	N 454	O 507	S 14	0	0	0
1	D	343	Total 2756	C 1779	N 454	O 509	S 14	0	1	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	131	SER	-	expression tag	UNP Q86X55
A	132	ILE	-	expression tag	UNP Q86X55
A	133	ALA	-	expression tag	UNP Q86X55
B	131	SER	-	expression tag	UNP Q86X55
B	132	ILE	-	expression tag	UNP Q86X55
B	133	ALA	-	expression tag	UNP Q86X55
C	131	SER	-	expression tag	UNP Q86X55
C	132	ILE	-	expression tag	UNP Q86X55
C	133	ALA	-	expression tag	UNP Q86X55
D	131	SER	-	expression tag	UNP Q86X55
D	132	ILE	-	expression tag	UNP Q86X55
D	133	ALA	-	expression tag	UNP Q86X55

- Molecule 2 is a protein called H3 peptide.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	F	9	Total 65	C 41	N 14	O 10	0	0	0
2	G	10	Total 70	C 44	N 15	O 11	0	0	0

*Continued on next page...*


Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	H	9	Total	C	N	O	0	0	0
			65	41	14	10			
2	I	10	Total	C	N	O	0	0	0
			70	44	15	11			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
F	0	ACE	-	acetylation	UNP P84243
F	19	NH2	-	amidation	UNP P84243
G	0	ACE	-	acetylation	UNP P84243
G	19	NH2	-	amidation	UNP P84243
H	0	ACE	-	acetylation	UNP P84243
H	19	NH2	-	amidation	UNP P84243
I	0	ACE	-	acetylation	UNP P84243
I	19	NH2	-	amidation	UNP P84243

- Molecule 3 is SINEFUNGIN (three-letter code: SFG) (formula: C<sub>15</sub>H<sub>23</sub>N<sub>7</sub>O<sub>5</sub>).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			27	15	7	5		
3	B	1	Total	C	N	O	0	0
			27	15	7	5		
3	C	1	Total	C	N	O	0	0
			27	15	7	5		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	
			Total	C	N			O
3	D	1	27	15	7	5	0	0

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
4	A	1	6	3	3	0	0
4	C	1	6	3	3	0	0

- Molecule 5 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total O S 5 4 1	0	0
5	B	1	Total O S 5 4 1	0	0
5	C	1	Total O S 5 4 1	0	0
5	D	1	Total O S 5 4 1	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	94	Total O 94 94	0	0
6	B	120	Total O 120 120	0	0
6	C	109	Total O 109 109	0	0
6	D	113	Total O 113 113	0	0
6	F	2	Total O 2 2	0	0
6	G	8	Total O 8 8	0	0
6	H	3	Total O 3 3	0	0
6	I	9	Total O 9 9	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Histone-arginine methyltransferase CARM1




- Molecule 1: Histone-arginine methyltransferase CARM1




- Molecule 1: Histone-arginine methyltransferase CARM1





- Molecule 1: Histone-arginine methyltransferase CARM1










- Molecule 2: H3 peptide




- Molecule 2: H3 peptide



- Molecule 2: H3 peptide



- Molecule 2: H3 peptide



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	75.72Å 98.85Å 208.67Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 2.05 46.14 – 2.05	Depositor EDS
% Data completeness (in resolution range)	89.8 (50.00-2.05) 89.8 (46.14-2.05)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.27 (at 2.05Å)	Xtrriage
Refinement program	REFMAC 5.8.0073	Depositor
R, $R_{free}$	0.211 , 0.263 0.217 , 0.267	Depositor DCC
$R_{free}$ test set	4324 reflections (4.86%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	30.2	Xtrriage
Anisotropy	0.100	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 36.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	11892	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	34.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 79.70 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 5.2649e-07. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, SO4, ACE, SFG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.52	0/2829	0.70	0/3834
1	B	0.51	0/2837	0.68	0/3844
1	C	0.54	0/2821	0.68	0/3823
1	D	0.52	0/2829	0.68	0/3834
2	F	0.47	0/63	0.78	0/82
2	G	0.52	0/68	0.75	0/89
2	H	0.55	0/63	0.72	0/82
2	I	0.44	0/68	0.64	0/89
All	All	0.52	0/11578	0.69	0/15677

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2756	0	2703	19	0
1	B	2761	0	2712	16	0
1	C	2751	0	2699	16	0
1	D	2756	0	2703	24	0
2	F	65	0	76	4	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	G	70	0	81	2	0
2	H	65	0	76	4	0
2	I	70	0	81	0	0
3	A	27	0	22	2	0
3	B	27	0	22	0	0
3	C	27	0	22	0	0
3	D	27	0	22	0	0
4	A	6	0	8	0	0
4	C	6	0	8	0	0
5	A	5	0	0	0	0
5	B	5	0	0	1	0
5	C	5	0	0	1	0
5	D	5	0	0	1	0
6	A	94	0	0	4	0
6	B	120	0	0	3	0
6	C	109	0	0	1	0
6	D	113	0	0	4	0
6	F	2	0	0	0	0
6	G	8	0	0	0	0
6	H	3	0	0	0	0
6	I	9	0	0	0	0
All	All	11892	0	11235	78	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 78 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:468[A]:ASP:OD1	6:B:601:HOH:O	1.82	0.97
1:C:265:ASN:HD21	2:H:0:ACE:H3	1.42	0.85
1:D:272:TYR:OH	1:D:287:PRO:O	2.09	0.71
1:A:294:LEU:HD12	1:A:376:PHE:HZ	1.55	0.70
1:D:474:PHE:HB3	6:D:603:HOH:O	1.94	0.66

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	342/349 (98%)	329 (96%)	12 (4%)	1 (0%)	41	31
1	B	343/349 (98%)	332 (97%)	10 (3%)	1 (0%)	41	31
1	C	341/349 (98%)	328 (96%)	12 (4%)	1 (0%)	41	31
1	D	342/349 (98%)	328 (96%)	13 (4%)	1 (0%)	41	31
2	F	7/20 (35%)	7 (100%)	0	0	100	100
2	G	8/20 (40%)	8 (100%)	0	0	100	100
2	H	7/20 (35%)	7 (100%)	0	0	100	100
2	I	8/20 (40%)	8 (100%)	0	0	100	100
All	All	1398/1476 (95%)	1347 (96%)	47 (3%)	4 (0%)	41	31

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	258	PRO
1	B	258	PRO
1	D	258	PRO
1	A	258	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	300/303 (99%)	295 (98%)	5 (2%)	60	57

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	301/303 (99%)	297 (99%)	4 (1%)	69	67
1	C	299/303 (99%)	295 (99%)	4 (1%)	69	67
1	D	300/303 (99%)	296 (99%)	4 (1%)	69	67
2	F	6/12 (50%)	6 (100%)	0	100	100
2	G	6/12 (50%)	6 (100%)	0	100	100
2	H	6/12 (50%)	6 (100%)	0	100	100
2	I	6/12 (50%)	6 (100%)	0	100	100
All	All	1224/1260 (97%)	1207 (99%)	17 (1%)	67	65

5 of 17 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	417	GLN
1	B	477	THR
1	D	249	GLU
1	B	257	GLU
1	D	257	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 25 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	164	GLN
1	C	265	ASN
1	D	471	ASN
1	C	178	GLN
1	C	368	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

10 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	SFG	A	501	-	22,29,29	1.13	3 (13%)	18,42,42	1.52	3 (16%)
5	SO4	C	503	-	4,4,4	0.49	0	6,6,6	0.42	0
3	SFG	D	501	-	22,29,29	1.06	1 (4%)	18,42,42	1.58	4 (22%)
5	SO4	D	502	-	4,4,4	0.52	0	6,6,6	0.42	0
5	SO4	B	502	-	4,4,4	0.35	0	6,6,6	0.40	0
4	GOL	C	502	-	5,5,5	0.25	0	5,5,5	0.28	0
5	SO4	A	503	-	4,4,4	0.58	0	6,6,6	0.34	0
4	GOL	A	502	-	5,5,5	0.26	0	5,5,5	0.33	0
3	SFG	C	501	-	22,29,29	0.92	1 (4%)	18,42,42	1.43	3 (16%)
3	SFG	B	501	-	22,29,29	1.09	3 (13%)	18,42,42	1.57	3 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	SFG	A	501	-	-	2/9/33/33	0/3/3/3
3	SFG	D	501	-	-	3/9/33/33	0/3/3/3
4	GOL	C	502	-	-	4/4/4/4	-
4	GOL	A	502	-	-	2/4/4/4	-
3	SFG	C	501	-	-	2/9/33/33	0/3/3/3
3	SFG	B	501	-	-	2/9/33/33	0/3/3/3

The worst 5 of 8 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	D	501	SFG	C2'-C1'	-2.75	1.49	1.53
3	A	501	SFG	O4'-C1'	2.50	1.44	1.41
3	B	501	SFG	C2'-C1'	-2.33	1.50	1.53
3	B	501	SFG	C2-N3	2.32	1.35	1.32
3	A	501	SFG	C5-C4	2.18	1.46	1.40

The worst 5 of 13 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	501	SFG	N3-C2-N1	-4.24	122.06	128.68
3	B	501	SFG	N3-C2-N1	-3.73	122.84	128.68
3	D	501	SFG	N3-C2-N1	-3.59	123.07	128.68
3	C	501	SFG	N3-C2-N1	-3.52	123.17	128.68
3	C	501	SFG	O4'-C1'-C2'	-2.90	102.68	106.93

There are no chirality outliers.

5 of 15 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	501	SFG	NE-CD-CG-CB
3	A	501	SFG	C5'-CD-CG-CB
3	D	501	SFG	NE-CD-CG-CB
3	D	501	SFG	C5'-CD-CG-CB
4	C	502	GOL	C1-C2-C3-O3

There are no ring outliers.


4 monomers are involved in 5 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	501	SFG	2	0
5	C	503	SO4	1	0
5	D	502	SO4	1	0
5	B	502	SO4	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring



in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2	OWAB(Å <sup>2</sup> )	Q < 0.9
1	A	343/349 (98%)	-0.10	5 (1%) 73 76	24, 34, 50, 65	0
1	B	343/349 (98%)	-0.24	2 (0%) 89 91	23, 31, 45, 71	0
1	C	343/349 (98%)	-0.21	4 (1%) 79 81	22, 32, 44, 62	0
1	D	343/349 (98%)	-0.00	5 (1%) 73 76	21, 35, 54, 68	0
2	F	8/20 (40%)	1.23	2 (25%) 0 0	54, 63, 74, 79	0
2	G	9/20 (45%)	0.07	0 100 100	33, 40, 49, 56	0
2	H	8/20 (40%)	1.08	2 (25%) 0 0	47, 57, 61, 68	0
2	I	9/20 (45%)	-0.35	0 100 100	33, 36, 48, 49	0
All	All	1406/1476 (95%)	-0.13	20 (1%) 75 78	21, 33, 51, 79	0

The worst 5 of 20 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	283	GLY	4.3
1	A	143	GLU	4.1
1	D	476	TYR	3.9
1	A	139	GLU	3.4
1	B	476	TYR	2.9

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
5	SO4	C	503	5/5	0.92	0.24	56,58,59,59	0
5	SO4	A	503	5/5	0.92	0.26	53,55,58,60	0
4	GOL	A	502	6/6	0.92	0.10	54,55,56,58	0
4	GOL	C	502	6/6	0.93	0.14	52,54,54,58	0
3	SFG	A	501	27/27	0.94	0.11	28,31,34,34	0
3	SFG	D	501	27/27	0.94	0.13	26,30,33,34	0
5	SO4	B	502	5/5	0.94	0.15	46,50,54,57	0
3	SFG	C	501	27/27	0.94	0.11	25,29,33,34	0
3	SFG	B	501	27/27	0.94	0.11	21,24,26,27	0
5	SO4	D	502	5/5	0.97	0.11	46,46,52,52	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







**Electron density around SFG D 501:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around SFG C 501:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [i](#)

There are no such residues in this entry.