

wwPDB X-ray Structure Validation Summary Report (i)

Oct 7, 2023 – 11:23 PM EDT

PDB ID : 6DXX

Title: Human N-acylethanolamine-hydrolyzing acid amidase (NAAA) in complex

with non-covalent benzothiazole-piperazine inhibitor ARN19702, in presence

of Triton X-100

Authors : Gorelik, A.; Gebai, A.; Illes, K.; Piomelli, D.; Nagar, B.

Deposited on : 2018-07-01

Resolution : 2.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul: 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.35.1

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac : 5.8.0158

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.35.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries},{\rm resolution\ range}(\mathring{\rm A})) \end{array}$
R_{free}	130704	2808 (2.70-2.70)
Clashscore	141614	3122 (2.70-2.70)
Ramachandran outliers	138981	3069 (2.70-2.70)
Sidechain outliers	138945	3069 (2.70-2.70)
RSRZ outliers	127900	2737 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	A	107	77%	14%	9%
1	С	107	78%	13%	9%
1	Е	107	15% 76%	14%	• 9%
2	В	234	94%		



Continued from previous page...

Mol	Chain	Length	Quality	of chain
2	D	234	92%	7% •
2	F	234	87%	11% .
3	G	2	50%	50%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	CL	D	404	-	-	X	-



2 Entry composition (i)

There are 9 unique types of molecules in this entry. The entry contains 16346 atoms, of which 8022 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called N-acylethanolamine-hydrolyzing acid amidase subunit alpha.

Mol	Chain	Residues			Aton	ns			ZeroOcc	AltConf	Trace
1	Λ	97	Total	С	Н	N	О	S	0	0	0
1	A	91	1552	502	780	131	134	5	0	U	
1	С	97	Total	С	Н	N	О	S	0	0	0
1		91	1553	502	781	131	134	5	0	U	
1	E	97	Total	С	Н	N	О	S	0	0	0
1	E	91	1553	502	781	131	134	5		U	

There are 30 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	19	ASP	-	expression tag	UNP Q02083
A	20	ARG	-	expression tag	UNP Q02083
A	21	HIS	-	expression tag	UNP Q02083
A	22	HIS	-	expression tag	UNP Q02083
A	23	HIS	=	expression tag	UNP Q02083
A	24	HIS	-	expression tag	UNP Q02083
A	25	HIS	ı	expression tag	UNP Q02083
A	26	HIS	-	expression tag	UNP Q02083
A	27	LYS	ı	expression tag	UNP Q02083
A	28	LEU	-	expression tag	UNP Q02083
С	19	ASP	-	expression tag	UNP Q02083
С	20	ARG	-	expression tag	UNP Q02083
С	21	HIS	-	expression tag	UNP Q02083
С	22	HIS	-	expression tag	UNP Q02083
С	23	HIS	-	expression tag	UNP Q02083
С	24	HIS	-	expression tag	UNP Q02083
С	25	HIS	-	expression tag	UNP Q02083
С	26	HIS	-	expression tag	UNP Q02083
С	27	LYS	=	expression tag	UNP Q02083
С	28	LEU	-	expression tag	UNP Q02083
Е	19	ASP	=	expression tag	UNP Q02083
Е	20	ARG	ı	expression tag	UNP Q02083
Е	21	HIS	-	expression tag	UNP Q02083



Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
E	22	HIS	-	expression tag	UNP Q02083
E	23	HIS	-	expression tag	UNP Q02083
E	24	HIS	-	expression tag	UNP Q02083
E	25	HIS	-	expression tag	UNP Q02083
E	26	HIS	-	expression tag	UNP Q02083
E	27	LYS	-	expression tag	UNP Q02083
E	28	LEU	-	expression tag	UNP Q02083

• Molecule 2 is a protein called N-acylethanolamine-hydrolyzing acid amidase subunit beta.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	В	231		С		N	О	S	0	0	0
	В	201	3637	1177	1800	323	334	3		Ü	
9	D	231	Total	С	Η	N	O	\mathbf{S}	0	1	0
	D	201	3658	1182	1813	326	334	3		1	
2	F	231	Total	С	Н	N	О	S	0	1	0
2	Г	231	3659	1182	1814	326	334	3		1	U


• Molecule 3 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-a cetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
2	C	9	Total	С	Н	N	О	0	0	0
3	G	2	55	16	27	2	10	0	0	U

• Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).





Mol	Chain	Residues		At	oms			ZeroOcc	AltConf
4	Λ	1	Total	С	Н	N	О	0	0
4	A	1	28	8	14	1	5	0	U
1	В	1	Total	С	Н	N	О	0	0
4	Ъ	1	28	8	14	1	5		U
1	С	1	Total	С	Н	N	О	0	0
$\frac{4}{}$		1	28	8	14	1	5	0	U
1	D	1	Total	С	Н	N	О	0	0
4	ע	1	28	8	14	1	5	0	U
1	E	1	Total	С	Н	N	О	0	0
4	ند	1	28	8	14	1	5		0


• Molecule 5 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total Cl 1 1	0	0
5	В	2	Total Cl 2 2	0	0
5	С	3	Total Cl 3 3	0	0
5	D	4	Total Cl 4 4	0	0
5	E	1	Total Cl 1 1	0	0
5	F	1	Total Cl 1 1	0	0

 $\bullet \ \, \text{Molecule 6 is 2-} \\ \{2\text{-}[4\text{-}(1,1,3,3\text{-}\text{TETRAMETHYLBUTYL})\text{PHENOXY}]\\ \text{ETHOXY}\}\\ \text{ETHANO}\\$




L (three-letter code: TON) (formula: $C_{18}H_{30}O_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	Λ	1	Total	С	Н	О	0	0
0	Λ	1	51	18	30	3	0	U
6	C	1	Total	С	Н	О	0	0
0	C	1	51	18	30	3	0	U
6	E	1	Total	С	Н	О	0	0
0	E	1	51	18	30	3		U


 $\bullet \ \, \text{Molecule 7 is [2-(ethylsulfonyl)phenyl][(2S)-4-(6-fluoro-1,3-benzothiazol-2-yl)-2-methylpiper azin-1-yl]methanone (three-letter code: WTF) (formula: $C_{21}H_{22}FN_3O_3S_2$). }$





Mol	Chain	Residues			Ato	oms				ZeroOcc	AltConf
7	В	1	Total	С	F	Η	N	О	S	0	0
'	Ъ	1	52	21	1	22	3	3	2		0
7	D	1	Total	С	F	Н	N	О	S	0	0
'	ע	1	52	21	1	22	3	3	2	0	0
7	Г	1	Total	С	F	Н	N	О	S	0	0
'	Г	1	52	21	1	22	3	3	2	0	0

• Molecule 8 is THIOCYANATE ION (three-letter code: SCN) (formula: CNS).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
8	С	1	Total	С	N	S	0	0
0		1	3	1	1	1	0	U
Q	D	1	Total	С	N	S	0	0
0	D	1	3	1	1	1	0	U
Q	D	1	Total	С	N	S	0	0
0	D	1	3	1	1	1	U	U
Q	F	1	Total	С	N	S	0	0
0	Г	1	3	1	1	1	0	U
Q	F	1	Total	С	N	S	0	0
0	Г	1	3	1	1	1	0	U
Q	F	1	Total	С	N	S	0	0
	I.	1	3	1	1	1		U

• Molecule 9 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
9	A	21	Total O 21 21	0	0



Continued from previous page...


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
9	В	61	Total O 61 61	0	0
9	С	19	Total O 19 19	0	0
9	D	44	Total O 44 44	0	0
9	E	17	Total O 17 17	0	0
9	F	38	Total O 38 38	0	0




3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: N-acylethanolamine-hydrolyzing acid amidase subunit alpha



• Molecule 2: N-acylethanolamine-hydrolyzing acid amidase subunit beta




• Molecule 2: N-acylethanolamine-hydrolyzing acid amidase subunit beta



• Molecule 2: N-acylethanolamine-hydrolyzing acid amidase subunit beta





 \bullet Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain G: 50% 50%





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	76.61Å 99.80Å 92.76Å	Donositor
a, b, c, α , β , γ	90.00° 96.13° 90.00°	Depositor
Resolution (Å)	46.11 - 2.70	Depositor
Resolution (A)	48.75 - 2.70	EDS
% Data completeness	77.1 (46.11-2.70)	Depositor
(in resolution range)	78.6 (48.75-2.70)	EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	0.98 (at 2.69Å)	Xtriage
Refinement program	PHENIX (1.13_2998)	Depositor
D D.	0.206 , 0.246	Depositor
R, R_{free}	0.206 , 0.246	DCC
R_{free} test set	1944 reflections (5.76%)	wwPDB-VP
Wilson B-factor (Å ²)	42.1	Xtriage
Anisotropy	0.151	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.39, 40.5	EDS
L-test for twinning ²	$ < L > = 0.47, < L^2> = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	16346	wwPDB-VP
Average B, all atoms (Å ²)	52.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 4.83% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, SCN, CL, TON, WTF

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Chain		Boı	nd lengths	Bond	angles
MIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5
1	A	0.31	0/792	0.48	0/1079
1	С	0.36	0/792	0.50	0/1079
1	Е	0.52	3/792 (0.4%)	0.49	0/1079
2	В	0.30	0/1889	0.48	0/2577
2	D	0.29	0/1900	0.48	0/2591
2	F	0.31	0/1900	0.48	0/2591
All	All	0.33	3/8065 (0.0%)	0.48	0/10996

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\operatorname{Observed}(\operatorname{\AA})$	$\operatorname{Ideal}(ext{\AA})$
1	Е	45	GLU	CG-CD	-7.22	1.41	1.51
1	Е	45	GLU	CD-OE1	-6.56	1.18	1.25
1	Е	45	GLU	CD-OE2	-6.42	1.18	1.25

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	772	780	780	13	0
1	С	772	781	781	13	0



Continued from previous page...

Mol	Chain	Non-H		H(added)	Clashes	Symm-Clashes
1	Ε	772	781	781	19	2
2	В	1837	1800	1800	6	0
2	D	1845	1813	1813	11	0
2	F	1845	1814	1814	22	2
3	G	28	27	25	0	0
4	A	14	14	13	0	0
4	В	14	14	13	0	0
4	С	14	14	13	0	0
4	D	14	14	13	0	0
4	Е	14	14	13	0	0
5	A	1	0	0	0	0
5	В	2	0	0	0	0
5	С	3	0	0	1	0
5	D	4	0	0	3	0
5	Ε	1	0	0	0	0
5	F	1	0	0	0	0
6	A	21	30	30	5	0
6	С	21	30	30	3	0
6	Ε	21	30	30	7	0
7	В	30	22	0	0	0
7	D	30	22	0	1	0
7	F	30	22	0	0	0
8	С	3	0	0	0	0
8	D	6	0	0	0	0
8	F	9	0	0	0	0
9	A	21	0	0	2	0
9	В	61	0	0	0	0
9	С	19	0	0	1	0
9	D	44	0	0	2	0
9	Е	17	0	0	1	0
9	F	38	0	0	4	0
All	All	8324	8022	7949	86	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 86 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	$egin{aligned} ext{Clash} \ ext{overlap } (ext{Å}) \end{aligned}$
1:A:66:GLN:NE2	1:A:70:ASP:OD2	2.11	0.84
1:A:78:VAL:HG12	1:A:82:LYS:HE3	1.59	0.83



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:43:VAL:HG12	1:E:45:GLU:OE1	1.86	0.74
2:F:153:ARG:NH2	9:F:502:HOH:O	2.23	0.70
1:E:91:LEU:O	9:E:301:HOH:O	2.10	0.70

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	$egin{aligned} ext{Clash} \ ext{overlap } (ext{Å}) \end{aligned}$
1:E:45:GLU:OE2	2:F:296:LYS:HZ1[2_746]	1.19	0.41
1:E:45:GLU:OE2	2:F:296:LYS:NZ[2_746]	1.97	0.23

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	A	95/107 (89%)	93 (98%)	2 (2%)	0	100	100
1	C	95/107 (89%)	93 (98%)	2 (2%)	0	100	100
1	E	95/107 (89%)	93 (98%)	2 (2%)	0	100	100
2	В	229/234 (98%)	221 (96%)	8 (4%)	0	100	100
2	D	230/234~(98%)	221 (96%)	9 (4%)	0	100	100
2	F	230/234 (98%)	220 (96%)	10 (4%)	0	100	100
All	All	974/1023 (95%)	941 (97%)	33 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.



The Analysed column shows the	number o	of residues	for	which	the	${\rm sidechain}$	conformation	was
analysed, and the total number of	residues.							

Mol	Chain	Analysed	Rotameric	Outliers	Percentile	\mathbf{s}
1	A	86/96 (90%)	84 (98%)	2 (2%)	50 78	
1	С	86/96 (90%)	85 (99%)	1 (1%)	71 88	
1	Е	86/96 (90%)	85 (99%)	1 (1%)	71 88	
2	В	194/197 (98%)	193 (100%)	1 (0%)	88 96	
2	D	195/197 (99%)	194 (100%)	1 (0%)	88 96	
2	F	195/197 (99%)	194 (100%)	1 (0%)	88 96	
All	All	842/879 (96%)	835 (99%)	7 (1%)	81 93	

5 of 7 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	С	77	HIS
2	D	142	ARG
2	F	142	ARG
1	Е	77	HIS
2	В	142	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	66	GLN
2	В	184	GLN
2	F	184	GLN
2	F	213	HIS

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.



5.5 Carbohydrates (i)

2 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Trino	Chain	Dag	T inle	Bo	ond leng	ths	Bond angles			
MIOI	Type	Chain	Res	Link	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2	
3	NAG	G	1	3,1	14,14,15	0.77	1 (7%)	17,19,21	0.50	0	
3	NAG	G	2	3	14,14,15	0.26	0	17,19,21	0.52	0	

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	G	1	3,1	-	2/6/23/26	0/1/1/1
3	NAG	G	2	3	-	0/6/23/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\operatorname{Observed}(\text{\AA})$	$\operatorname{Ideal}(ext{\AA})$
3	G	1	NAG	O5-C1	2.78	1.48	1.43

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	G	1	NAG	C4-C5-C6-O6
3	G	1	NAG	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry (i)

Of 29 ligands modelled in this entry, 12 are monoatomic - leaving 17 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Mol Type	Chain	Res	Link	Bond lengths			Bond angles			
IVIOI	Туре	Chain	nes	Lilik	Counts	RMSZ	# Z >2	Counts	RMSZ	# Z >2	
4	NAG	В	401	2	14,14,15	0.19	0	17,19,21	0.58	0	
4	NAG	Е	201	1	14,14,15	0.41	0	17,19,21	0.47	0	
8	SCN	F	403	-	1,2,2	0.85	0	0,1,1	-	-	



Mol	Type	Chain	Res	Link	В	ond leng	$_{ m gths}$	В	ond ang	gles
WIOI	Type	Chain	nes	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z >2
7	WTF	В	404	-	30,33,33	3.18	17 (56%)	42,49,49	2.99	15 (35%)
4	NAG	С	201	1	14,14,15	0.40	0	17,19,21	0.42	0
8	SCN	D	407	-	1,2,2	0.85	0	0,1,1	-	-
4	NAG	A	203	1	14,14,15	0.17	0	17,19,21	0.40	0
7	WTF	D	408	-	30,33,33	3.30	16 (53%)	42,49,49	3.08	15 (35%)
8	SCN	F	404	-	1,2,2	1.07	0	0,1,1	-	-
6	TON	С	206	-	21,21,21	0.89	1 (4%)	29,29,29	0.67	0
6	TON	Е	203	-	21,21,21	0.89	1 (4%)	29,29,29	0.72	0
8	SCN	D	406	-	1,2,2	0.83	0	0,1,1	-	-
4	NAG	D	401	2	14,14,15	0.85	1 (7%)	17,19,21	0.55	0
6	TON	A	205	-	21,21,21	0.83	1 (4%)	29,29,29	0.75	0
8	SCN	F	402	-	1,2,2	0.92	0	0,1,1	-	-
8	SCN	С	205	-	1,2,2	0.95	0	0,1,1	-	-
7	WTF	F	405		30,33,33	3.30	17 (56%)	42,49,49	2.96	14 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	В	401	2	-	2/6/23/26	0/1/1/1
4	NAG	E	201	1	-	0/6/23/26	0/1/1/1
7	WTF	В	404	-	-	5/19/34/34	0/4/4/4
4	NAG	С	201	1	-	2/6/23/26	0/1/1/1
4	NAG	A	203	1	-	2/6/23/26	0/1/1/1
7	WTF	D	408	-	-	6/19/34/34	0/4/4/4
6	TON	С	206	-	-	6/19/19/19	0/1/1/1
6	TON	Е	203	-	-	1/19/19/19	0/1/1/1
4	NAG	D	401	2	-	0/6/23/26	0/1/1/1
6	TON	A	205	_	_	4/19/19/19	0/1/1/1
7	WTF	F	405	-	-	6/19/34/34	0/4/4/4

The worst 5 of 54 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\operatorname{Observed}(\operatorname{\AA})$	Ideal(A)
7	F	405	WTF	C21-N17	8.96	1.46	1.32
7	D	408	WTF	C21-N17	8.67	1.45	1.32
7	В	404	WTF	C21-N17	8.06	1.44	1.32



Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\text{\AA})$	$\operatorname{Ideal}(ext{\AA})$
7	D	408	WTF	C26-C27	6.71	1.47	1.36
7	В	404	WTF	C26-C27	6.48	1.46	1.36

The worst 5 of 44 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$Ideal(^{o})$
7	D	408	WTF	O05-S03-O04	-14.91	101.01	118.44
7	В	404	WTF	O05-S03-O04	-14.40	101.61	118.44
7	F	405	WTF	O05-S03-O04	-14.14	101.92	118.44
7	D	408	WTF	C02-S03-C06	5.36	114.19	105.55
7	D	408	WTF	C16-C15-N14	5.21	120.15	110.46

There are no chirality outliers.

5 of 34 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
7	В	404	WTF	C01-C02-S03-C06
7	В	404	WTF	S25-C21-N17-C16
7	В	404	WTF	S25-C21-N17-C18
7	D	408	WTF	C01-C02-S03-O04
7	D	408	WTF	C01-C02-S03-O05

There are no ring outliers.


4 monomers are involved in 16 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	D	408	WTF	1	0
6	С	206	TON	3	0
6	Е	203	TON	7	0
6	A	205	TON	5	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient





equivalents in the CSD to analyse the geometry.
















5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	$\#\mathrm{RSRZ}{>}2$	$\mathbf{OWAB}(\mathrm{\AA}^2)$	Q < 0.9
1	A	97/107 (90%)	0.42	5 (5%) 27 25	31, 44, 95, 120	0
1	С	97/107 (90%)	0.68	15 (15%) 2 1	31, 49, 111, 124	0
1	E	97/107 (90%)	0.83	16 (16%) 1 1	37, 56, 103, 115	0
2	В	231/234 (98%)	-0.00	4 (1%) 70 72	24, 35, 61, 84	0
2	D	231/234 (98%)	0.16	10 (4%) 35 33	27, 41, 66, 96	0
2	F	231/234 (98%)	0.26	9 (3%) 39 38	28, 43, 67, 107	0
All	All	984/1023 (96%)	0.29	59 (5%) 21 20	24, 42, 86, 124	0

The worst 5 of 59 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	С	87	LEU	10.5
1	С	90	PHE	7.8
1	A	87	LEU	6.3
1	С	86	GLU	5.9
2	F	297	GLU	5.5

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.


6.3 Carbohydrates (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.



Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q<0.9
3	NAG	G	2	14/15	0.69	0.36	90,111,133,137	0
3	NAG	G	1	14/15	0.92	0.26	48,66,82,91	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q<0.9
5	CL	С	202	1/1	0.78	0.16	69,69,69,69	0



Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q < 0.9
8	SCN	С	205	3/3	0.80	0.31	63,63,70,95	0
5	CL	С	204	1/1	0.81	0.13	66,66,66,66	0
4	NAG	Е	201	14/15	0.83	0.28	60,72,85,88	0
4	NAG	В	401	14/15	0.84	0.17	40,50,76,76	0
8	SCN	D	407	3/3	0.84	0.14	62,62,66,69	0
8	SCN	D	406	3/3	0.86	0.30	52,52,56,58	0
6	TON	С	206	21/21	0.87	0.54	54,73,93,110	0
4	NAG	A	203	14/15	0.88	0.23	59,73,87,94	0
6	TON	Е	203	21/21	0.90	0.28	44,58,82,92	0
5	CL	D	402	1/1	0.90	0.14	51,51,51,51	0
5	CL	D	404	1/1	0.90	0.50	64,64,64,64	0
4	NAG	D	401	14/15	0.90	0.15	47,61,76,81	0
8	SCN	F	404	3/3	0.90	0.31	62,62,63,72	0
5	CL	В	402	1/1	0.92	0.10	37,37,37,37	0
5	CL	D	403	1/1	0.92	0.08	61,61,61,61	0
5	CL	С	203	1/1	0.92	0.10	38,38,38,38	0
5	CL	D	405	1/1	0.92	0.19	68,68,68,68	0
5	CL	В	403	1/1	0.92	0.07	50,50,50,50	0
5	CL	A	204	1/1	0.93	0.16	50,50,50,50	0
8	SCN	F	403	3/3	0.93	0.27	41,41,48,56	0
6	TON	A	205	21/21	0.93	0.20	38,51,90,96	0
4	NAG	С	201	14/15	0.94	0.17	42,56,80,80	0
7	WTF	F	405	30/30	0.95	0.22	31,44,56,62	0
7	WTF	D	408	30/30	0.95	0.21	33,44,57,60	0
5	CL	Е	202	1/1	0.96	0.17	49,49,49,49	0
7	WTF	В	404	30/30	0.96	0.21	24,37,52,62	0
5	CL	F	401	1/1	0.96	0.13	46,46,46,46	0
8	SCN	F	402	3/3	0.97	0.20	47,47,52,52	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around TON C 206: 2mF_o-DF_c (at 0.7 rmsd) in gray mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)



Electron density around TON A 205: 2mF_o-DF_c (at 0.7 rmsd) in gray mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)

Electron density around WTF F 405: 2mF_o-DF_c (at 0.7 rmsd) in gray mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)







6.5 Other polymers (i)

There are no such residues in this entry.

