



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 21, 2020 – 05:55 PM BST

PDB ID : 5DY9
Title : Y68T Hfq from *Methanococcus jannaschii* in complex with AMP
Authors : Nikulin, A.D.; Mikhailina, A.O.; Lekontseva, N.V.; Balobanov, V.A.;
Nikonova, E.Y.; Tishchenko, S.V.
Deposited on : 2015-09-24
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.13.1
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3398 (1.60-1.60)
Clashscore	141614	3665 (1.60-1.60)
Ramachandran outliers	138981	3564 (1.60-1.60)
Sidechain outliers	138945	3563 (1.60-1.60)
RSRZ outliers	127900	3321 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	71	
1	B	71	
1	C	71	
1	D	71	
1	E	71	
1	F	71	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	G	71	<p>% 76% 11% 21%</p>
1	H	71	<p>% 69% 11% 20%</p>
1	I	71	<p>% 80% 11% 17%</p>
1	J	71	<p>% 76% 11% 20%</p>
1	K	71	<p>% 75% 6% 20%</p>
1	L	71	<p>% 76% 11% 20%</p>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	CL	H	102	-	-	X	-
8	PEG	I	101	-	-	X	-

2 Entry composition i

There are 9 unique types of molecules in this entry. The entry contains 6910 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Hfq-like protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	56	471	301	83	86	1	0	2	0
1	B	58	478	307	83	87	1	0	1	0
1	C	57	465	298	81	85	1	0	0	0
1	D	56	458	293	80	84	1	0	0	0
1	E	60	498	323	85	89	1	0	1	0
1	F	58	473	304	82	86	1	0	0	0
1	G	56	455	292	79	83	1	0	0	0
1	H	57	505	323	90	91	1	0	4	0
1	I	59	488	313	85	89	1	0	1	0
1	J	57	478	305	83	89	1	0	2	0
1	K	57	473	302	83	87	1	0	1	0
1	L	57	473	303	84	85	1	0	1	0

There are 12 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	68	THR	TYR	engineered mutation	UNP Q58830
B	68	THR	TYR	engineered mutation	UNP Q58830
C	68	THR	TYR	engineered mutation	UNP Q58830
D	68	THR	TYR	engineered mutation	UNP Q58830
E	68	THR	TYR	engineered mutation	UNP Q58830

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
F	68	THR	TYR	engineered mutation	UNP Q58830
G	68	THR	TYR	engineered mutation	UNP Q58830
H	68	THR	TYR	engineered mutation	UNP Q58830
I	68	THR	TYR	engineered mutation	UNP Q58830
J	68	THR	TYR	engineered mutation	UNP Q58830
K	68	THR	TYR	engineered mutation	UNP Q58830
L	68	THR	TYR	engineered mutation	UNP Q58830

- Molecule 2 is ADENOSINE MONOPHOSPHATE (three-letter code: AMP) (formula: $C_{10}H_{14}N_5O_7P$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	23	10	5	7	1	0	0
2	E	1	23	10	5	7	1	0	0
2	G	1	23	10	5	7	1	0	0

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	D	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	C	1	Total	O	S	0	0
			5	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	E	1	Total	O	S	0	0
			5	4	1		
4	F	1	Total	O	S	0	0
			5	4	1		
4	J	1	Total	O	S	0	0
			5	4	1		

- Molecule 5 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (three-letter code: TRS) (formula: C₄H₁₂NO₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	E	1	Total	C	N	O	0	0
			8	4	1	3		
5	J	1	Total	C	N	O	0	0
			8	4	1	3		
5	L	1	Total	C	N	O	0	0
			8	4	1	3		

- Molecule 6 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	H	1	Total	Cl	0	0
			1	1		
6	I	1	Total	Cl	0	0
			1	1		
6	K	2	Total	Cl	0	0
			2	2		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	E	1	Total Cl 1 1	0	0

- Molecule 7 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	K	1	Total Na 1 1	0	0
7	F	1	Total Na 1 1	0	0

- Molecule 8 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C₄H₁₀O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	G	1	Total C O 7 4 3	0	0
8	H	1	Total C O 7 4 3	0	0
8	I	1	Total C O 7 4 3	0	0

- Molecule 9 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
9	A	86	Total O 86 86	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
9	B	82	Total O 82 82	0	0
9	C	83	Total O 83 83	0	0
9	D	85	Total O 85 85	0	0
9	E	97	Total O 97 97	0	0
9	F	92	Total O 92 92	0	0
9	G	71	Total O 71 71	0	0
9	H	88	Total O 88 88	0	0
9	I	84	Total O 84 84	0	0
9	J	98	Total O 98 98	0	0
9	K	80	Total O 80 80	0	0
9	L	90	Total O 90 90	0	0



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: Hfq-like protein




- Molecule 1: Hfq-like protein





- Molecule 1: Hfq-like protein





- Molecule 1: Hfq-like protein





- Molecule 1: Hfq-like protein



- Molecule 1: Hfq-like protein





- Molecule 1: Hfq-like protein




- Molecule 1: Hfq-like protein





- Molecule 1: Hfq-like protein





- Molecule 1: Hfq-like protein



- Molecule 1: Hfq-like protein



- Molecule 1: Hfq-like protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	60.64Å 67.64Å 91.17Å 90.00° 90.95° 90.00°	Depositor
Resolution (Å)	50.00 – 1.60 45.58 – 1.60	Depositor EDS
% Data completeness (in resolution range)	99.2 (50.00-1.60) 99.2 (45.58-1.60)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.05	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.90 (at 1.60Å)	Xtrriage
Refinement program	PHENIX 1.9_1692	Depositor
R, R_{free}	0.169 , 0.201 0.171 , 0.201	Depositor DCC
R_{free} test set	2100 reflections (2.17%)	wwPDB-VP
Wilson B-factor (Å ²)	19.0	Xtrriage
Anisotropy	0.272	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 44.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.022 for h,-k,-l	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	6910	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.00% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, CL, NA, SO4, AMP, TRS, PEG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.37	0/483	0.54	0/646
1	B	0.34	0/488	0.54	0/655
1	C	0.35	0/472	0.53	0/632
1	D	0.33	0/464	0.53	0/621
1	E	0.35	0/509	0.58	0/683
1	F	0.37	0/480	0.53	0/644
1	G	0.34	0/461	0.53	0/617
1	H	0.38	0/516	0.58	0/690
1	I	0.35	0/495	0.54	0/665
1	J	0.32	0/488	0.55	0/654
1	K	0.36	0/480	0.53	0/643
1	L	0.36	0/483	0.55	0/646
All	All	0.35	0/5819	0.55	0/7796

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	471	0	483	4	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	478	0	490	3	0
1	C	465	0	474	1	0
1	D	458	0	466	5	0
1	E	498	0	515	3	0
1	F	473	0	484	6	0
1	G	455	0	462	2	0
1	H	505	0	512	8	0
1	I	488	0	498	2	0
1	J	478	0	483	7	0
1	K	473	0	479	3	0
1	L	473	0	487	2	0
2	A	23	0	12	2	0
2	E	23	0	12	3	0
2	G	23	0	12	1	0
3	B	12	0	16	0	0
3	D	6	0	8	0	0
4	C	5	0	0	1	0
4	E	5	0	0	0	0
4	F	5	0	0	0	0
4	J	5	0	0	0	0
5	E	8	0	12	1	0
5	J	8	0	12	2	0
5	L	8	0	12	1	0
6	E	1	0	0	0	0
6	H	1	0	0	2	0
6	I	1	0	0	0	0
6	K	2	0	0	1	0
7	F	1	0	0	0	0
7	K	1	0	0	0	0
8	G	7	0	10	1	0
8	H	7	0	10	2	0
8	I	7	0	10	4	0
9	A	86	0	0	2	0
9	B	82	0	0	3	2
9	C	83	0	0	2	2
9	D	85	0	0	3	2
9	E	97	0	0	6	0
9	F	92	0	0	5	0
9	G	71	0	0	3	0
9	H	88	0	0	5	0
9	I	84	0	0	2	0
9	J	98	0	0	4	1

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
9	K	80	0	0	1	1
9	L	90	0	0	1	0
All	All	6910	0	5959	55	4

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (55) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
9:J:238:HOH:O	6:K:102:CL:CL	2.23	0.94
1:J:21:ARG:NH2	9:J:201:HOH:O	2.03	0.90
1:J:21:ARG:NH1	9:J:202:HOH:O	2.06	0.87
1:K:36:GLU:OE2	9:K:201:HOH:O	1.95	0.84
8:I:101:PEG:O1	9:I:201:HOH:O	1.97	0.81
2:E:101:AMP:O5'	9:E:201:HOH:O	2.00	0.80
4:C:101:SO4:O4	9:C:201:HOH:O	1.99	0.79
1:A:16:ASN:OD1	1:F:47:ASN:ND2	2.19	0.76
1:F:14:ILE:N	9:F:202:HOH:O	2.20	0.73
2:A:101:AMP:H5'2	9:A:201:HOH:O	1.88	0.72
1:B:36:GLU:OE1	9:B:201:HOH:O	2.07	0.71
1:H:27:LYS:HG3	9:H:202:HOH:O	1.90	0.71
1:D:18:GLU:O	9:D:201:HOH:O	2.07	0.71
1:D:56:ASP:OD1	9:D:202:HOH:O	2.08	0.70
1:A:18:GLU:HG2	1:A:21:ARG:HE	1.57	0.69
1:F:18:GLU:OE1	9:F:201:HOH:O	2.11	0.68
1:H:70:GLU:OE2	9:H:201:HOH:O	2.11	0.67
2:E:101:AMP:O3P	9:E:202:HOH:O	2.12	0.67
1:B:14:ILE:HD11	9:B:259:HOH:O	1.95	0.66
2:G:101:AMP:H5'2	9:G:202:HOH:O	1.97	0.64
1:E:70:GLU:OE2	9:E:203:HOH:O	2.16	0.63
1:F:47:ASN:OD1	9:F:201:HOH:O	2.16	0.59
1:F:27:LYS:NZ	9:F:204:HOH:O	2.34	0.59
8:G:102:PEG:H41	9:G:249:HOH:O	2.05	0.56
1:E:48[B]:TYR:HE1	9:E:202:HOH:O	1.88	0.54
1:G:31:PHE:CG	5:L:101:TRS:H31	2.43	0.54
1:G:21:ARG:NH2	9:G:204:HOH:O	2.37	0.53
1:H:21[A]:ARG:NH1	9:H:206:HOH:O	2.41	0.52
1:H:48[B]:TYR:HE1	9:I:208:HOH:O	1.92	0.52
6:H:102:CL:CL	9:H:203:HOH:O	2.56	0.52
8:I:101:PEG:C1	1:J:22:ARG:HH11	2.24	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:32:LEU:CD1	5:E:102:TRS:HN1	2.24	0.50
1:J:31:PHE:CG	5:J:101:TRS:H11	2.46	0.50
1:H:46:SER:HA	8:H:101:PEG:H42	1.94	0.49
1:D:22:ARG:HG3	9:D:201:HOH:O	2.13	0.48
1:L:54:VAL:O	1:L:57[B]:ARG:HG2	2.13	0.48
1:J:31:PHE:CD2	5:J:101:TRS:H11	2.48	0.47
1:C:39:ASP:OD1	9:C:202:HOH:O	2.20	0.47
1:H:21[B]:ARG:NH1	1:H:47:ASN:OD1	2.48	0.47
1:D:23:LEU:HA	1:D:26:LYS:HD2	1.97	0.46
1:A:22:ARG:HB2	9:A:219:HOH:O	2.14	0.46
1:L:70:GLU:OE2	9:L:201:HOH:O	2.21	0.45
1:F:15:PRO:HG2	9:F:201:HOH:O	2.16	0.45
1:B:57:ARG:NH2	9:B:203:HOH:O	2.27	0.44
1:I:14:ILE:HD11	9:J:219:HOH:O	2.17	0.44
8:H:101:PEG:H32	1:I:17:PHE:CD2	2.53	0.43
1:K:40:ALA:HB1	1:K:52:VAL:HB	2.01	0.42
1:H:39:ASP:OD1	9:H:202:HOH:O	2.21	0.41
2:E:101:AMP:O2P	9:E:204:HOH:O	2.21	0.41
8:I:101:PEG:H12	1:J:22:ARG:HH11	1.86	0.41
1:A:18:GLU:OE1	2:A:101:AMP:H2	2.04	0.40
1:H:57[A]:ARG:NH2	6:H:102:CL:CL	2.86	0.40
8:I:101:PEG:H11	1:J:22:ARG:HH11	1.86	0.40
1:K:53:LYS:HE3	1:K:53:LYS:HB3	1.86	0.40
1:E:48[B]:TYR:CE1	9:E:202:HOH:O	2.57	0.40

All (4) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
9:B:266:HOH:O	9:D:268:HOH:O[2_547]	1.67	0.53
9:B:233:HOH:O	9:D:269:HOH:O[2_547]	1.91	0.29
9:C:269:HOH:O	9:K:278:HOH:O[1_554]	1.96	0.24
9:C:270:HOH:O	9:J:257:HOH:O[1_554]	2.13	0.07

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	56/71 (79%)	56 (100%)	0	0	100	100
1	B	57/71 (80%)	56 (98%)	1 (2%)	0	100	100
1	C	55/71 (78%)	55 (100%)	0	0	100	100
1	D	54/71 (76%)	53 (98%)	1 (2%)	0	100	100
1	E	59/71 (83%)	59 (100%)	0	0	100	100
1	F	56/71 (79%)	56 (100%)	0	0	100	100
1	G	54/71 (76%)	53 (98%)	1 (2%)	0	100	100
1	H	59/71 (83%)	58 (98%)	1 (2%)	0	100	100
1	I	58/71 (82%)	57 (98%)	1 (2%)	0	100	100
1	J	57/71 (80%)	56 (98%)	1 (2%)	0	100	100
1	K	56/71 (79%)	56 (100%)	0	0	100	100
1	L	56/71 (79%)	56 (100%)	0	0	100	100
All	All	677/852 (80%)	671 (99%)	6 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	51/64 (80%)	51 (100%)	0	100	100
1	B	52/64 (81%)	52 (100%)	0	100	100
1	C	50/64 (78%)	50 (100%)	0	100	100
1	D	49/64 (77%)	49 (100%)	0	100	100
1	E	54/64 (84%)	54 (100%)	0	100	100
1	F	51/64 (80%)	51 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	G	48/64 (75%)	48 (100%)	0	100	100
1	H	54/64 (84%)	54 (100%)	0	100	100
1	I	53/64 (83%)	53 (100%)	0	100	100
1	J	52/64 (81%)	52 (100%)	0	100	100
1	K	51/64 (80%)	51 (100%)	0	100	100
1	L	51/64 (80%)	51 (100%)	0	100	100
All	All	616/768 (80%)	616 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 23 ligands modelled in this entry, 7 are monoatomic - leaving 16 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	GOL	B	102	-	5,5,5	0.30	0	5,5,5	0.45	0
2	AMP	G	101	-	22,25,25	0.90	1 (4%)	25,38,38	1.13	2 (8%)
8	PEG	H	101	-	6,6,6	0.47	0	5,5,5	0.36	0
4	SO4	J	102	-	4,4,4	0.14	0	6,6,6	0.08	0
5	TRS	L	101	-	7,7,7	0.30	0	9,9,9	0.27	0
4	SO4	E	103	-	4,4,4	0.13	0	6,6,6	0.13	0
2	AMP	E	101	-	22,25,25	0.88	1 (4%)	25,38,38	1.21	2 (8%)
3	GOL	D	101	-	5,5,5	0.35	0	5,5,5	0.29	0
8	PEG	I	101	-	6,6,6	0.41	0	5,5,5	0.49	0
5	TRS	J	101	-	7,7,7	0.30	0	9,9,9	0.38	0
2	AMP	A	101	-	22,25,25	0.90	1 (4%)	25,38,38	1.13	2 (8%)
4	SO4	F	101	-	4,4,4	0.17	0	6,6,6	0.12	0
3	GOL	B	101	-	5,5,5	0.33	0	5,5,5	0.27	0
8	PEG	G	102	-	6,6,6	0.45	0	5,5,5	0.51	0
4	SO4	C	101	-	4,4,4	0.10	0	6,6,6	0.12	0
5	TRS	E	102	-	7,7,7	0.59	0	9,9,9	1.34	1 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	AMP	A	101	-	-	0/6/26/26	0/3/3/3
2	AMP	G	101	-	-	0/6/26/26	0/3/3/3
8	PEG	H	101	-	-	3/4/4/4	-
2	AMP	E	101	-	-	2/6/26/26	0/3/3/3
3	GOL	D	101	-	-	1/4/4/4	-
5	TRS	J	101	-	-	5/9/9/9	-
3	GOL	B	102	-	-	2/4/4/4	-
5	TRS	L	101	-	-	3/9/9/9	-
3	GOL	B	101	-	-	2/4/4/4	-
8	PEG	G	102	-	-	2/4/4/4	-
8	PEG	I	101	-	-	4/4/4/4	-
5	TRS	E	102	-	-	3/9/9/9	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	G	101	AMP	C5-C4	2.63	1.47	1.40
2	A	101	AMP	C5-C4	2.54	1.47	1.40
2	E	101	AMP	C5-C4	2.44	1.47	1.40

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	101	AMP	N3-C2-N1	-3.14	123.77	128.68
5	E	102	TRS	O1-C1-C	-2.94	101.68	111.00
2	G	101	AMP	C4-C5-N7	-2.84	106.44	109.40
2	G	101	AMP	N3-C2-N1	-2.69	124.47	128.68
2	A	101	AMP	N3-C2-N1	-2.58	124.64	128.68
2	A	101	AMP	C4-C5-N7	-2.55	106.74	109.40
2	E	101	AMP	C4-C5-N7	-2.50	106.80	109.40

There are no chirality outliers.

All (27) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	J	101	TRS	N-C-C1-O1
5	E	102	TRS	C2-C-C1-O1
5	E	102	TRS	C3-C-C1-O1
5	E	102	TRS	N-C-C1-O1
3	B	101	GOL	O1-C1-C2-C3
2	E	101	AMP	O4'-C4'-C5'-O5'
3	B	101	GOL	O1-C1-C2-O2
8	H	101	PEG	O2-C3-C4-O4
3	B	102	GOL	O1-C1-C2-C3
3	D	101	GOL	O1-C1-C2-C3
8	H	101	PEG	O1-C1-C2-O2
3	B	102	GOL	O1-C1-C2-O2
5	L	101	TRS	C3-C-C2-O2
8	G	102	PEG	O2-C3-C4-O4
5	J	101	TRS	C1-C-C3-O3
5	J	101	TRS	N-C-C3-O3
5	L	101	TRS	C1-C-C2-O2
5	L	101	TRS	N-C-C2-O2
8	I	101	PEG	C1-C2-O2-C3
2	E	101	AMP	C3'-C4'-C5'-O5'
5	J	101	TRS	C2-C-C1-O1
5	J	101	TRS	C2-C-C3-O3
8	H	101	PEG	C4-C3-O2-C2

Continued on next page...

Continued from previous page...



Mol	Chain	Res	Type	Atoms
8	I	101	PEG	O1-C1-C2-O2
8	G	102	PEG	C1-C2-O2-C3
8	I	101	PEG	C4-C3-O2-C2
8	I	101	PEG	O2-C3-C4-O4


There are no ring outliers.

10 monomers are involved in 18 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	G	101	AMP	1	0
8	H	101	PEG	2	0
5	L	101	TRS	1	0
2	E	101	AMP	3	0
8	I	101	PEG	4	0
5	J	101	TRS	2	0
2	A	101	AMP	2	0
8	G	102	PEG	1	0
4	C	101	SO4	1	0
5	E	102	TRS	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	56/71 (78%)	-0.33	1 (1%) 68 67	13, 21, 35, 49	0
1	B	58/71 (81%)	-0.36	1 (1%) 70 69	14, 21, 33, 43	0
1	C	57/71 (80%)	-0.37	1 (1%) 68 67	14, 20, 30, 43	0
1	D	56/71 (78%)	-0.36	2 (3%) 42 40	14, 22, 39, 58	0
1	E	60/71 (84%)	-0.55	1 (1%) 70 69	12, 20, 33, 44	0
1	F	58/71 (81%)	-0.36	1 (1%) 70 69	13, 19, 34, 45	0
1	G	56/71 (78%)	-0.13	1 (1%) 68 67	14, 24, 41, 50	0
1	H	57/71 (80%)	-0.35	1 (1%) 68 67	12, 18, 32, 51	0
1	I	59/71 (83%)	-0.33	1 (1%) 70 69	13, 18, 33, 49	0
1	J	57/71 (80%)	-0.36	0 100 100	14, 24, 34, 53	0
1	K	57/71 (80%)	-0.31	1 (1%) 68 67	14, 22, 36, 49	0
1	L	57/71 (80%)	-0.47	1 (1%) 68 67	13, 21, 31, 44	0
All	All	688/852 (80%)	-0.36	12 (1%) 70 69	12, 21, 37, 58	0

All (12) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	I	13	VAL	4.8
1	G	17	PHE	3.7
1	B	14	ILE	3.6
1	K	15	PRO	2.9
1	L	15	PRO	2.8
1	D	17	PHE	2.7
1	D	16	ASN	2.7
1	F	14	ILE	2.7
1	C	15	PRO	2.5
1	H	48[A]	TYR	2.2
1	A	16	ASN	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	E	48[A]	TYR	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	AMP	A	101	23/23	0.65	0.26	25,34,40,45	23
6	CL	K	102	1/1	0.67	0.09	53,53,53,53	0
6	CL	K	101	1/1	0.70	0.07	69,69,69,69	0
2	AMP	E	101	23/23	0.70	0.21	23,28,40,44	23
8	PEG	G	102	7/7	0.70	0.16	28,33,35,46	0
8	PEG	I	101	7/7	0.76	0.23	31,37,45,45	0
3	GOL	B	101	6/6	0.78	0.14	41,43,48,50	0
8	PEG	H	101	7/7	0.80	0.18	26,35,44,44	0
7	NA	K	103	1/1	0.83	0.10	48,48,48,48	0
2	AMP	G	101	23/23	0.83	0.15	21,26,33,35	23
5	TRS	L	101	8/8	0.85	0.20	25,36,40,42	0
4	SO4	C	101	5/5	0.86	0.12	28,34,43,48	5
5	TRS	E	102	8/8	0.87	0.22	27,32,34,41	0
3	GOL	B	102	6/6	0.89	0.13	33,39,41,41	0
3	GOL	D	101	6/6	0.92	0.11	28,35,38,46	0
7	NA	F	102	1/1	0.92	0.07	35,35,35,35	1
5	TRS	J	101	8/8	0.94	0.16	25,35,38,50	0
6	CL	H	102	1/1	0.94	0.09	40,40,40,40	0
4	SO4	E	103	5/5	0.94	0.09	32,35,39,40	5
4	SO4	J	102	5/5	0.98	0.15	39,42,45,55	0
4	SO4	F	101	5/5	0.98	0.11	33,36,39,43	0
6	CL	E	104	1/1	0.99	0.04	20,20,20,20	1


Continued on next page...


Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	CL	I	102	1/1	0.99	0.08	16,16,16,16	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







6.5 Other polymers [i](#)

There are no such residues in this entry.