

# wwPDB X-ray Structure Validation Summary Report (i)

May 29, 2020 – 06:47 pm BST

PDB ID : 3E4O

Title: Crystal structure of succinate bound state DctB

Authors: Zhou, Y.F.; Nan, J.; Nan, B.Y.; Liang, Y.H.; Panjikar, S.; Su, X.D.

Deposited on : 2008-08-12

Resolution : 2.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13 EDS : 2.11

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac: 5.8.0158

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001)

Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.11

## 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$ 

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \text{Whole archive} \\ (\#\text{Entries}) \end{array}$	$\begin{array}{c} {\rm Similar \; resolution} \\ (\#{\rm Entries, \; resolution \; range(\AA)}) \end{array}$
$R_{free}$	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain				
1	A	305	70%	12%	•	17%	
1	В	305	66%	13%		17%	



# 2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 4107 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called C4-dicarboxylate transport sensor protein dctB.

$\mathbf{Mol}$	Chain	Residues	${f Atoms}$			ZeroOcc	AltConf	Trace		
1	A	254	Total 1945	C 1232	N 349	O 360	Se 4	0	0	0
1	В	254	Total 1945	C 1232	N 349	O 360	Se 4	0	0	0

There are 72 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	8	MET	-	EXPRESSION TAG	UNP P13633
A	9	GLY	_	EXPRESSION TAG	UNP P13633
A	10	SER	_	EXPRESSION TAG	UNP P13633
A	11	SER	-	EXPRESSION TAG	UNP P13633
A	12	HIS	-	EXPRESSION TAG	UNP P13633
A	13	HIS	-	EXPRESSION TAG	UNP P13633
A	14	HIS	-	EXPRESSION TAG	UNP P13633
A	15	HIS	-	EXPRESSION TAG	UNP P13633
A	16	HIS	-	EXPRESSION TAG	UNP P13633
A	17	HIS	-	EXPRESSION TAG	UNP P13633
A	18	SER	-	EXPRESSION TAG	UNP P13633
A	19	SER	-	EXPRESSION TAG	UNP P13633
A	20	GLY	-	EXPRESSION TAG	UNP P13633
A	21	LEU	-	EXPRESSION TAG	UNP P13633
A	22	VAL	_	EXPRESSION TAG	UNP P13633
A	23	PRO	_	EXPRESSION TAG	UNP P13633
A	24	ARG	-	EXPRESSION TAG	UNP P13633
A	25	GLY	_	EXPRESSION TAG	UNP P13633
A	26	SER	_	EXPRESSION TAG	UNP P13633
A	27	HIS	-	EXPRESSION TAG	UNP P13633
A	28	MSE	-	EXPRESSION TAG	UNP P13633
A	29	ALA	-	EXPRESSION TAG	UNP P13633
A	30	SER	-	EXPRESSION TAG	UNP P13633
A	31	MSE	-	EXPRESSION TAG	UNP P13633
A	32	THR	_	EXPRESSION TAG	UNP P13633

Continued on next page...



Continued from previous page...

Chain	Residue	Modelled  Modelled	Actual	Comment	Reference
A	33	GLY		EXPRESSION TAG	UNP P13633
A	34	GLY	-	EXPRESSION TAG	UNP P13633
A	35	GLU	-	EXPRESSION TAG	UNP P13633
A	36	GLU	-	EXPRESSION TAG	UNP P13633
A	37	MSE	_	EXPRESSION TAG	UNP P13633
A	38	GLY	-	EXPRESSION TAG	UNP P13633
A	39	ARG	-	EXPRESSION TAG	UNP P13633
A	40	GLY	_	EXPRESSION TAG	UNP P13633
A	41	SER	-	EXPRESSION TAG	UNP P13633
A	174	LYS	ASN	SEE REMARK 999	UNP P13633
A	309	ASN	LYS	SEE REMARK 999	UNP P13633
В	8	MET	_	EXPRESSION TAG	UNP P13633
В	9	GLY	_	EXPRESSION TAG	UNP P13633
В	10	SER	-	EXPRESSION TAG	UNP P13633
В	11	SER	-	EXPRESSION TAG	UNP P13633
В	12	HIS	-	EXPRESSION TAG	UNP P13633
В	13	HIS	_	EXPRESSION TAG	UNP P13633
В	14	HIS	_	EXPRESSION TAG	UNP P13633
В	15	HIS	_	EXPRESSION TAG	UNP P13633
В	16	HIS	_	EXPRESSION TAG	UNP P13633
В	17	HIS	_	EXPRESSION TAG	UNP P13633
В	18	SER	-	EXPRESSION TAG	UNP P13633
В	19	SER	-	EXPRESSION TAG	UNP P13633
В	20	GLY	-	EXPRESSION TAG	UNP P13633
В	21	LEU	_	EXPRESSION TAG	UNP P13633
В	22	VAL	_	EXPRESSION TAG	UNP P13633
В	23	PRO	_	EXPRESSION TAG	UNP P13633
В	24	ARG	_	EXPRESSION TAG	UNP P13633
В	25	GLY	-	EXPRESSION TAG	UNP P13633
В	26	SER	_	EXPRESSION TAG	UNP P13633
В	27	HIS	-	EXPRESSION TAG	UNP P13633
В	28	MSE	-	EXPRESSION TAG	UNP P13633
В	29	ALA	-	EXPRESSION TAG	UNP P13633
В	30	SER	-	EXPRESSION TAG	UNP P13633
В	31	MSE	-	EXPRESSION TAG	UNP P13633
В	32	THR	-	EXPRESSION TAG	UNP P13633
В	33	GLY	-	EXPRESSION TAG	UNP P13633
В	34	GLY	-	EXPRESSION TAG	UNP P13633
В	35	GLU	-	EXPRESSION TAG	UNP P13633
В	36	GLU	-	EXPRESSION TAG	UNP P13633
В	37	MSE	-	EXPRESSION TAG	UNP P13633
В	38	GLY	-	EXPRESSION TAG	UNP P13633

Continued on next page...




$\alpha$	7 .	e	•	
Lanting	ed 1	trom	previous	naae
Constitu	uu i	10116	$\rho_1 \cup \sigma_0 \cup \sigma_0$	$\rho u q c \dots$

Chain	Residue	Modelled	Actual	${f Comment}$	Reference
В	39	ARG	_	EXPRESSION TAG	UNP P13633
В	40	GLY	-	EXPRESSION TAG	UNP P13633
В	41	SER	-	EXPRESSION TAG	UNP P13633
В	174	LYS	ASN	SEE REMARK 999	UNP P13633
В	309	ASN	LYS	SEE REMARK 999	UNP P13633

• Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	В	1	Total Mg 1 1	0	0
2	A	1	Total Mg 1 1	0	0

 $\bullet$  Molecule 3 is SUCCINIC ACID (three-letter code: SIN) (formula:  $\mathrm{C_4H_6O_4}).$ 



-	Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
	3	A	1	Total C O 8 4 4	0	0
	3	A	1	Total C O 8 4 4	0	0

• Molecule 4 is water.




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	110	Total O 110 110	0	0
4	В	89	Total O 89 89	0	0




# 3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: C4-dicarboxylate transport sensor protein dctB



• Molecule 1: C4-dicarboxylate transport sensor protein dctB





# 4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants	101.89Å 66.59Å 93.60Å	Danagitan
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $94.71^{\circ}$ $90.00^{\circ}$	Depositor
Resolution (Å)	20.00 - 2.30	Depositor
Resolution (A)	48.79 - 2.30	EDS
% Data completeness	100.0 (20.00-2.30)	Depositor
(in resolution range)	99.6 (48.79-2.30)	EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.58 (at 2.29Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
D D.	0.200 , 0.252	Depositor
$R, R_{free}$	0.203 , $0.256$	DCC
$R_{free}$ test set	1299 reflections $(4.67\%)$	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	35.9	Xtriage
Anisotropy	0.426	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$ , $B_{sol}(Å^2)$	0.34, 30.1	EDS
L-test for twinning <sup>2</sup>	$ < L >=0.46, < L^2>=0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	4107	wwPDB-VP
Average B, all atoms $(Å^2)$	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 7.15% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of <|L|>,  $< L^2>$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 5 Model quality (i)

## 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, SIN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Boı	nd lengths	Bond angles	
IVIOI	Chain	RMSZ	# Z  > 5	RMSZ	# Z >5
1	A	0.69	1/1981 (0.1%)	0.74	1/2687 (0.0%)
1	В	0.63	1/1981 (0.1%)	0.70	0/2687
All	All	0.66	$2/3962 \ (0.1\%)$	0.72	1/5374 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	Α	0	1

#### All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	$\operatorname{Observed}(\text{\AA})$	$\operatorname{Ideal}(\text{\AA})$
1	В	169	MSE	SE-CE	-5.87	1.60	1.95
1	A	169	MSE	SE-CE	-5.85	1.60	1.95

#### All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	$\mathbf{Observed}(^o)$	$\mathbf{Ideal}(^o)$
1	A	150	ALA	N-CA-C	-7.86	89.77	111.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group	
1	Α	149	TYR	Peptide	



### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1945	0	1975	26	0
1	В	1945	0	1975	44	1
2	A	1	0	0	0	0
2	В	1	0	0	0	0
3	A	16	0	8	0	0
4	A	110	0	0	4	0
4	В	89	0	0	2	1
All	All	4107	0	3958	68	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 68 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$egin{aligned}  ext{Interatomic} \  ext{distance} \ ( ext{Å}) \end{aligned}$	$egin{aligned}  ext{Clash} \  ext{overlap } ( ext{Å}) \end{aligned}$
1:A:294:MSE:HG3	1:B:301:ARG:HH21	1.15	1.03
1:B:182:ARG:HH21	1:B:182:ARG:HG2	1.24	1.02
1:B:294:MSE:HG2	1:B:301:ARG:NH2	1.76	1.00
1:B:182:ARG:HH21	1:B:182:ARG:CG	1.92	0.83
1:B:294:MSE:HG2	1:B:301:ARG:CZ	2.09	0.80

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	$egin{aligned}  ext{Interatomic} \  ext{distance} & ( ext{Å}) \end{aligned}$	$egin{aligned}  ext{Clash} \  ext{overlap } ( ext{Å}) \end{aligned}$
1:B:284:SER:OG	4:B:607:HOH:O[4_546]	2.19	0.01

## 5.3 Torsion angles (i)

### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.



The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	$\mathbf{ntiles}$
1	A	252/305~(83%)	248 (98%)	4 (2%)	0	100	100
1	В	252/305~(83%)	247 (98%)	5 (2%)	0	100	100
All	All	504/610 (83%)	495 (98%)	9 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric Outlie		Percentiles
1	A	203/235~(86%)	193 (95%)	10 (5%)	25 35
1	В	203/235~(86%)	184 (91%)	19 (9%)	8 10
All	All	406/470 (86%)	377 (93%)	29 (7%)	14 19

5 of 29 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	В	105	ASP
1	В	163	MSE
1	В	295	VAL
1	В	114	LEU
1	В	182	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	208	GLN
1	В	79	GLN
1	В	108	ASN
1	В	304	GLN



#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry (i)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	B	ond leng	$_{ m gths}$	В	ond ang	gles
MIGI	туре	Chain	nes	es Link	Counts	RMSZ	# Z >2	Counts	RMSZ	# Z  > 2
3	SIN	A	801	_	1,7,7	0.17	0	2,8,8	1.40	0
3	SIN	A	701	-	1,7,7	0.04	0	2,8,8	0.77	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

$\mathbf{Mol}$	$\mathbf{Type}$	Chain	${f Res}$	Link	Chirals	Torsions	Rings
3	SIN	A	801	-	-	0/1/5/5	-
3	SIN	A	701	-	-	0/1/5/5	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.



There are no ring outliers.

No monomer is involved in short contacts.

# 5.7 Other polymers (i)

There are no such residues in this entry.

## 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



## 6 Fit of model and data (i)

### 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	$\#\mathrm{RSRZ}{>}2$		$OWAB(A^2)$	Q < 0.9
1	A	250/305~(81%)	-0.12	2 (0%)	86 89	21, 34, 46, 54	0
1	В	250/305~(81%)	-0.00	1 (0%)	92 95	21, 35, 50, 58	2 (0%)
All	All	500/610 (81%)	-0.06	3 (0%)	89 92	21, 35, 49, 58	2 (0%)

#### All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	239	ILE	3.3
1	В	238	PRO	2.7
1	A	243	ARG	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates (i)

There are no carbohydrates in this entry.

### 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q < 0.9
2	MG	В	601	1/1	0.97	0.05	38,38,38,38	0
2	MG	A	501	1/1	0.97	0.05	34,34,34,34	0

Continued on next page...



Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	${f B\text{-factors}}({f \AA}^2)$	Q < 0.9
3	SIN	A	701	8/8	0.97	0.10	26,28,29,30	0
3	SIN	A	801	8/8	0.98	0.07	19,21,22,23	0

# 6.5 Other polymers (i)

There are no such residues in this entry.

