

# wwPDB X-ray Structure Validation Summary Report (i)

Jun 14, 2020 – 12:56 am BST

PDB ID : 1E7L

Title: Endonuclease VII (EndoVII) N62D mutant from phage T4

Authors: Raaijmakers, H.C.A.; Vix, O.; Toro, I.; Suck, D.

Deposited on : 2000-08-29

Resolution : 1.32 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : NOT EXECUTED EDS : NOT EXECUTED

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.11

## 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$ 

The reported resolution of this entry is 1.32 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
Metric	$(\# \mathrm{Entries})$	$(\#  ext{Entries},  ext{resolution range}( ext{Å}))$
Clashscore	141614	1667 (1.34-1.30)
Ramachandran outliers	138981	1615 (1.34-1.30)
Sidechain outliers	138945	1615 (1.34-1.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Note EDS was not executed.

Mol	Chain	Length	Quality of chain		
1	A	157	86%	13%	•
1	В	157	87%	13%	



# 2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 3257 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


• Molecule 1 is a protein called RECOMBINATION ENDONUCLEASE VII.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	٨	157	Total	С	N	О	S	0	22	0
1	Α	157	1378	866	243	258	11	U	22	U
1	D	157	Total	С	N	О	S	0	19	0
1	Ъ	197	1369	857	243	259	10	0	19	U

There are 2 discrepancies between the modelled and reference sequences:

	Chain	Residue	Modelled	Actual	Comment	Reference
	Α	62	ASP	ASN	engineered mutation	UNP P13340
ĺ	В	62	ASP	ASN	engineered mutation	UNP P13340

• Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	${f Atoms}$		ZeroOcc	AltConf
2	A	1	Total 0	O S 8 2	0	1

Continued on next page...



Continued from previous page...

Mol	Chain	Residues	Ato	ms		ZeroOcc	AltConf	
2	A	1	Total	О	S	0	0	
	A	1	5	4	1	0	0	
2	A	1	Total	О	S	0	0	
	Λ	1	5	4	1	U	U	
2	A	1	Total	Ο	S	0	0	
	11	1	5	4	1	U	U	
$\frac{1}{2}$	A	1	Total	Ο	S	0	0	
	11	1	5	4	1	· ·	0	
$\frac{1}{2}$	A	1	Total	Ο	S	0	0	
	11	-	5	4	1	Ü		
2	A	1	Total	Ο	$\mathbf{S}$	0	0	
_		_	5	4	1	Ü		
2	В	1	Total	O	S	0	0	
			5	4	1	_	-	
2	В	1	Total	O	S	0	0	
			5	4	1			
2	В	1	Total	O	S	0	1	
			10	8	2			
2	В	1	Total	O	S	0	0	
			5 T-4-1	4	1			
2	В	1	Total 5	O	S 1	0	0	
			Total	$\frac{4}{O}$	$\frac{1}{S}$			
2	В	1	10tai   5	4	5 1	0	0	
			Total	$\frac{4}{O}$	$\frac{1}{S}$			
2	В	1	5	4	1	0	0	

• Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	В	1	Total Zn 1 1	0	0
3	A	1	Total Zn 1 1	0	0

#### • Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	193	Total O 193 193	0	0
4	В	235	Total O 235 235	0	0




# 3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: RECOMBINATION ENDONUCLEASE VII





# 4 Data and refinement statistics (i)

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source	
Space group	P 1 21 1	Depositor	
Cell constants	58.24Å 35.83Å 91.93Å	Depositor	
a, b, c, $\alpha$ , $\beta$ , $\gamma$	90.00° 103.93° 90.00°	Depositor	
Resolution (Å)	35.00 - 1.32	Depositor	
% Data completeness	96.5 (35.00-1.32)	Depositor	
(in resolution range)	30.9 (39.00 1.32)	Depositor	
$R_{merge}$	(Not available)	Depositor	
$R_{sym}$	0.06	Depositor	
Refinement program	REFMAC	Depositor	
$R, R_{free}$	0.146 , 0.186	Depositor	
Estimated twinning fraction	No twinning to report.	Xtriage	
Total number of atoms	3257	wwPDB-VP	
Average B, all atoms (Å <sup>2</sup> )	30.0	wwPDB-VP	



# 5 Model quality (i)

### 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
MIOI	Chain	RMSZ	# Z >5	RMSZ	# Z  > 5	
1	A	0.56	0/1491	0.95	7/1981 (0.4%)	
1	В	0.52	0/1471	0.84	3/1955~(0.2%)	
All	All	0.54	0/2962	0.90	$10/3936 \ (0.3\%)$	

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	Α	0	1

There are no bond length outliers.

The worst 5 of 10 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
1	A	28	ARG	CD-NE-CZ	9.54	136.95	123.60
1	A	54	ARG	NE-CZ-NH2	-7.21	116.70	120.30
1	В	131	ARG	CD-NE-CZ	6.66	132.93	123.60
1	В	131	ARG	NE-CZ-NH1	6.43	123.52	120.30
1	A	153[A]	ARG	CG-CD-NE	6.06	124.53	111.80

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	92[B]	LEU	Mainchain



#### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1378	0	1386	46	0
1	В	1369	0	1366	49	0
2	A	40	0	0	2	0
2	В	40	0	0	1	0
3	A	1	0	0	0	0
3	В	1	0	0	0	0
4	A	193	0	0	11	0
4	В	235	0	0	11	0
All	All	3257	0	2752	75	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

The worst 5 of 75 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$egin{array}{c}  ext{Interatomic} \  ext{distance} \ ( ext{Å}) \end{array}$	$egin{aligned}  ext{Clash} \  ext{overlap } ( ext{Å}) \end{aligned}$
1:B:118[B]:ARG:CG	1:B:118[B]:ARG:CD	1.83	1.53
1:B:118[B]:ARG:CB	1:B:118[B]:ARG:CG	1.86	1.52
1:A:52[B]:LYS:CE	1:A:52[B]:LYS:NZ	1.82	1.41
1:A:89[B]:GLU:OE1	1:B:92[B]:LEU:CD2	1.95	1.14
1:A:89[B]:GLU:OE1	1:B:92[B]:LEU:HD21	1.53	1.04

There are no symmetry-related clashes.

### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	$\mathbf{ntiles}$
1	A	178/157 (113%)	178 (100%)	0	0	100	100
1	В	174/157 (111%)	172 (99%)	2 (1%)	0	100	100
All	All	352/314 (112%)	350 (99%)	2 (1%)	0	100	100

There are no Ramachandran outliers to report.

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	${f Analysed}$	Rotameric	Outliers	Percentiles		
1	A	$161/138 \; (117\%)$	161 (100%)	0	100	100	
1	В	158/138 (114%)	158 (100%)	0	100	100	
All	All	319/276 (116%)	319 (100%)	0	100	100	

There are no protein residues with a non-rotameric sidechain to report.

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates (i)

There are no carbohydrates in this entry.



### 5.6 Ligand geometry (i)

Of 18 ligands modelled in this entry, 2 are monoatomic - leaving 16 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
WIOI	Type	Chain	nes	Lilik	Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	SO4	A	1160	-	4,4,4	0.71	0	6,6,6	0.44	0
2	SO4	A	1161	-	4,4,4	0.60	0	6,6,6	0.05	0
2	SO4	В	1158	_	4,4,4	0.51	0	6,6,6	0.30	0
2	SO4	A	1158[A]	_	4,4,4	0.59	0	6,6,6	0.17	0
2	SO4	В	1163	_	4,4,4	0.60	0	6,6,6	0.10	0
2	SO4	A	1163	_	4,4,4	0.66	0	6,6,6	0.15	0
2	SO4	В	1162	-	4,4,4	0.59	0	6,6,6	0.17	0
2	SO4	В	1160[B]	_	4,4,4	0.64	0	6,6,6	0.14	0
2	SO4	A	1158[B]	_	4,4,4	0.61	0	6,6,6	0.17	0
2	SO4	A	1159	_	4,4,4	0.62	0	6,6,6	0.09	0
2	SO4	В	1159	_	4,4,4	0.81	0	6,6,6	0.42	0
2	SO4	A	1164	-	4,4,4	0.62	0	6,6,6	0.06	0
2	SO4	В	1164	-	4,4,4	0.63	0	6,6,6	0.08	0
2	SO4	В	1160[A]	-	4,4,4	0.57	0	6,6,6	0.12	0
2	SO4	A	1162	-	4,4,4	0.61	0	6,6,6	0.05	0
2	SO4	В	1161	_	4,4,4	0.64	0	6,6,6	0.09	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1161	SO4	1	0
2	A	1158[A]	SO4	1	0
2	В	1160[B]	SO4	1	0



# 5.7 Other polymers (i)

There are no such residues in this entry.

# 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



## 6 Fit of model and data (i)

#### 6.1 Protein, DNA and RNA chains (i)

EDS was not executed - this section is therefore empty.

#### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS was not executed - this section is therefore empty.

### 6.3 Carbohydrates (i)

EDS was not executed - this section is therefore empty.

### 6.4 Ligands (i)

EDS was not executed - this section is therefore empty.

#### 6.5 Other polymers (i)

EDS was not executed - this section is therefore empty.

