



wwPDB X-ray Structure Validation Summary Report ⓘ

Sep 19, 2023 – 12:09 PM EDT

PDB ID : 5E84
Title : ATP-bound state of BiP
Authors : Liu, Q.; Yang, J.; Nune, M.; Zong, Y.; Zhou, L.
Deposited on : 2015-10-13
Resolution : 2.99 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : 1.13
EDS : 2.35.1
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.99 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2092 (3.00-3.00)
Clashscore	141614	2416 (3.00-3.00)
Ramachandran outliers	138981	2333 (3.00-3.00)
Sidechain outliers	138945	2336 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	606	84% 14% .
1	B	606	89% 9% .
1	C	606	88% 11% .
1	D	606	86% 13% .
1	E	606	85% 14% .
1	F	606	85% 13% .

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	MG	D	804	-	-	X	-
5	SO4	A	808	-	-	X	-
5	SO4	C	806	-	-	X	-
5	SO4	D	808	-	-	X	-
5	SO4	E	806	-	-	X	-
5	SO4	F	806	-	-	X	-

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 28484 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 78 kDa glucose-regulated protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	606	4710	2955	806	938	11	8	0	0
1	B	606	4700	2948	806	935	11	8	0	0
1	C	606	4700	2948	806	935	11	8	0	0
1	D	606	4700	2948	806	935	11	8	0	0
1	E	606	4696	2945	805	935	11	8	0	0
1	F	606	4700	2948	806	935	11	8	0	0

There are 54 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	24	SER	-	expression tag	UNP P11021
A	229	ALA	THR	conflict	UNP P11021
A	?	-	THR	deletion	UNP P11021
A	?	-	ALA	deletion	UNP P11021
A	?	-	SER	deletion	UNP P11021
A	?	-	ASP	deletion	UNP P11021
A	453	VAL	ASN	conflict	UNP P11021
A	454	GLY	GLN	conflict	UNP P11021
A	455	GLY	PRO	conflict	UNP P11021
B	24	SER	-	expression tag	UNP P11021
B	229	ALA	THR	conflict	UNP P11021
B	?	-	THR	deletion	UNP P11021
B	?	-	ALA	deletion	UNP P11021
B	?	-	SER	deletion	UNP P11021
B	?	-	ASP	deletion	UNP P11021
B	453	VAL	ASN	conflict	UNP P11021
B	454	GLY	GLN	conflict	UNP P11021

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	455	GLY	PRO	conflict	UNP P11021
C	24	SER	-	expression tag	UNP P11021
C	229	ALA	THR	conflict	UNP P11021
C	?	-	THR	deletion	UNP P11021
C	?	-	ALA	deletion	UNP P11021
C	?	-	SER	deletion	UNP P11021
C	?	-	ASP	deletion	UNP P11021
C	453	VAL	ASN	conflict	UNP P11021
C	454	GLY	GLN	conflict	UNP P11021
C	455	GLY	PRO	conflict	UNP P11021
D	24	SER	-	expression tag	UNP P11021
D	229	ALA	THR	conflict	UNP P11021
D	?	-	THR	deletion	UNP P11021
D	?	-	ALA	deletion	UNP P11021
D	?	-	SER	deletion	UNP P11021
D	?	-	ASP	deletion	UNP P11021
D	453	VAL	ASN	conflict	UNP P11021
D	454	GLY	GLN	conflict	UNP P11021
D	455	GLY	PRO	conflict	UNP P11021
E	24	SER	-	expression tag	UNP P11021
E	229	ALA	THR	conflict	UNP P11021
E	?	-	THR	deletion	UNP P11021
E	?	-	ALA	deletion	UNP P11021
E	?	-	SER	deletion	UNP P11021
E	?	-	ASP	deletion	UNP P11021
E	453	VAL	ASN	conflict	UNP P11021
E	454	GLY	GLN	conflict	UNP P11021
E	455	GLY	PRO	conflict	UNP P11021
F	24	SER	-	expression tag	UNP P11021
F	229	ALA	THR	conflict	UNP P11021
F	?	-	THR	deletion	UNP P11021
F	?	-	ALA	deletion	UNP P11021
F	?	-	SER	deletion	UNP P11021
F	?	-	ASP	deletion	UNP P11021
F	453	VAL	ASN	conflict	UNP P11021
F	454	GLY	GLN	conflict	UNP P11021
F	455	GLY	PRO	conflict	UNP P11021

- Molecule 2 is ADENOSINE-5'-TRIPHOSPHATE (three-letter code: ATP) (formula: $C_{10}H_{16}N_5O_{13}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	Total	C	N	O	P	0	0
			31	10	5	13	3		
2	B	1	Total	C	N	O	P	0	0
			31	10	5	13	3		
2	C	1	Total	C	N	O	P	0	0
			31	10	5	13	3		
2	D	1	Total	C	N	O	P	0	0
			31	10	5	13	3		
2	E	1	Total	C	N	O	P	0	0
			31	10	5	13	3		
2	F	1	Total	C	N	O	P	0	0
			31	10	5	13	3		


- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Zn		
3	A	3	Total	Zn	0	0
			3	3		
3	B	3	Total	Zn	0	0
			3	3		
3	C	3	Total	Zn	0	0
			3	3		
3	D	3	Total	Zn	0	0
			3	3		
3	E	3	Total	Zn	0	0
			3	3		
3	F	3	Total	Zn	0	0
			3	3		

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	3	Total 3	Mg 3	0	0
4	B	3	Total 3	Mg 3	0	0
4	C	1	Total 1	Mg 1	0	0
4	D	3	Total 3	Mg 3	0	0
4	E	1	Total 1	Mg 1	0	0
4	F	1	Total 1	Mg 1	0	0

- Molecule 5 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total 5	O 4	S 1	0	0
5	B	1	Total 5	O 4	S 1	0	0
5	C	1	Total 5	O 4	S 1	0	0
5	D	1	Total 5	O 4	S 1	0	0
5	E	1	Total 5	O 4	S 1	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	F	1	Total	O	S	0	0
			5	4	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	7	Total	O	0	0
			7	7		
6	B	5	Total	O	0	0
			5	5		
6	C	4	Total	O	0	0
			4	4		
6	D	4	Total	O	0	0
			4	4		
6	E	9	Total	O	0	0
			9	9		
6	F	3	Total	O	0	0
			3	3		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: 78 kDa glucose-regulated protein




- Molecule 1: 78 kDa glucose-regulated protein




- Molecule 1: 78 kDa glucose-regulated protein




- Molecule 1: 78 kDa glucose-regulated protein



- Molecule 1: 78 kDa glucose-regulated protein



- Molecule 1: 78 kDa glucose-regulated protein



4 Data and refinement statistics i

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	222.47Å 222.47Å 209.46Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	40.76 – 2.99 39.65 – 2.99	Depositor EDS
% Data completeness (in resolution range)	97.6 (40.76-2.99) 97.6 (39.65-2.99)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.42 (at 3.01Å)	Xtrriage
Refinement program	REFMAC 5.7.0032	Depositor
R, R_{free}	0.243 , 0.287 0.241 , 0.285	Depositor DCC
R_{free} test set	5907 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	67.3	Xtrriage
Anisotropy	0.115	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 27.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.42$, $\langle L^2 \rangle = 0.24$	Xtrriage
Estimated twinning fraction	0.309 for -h,-k,l	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	28484	wwPDB-VP
Average B, all atoms (Å ²)	77.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.31% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: ATP, ZN, SO4, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.75	9/4777 (0.2%)	0.86	7/6446 (0.1%)
1	B	0.60	4/4767 (0.1%)	0.73	5/6432 (0.1%)
1	C	0.56	1/4767 (0.0%)	0.71	2/6432 (0.0%)
1	D	0.67	3/4767 (0.1%)	0.78	8/6432 (0.1%)
1	E	0.66	1/4763 (0.0%)	0.73	4/6428 (0.1%)
1	F	0.61	3/4767 (0.1%)	0.75	6/6432 (0.1%)
All	All	0.65	21/28608 (0.1%)	0.76	32/38602 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	C	0	3
1	D	2	2
1	F	0	3
All	All	2	9

The worst 5 of 21 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	E	576	GLU	CB-CG	25.85	2.01	1.52
1	D	577	LYS	CB-CG	-17.80	1.04	1.52
1	F	577	LYS	CB-CG	-13.61	1.15	1.52
1	B	577	LYS	CB-CG	-13.20	1.17	1.52
1	D	576	GLU	CB-CG	-12.78	1.27	1.52

The worst 5 of 32 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	F	556	ASP	CB-CG-OD1	11.50	128.65	118.30
1	A	556	ASP	CB-CG-OD1	11.41	128.57	118.30
1	D	577	LYS	CA-CB-CG	9.73	134.82	113.40
1	A	599	GLU	OE1-CD-OE2	-9.65	111.72	123.30
1	E	576	GLU	CA-CB-CG	-9.10	93.39	113.40

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
1	D	488	ARG	CA
1	D	514	THR	CA

5 of 9 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	488	ARG	Peptide
1	C	482	GLY	Peptide
1	C	483	ILE	Peptide
1	C	484	PRO	Peptide
1	D	487	PRO	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4710	0	4736	47	4
1	B	4700	0	4713	33	2
1	C	4700	0	4713	37	1
1	D	4700	0	4713	52	0
1	E	4696	0	4702	51	4
1	F	4700	0	4713	58	1
2	A	31	0	12	0	0
2	B	31	0	12	0	0
2	C	31	0	12	0	0
2	D	31	0	12	1	0
2	E	31	0	12	1	0
2	F	31	0	12	1	0
3	A	3	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	3	0	0	0	0
3	C	3	0	0	0	0
3	D	3	0	0	0	0
3	E	3	0	0	0	0
3	F	3	0	0	0	0
4	A	3	0	0	1	2
4	B	3	0	0	0	0
4	C	1	0	0	0	1
4	D	3	0	0	0	2
4	E	1	0	0	0	0
4	F	1	0	0	0	0
5	A	5	0	0	6	0
5	B	5	0	0	1	0
5	C	5	0	0	6	0
5	D	5	0	0	5	0
5	E	5	0	0	7	0
5	F	5	0	0	5	0
6	A	7	0	0	2	0
6	B	5	0	0	4	1
6	C	4	0	0	0	0
6	D	4	0	0	1	0
6	E	9	0	0	3	0
6	F	3	0	0	1	0
All	All	28484	0	28362	262	12

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 262 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:303:HIS:ND1	1:F:556:ASP:OD1	1.97	0.95
1:D:556:ASP:OD2	1:F:303:HIS:CE1	2.26	0.89
1:B:552:LYS:HE2	1:B:556:ASP:OD2	1.75	0.87
1:D:556:ASP:OD2	1:F:303:HIS:ND1	2.12	0.82
1:A:415:VAL:HG11	1:D:417:LEU:HD21	1.63	0.81

The worst 5 of 12 symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:317:ASP:OD2	4:D:804:MG:MG[6_654]	1.44	0.76
4:A:803:MG:MG	4:C:803:MG:MG[2_554]	1.48	0.72
1:C:317:ASP:OD2	4:A:804:MG:MG[6_554]	1.50	0.70
1:A:303:HIS:ND1	1:A:556:ASP:OD1[6_554]	1.54	0.66
1:A:595:GLU:OE2	1:A:595:GLU:OE2[5_554]	1.62	0.58

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	604/606 (100%)	566 (94%)	29 (5%)	9 (2%)	10	42
1	B	604/606 (100%)	569 (94%)	32 (5%)	3 (0%)	29	68
1	C	604/606 (100%)	562 (93%)	36 (6%)	6 (1%)	15	53
1	D	604/606 (100%)	561 (93%)	34 (6%)	9 (2%)	10	42
1	E	604/606 (100%)	560 (93%)	38 (6%)	6 (1%)	15	53
1	F	604/606 (100%)	564 (93%)	32 (5%)	8 (1%)	12	45
All	All	3624/3636 (100%)	3382 (93%)	201 (6%)	41 (1%)	14	50

5 of 41 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	408	ASP
1	A	516	ASN
1	C	483	ILE
1	C	517	LYS
1	D	408	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	515/518 (99%)	498 (97%)	17 (3%)	38	73
1	B	511/518 (99%)	495 (97%)	16 (3%)	40	75
1	C	511/518 (99%)	490 (96%)	21 (4%)	30	67
1	D	511/518 (99%)	494 (97%)	17 (3%)	38	73
1	E	510/518 (98%)	495 (97%)	15 (3%)	42	76
1	F	511/518 (99%)	495 (97%)	16 (3%)	40	75
All	All	3069/3108 (99%)	2967 (97%)	102 (3%)	38	73

5 of 102 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	146	SER
1	E	66	VAL
1	F	567	SER
1	D	176	PHE
1	D	528	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 11 such sidechains are listed below:

Mol	Chain	Res	Type
1	E	167	HIS
1	E	492	GLN
1	F	194	ASN
1	F	167	HIS
1	C	167	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 42 ligands modelled in this entry, 30 are monoatomic - leaving 12 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	ATP	A	801	3	26,33,33	0.92	0	31,52,52	1.60	7 (22%)
2	ATP	B	801	3	26,33,33	0.88	0	31,52,52	1.62	6 (19%)
5	SO4	F	806	-	4,4,4	0.33	0	6,6,6	0.34	0
5	SO4	D	808	-	4,4,4	0.37	0	6,6,6	0.25	0
2	ATP	C	801	3	26,33,33	0.87	0	31,52,52	1.64	7 (22%)
2	ATP	E	801	3	26,33,33	1.00	1 (3%)	31,52,52	1.42	5 (16%)
5	SO4	E	806	-	4,4,4	0.37	0	6,6,6	0.55	0
5	SO4	C	806	-	4,4,4	0.30	0	6,6,6	0.37	0
5	SO4	A	808	-	4,4,4	0.41	0	6,6,6	0.39	0
5	SO4	B	808	-	4,4,4	0.41	0	6,6,6	0.44	0
2	ATP	D	801	3	26,33,33	0.81	0	31,52,52	1.66	7 (22%)
2	ATP	F	801	3	26,33,33	1.10	2 (7%)	31,52,52	1.67	7 (22%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ATP	A	801	3	-	1/18/38/38	0/3/3/3
2	ATP	B	801	3	-	3/18/38/38	0/3/3/3
2	ATP	C	801	3	-	1/18/38/38	0/3/3/3
2	ATP	E	801	3	-	5/18/38/38	0/3/3/3
2	ATP	D	801	3	-	1/18/38/38	0/3/3/3
2	ATP	F	801	3	-	1/18/38/38	0/3/3/3

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	F	801	ATP	C5-C4	2.76	1.48	1.40
2	E	801	ATP	C5-C4	2.25	1.46	1.40
2	F	801	ATP	C2-N3	2.03	1.35	1.32

The worst 5 of 39 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	F	801	ATP	PA-O3A-PB	-5.17	115.09	132.83
2	C	801	ATP	N3-C2-N1	-4.57	121.54	128.68
2	B	801	ATP	PA-O3A-PB	-4.49	117.41	132.83
2	D	801	ATP	PA-O3A-PB	-4.08	118.83	132.83
2	B	801	ATP	N3-C2-N1	-3.78	122.77	128.68

There are no chirality outliers.

5 of 12 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	801	ATP	PB-O3B-PG-O3G
2	E	801	ATP	PB-O3B-PG-O2G
2	E	801	ATP	PB-O3B-PG-O3G
2	E	801	ATP	C5'-O5'-PA-O3A
2	E	801	ATP	PB-O3A-PA-O1A


There are no ring outliers.


9 monomers are involved in 33 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	F	806	SO4	5	0
5	D	808	SO4	5	0
2	E	801	ATP	1	0
5	E	806	SO4	7	0
5	C	806	SO4	6	0
5	A	808	SO4	6	0
5	B	808	SO4	1	0
2	D	801	ATP	1	0
2	F	801	ATP	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is


within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.














5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.


6.3 Carbohydrates [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.4 Ligands [i](#)




Unable to reproduce the depositors R factor - this section is therefore empty.

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around ATP B 801:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around ATP C 801:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around ATP D 801:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around ATP E 801:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.