



# Full wwPDB X-ray Structure Validation Report ⓘ

Nov 21, 2023 – 01:16 AM JST

PDB ID : 7E8N  
Title : Crystal structure of Type II citrate synthase (HyCS) from Hymenobacter sp. PAMC 26554  
Authors : Park, S.-H.; Lee, C.W.; Bae, D.-W.; Lee, J.H.  
Deposited on : 2021-03-02  
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4898 (2.20-2.20)
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)
RSRZ outliers	127900	4800 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	434	
1	B	434	
1	C	434	

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 10484 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Citrate synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	428	3359	2148	559	636	16	0	0	0
1	B	424	3338	2139	554	628	17	0	0	0
1	C	428	3354	2146	557	634	17	0	0	0

There are 9 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	GLY	-	expression tag	UNP A0A142HAE2
A	-1	SER	-	expression tag	UNP A0A142HAE2
A	0	HIS	-	expression tag	UNP A0A142HAE2
B	-2	GLY	-	expression tag	UNP A0A142HAE2
B	-1	SER	-	expression tag	UNP A0A142HAE2
B	0	HIS	-	expression tag	UNP A0A142HAE2
C	-2	GLY	-	expression tag	UNP A0A142HAE2
C	-1	SER	-	expression tag	UNP A0A142HAE2
C	0	HIS	-	expression tag	UNP A0A142HAE2

- Molecule 2 is CITRIC ACID (three-letter code: CIT) (formula: C<sub>6</sub>H<sub>8</sub>O<sub>7</sub>) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 13 6 7	0	0
2	B	1	Total C O 13 6 7	0	0
2	C	1	Total C O 13 6 7	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	160	Total O 160 160	0	0
3	B	141	Total O 141 141	0	0
3	C	93	Total O 93 93	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Citrate synthase



- Molecule 1: Citrate synthase



- Molecule 1: Citrate synthase





## 4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	134.34Å 166.17Å 135.34Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	41.58 – 2.20 41.54 – 2.20	Depositor EDS
% Data completeness (in resolution range)	99.6 (41.58-2.20) 99.6 (41.54-2.20)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	7.21 (at 2.20Å)	Xtrriage
Refinement program	REFMAC 5.8.0267	Depositor
R, $R_{free}$	0.190 , 0.235 0.196 , 0.239	Depositor DCC
$R_{free}$ test set	3848 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	37.6	Xtrriage
Anisotropy	0.107	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 48.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	10484	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	49.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.01% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CIT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.72	0/3437	0.81	0/4659
1	B	0.75	0/3414	0.88	3/4625 (0.1%)
1	C	0.72	0/3431	0.84	0/4650
All	All	0.73	0/10282	0.84	3/13934 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	403	ARG	NE-CZ-NH2	-8.76	115.92	120.30
1	B	232	GLU	N-CA-CB	-5.87	100.03	110.60
1	B	140	CYS	N-CA-CB	5.61	120.70	110.60

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3359	0	3321	13	0
1	B	3338	0	3312	29	0
1	C	3354	0	3315	33	0
2	A	13	0	5	0	0
2	B	13	0	5	0	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	C	13	0	5	4	0
3	A	160	0	0	0	0
3	B	141	0	0	4	0
3	C	93	0	0	1	0
All	All	10484	0	9963	72	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (72) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:340:LEU:HD12	1:C:340:LEU:O	1.76	0.84
1:B:348:GLN:O	1:B:352:THR:HG23	1.90	0.72
1:B:324:ALA:O	1:B:328:VAL:HG23	1.90	0.70
1:C:403:ARG:NH1	3:C:602:HOH:O	2.28	0.67
1:C:300:GLY:O	1:C:301:PHE:HB2	1.96	0.65
1:A:411:ARG:NH1	2:C:501:CIT:O4	2.32	0.62
1:A:271:GLN:HG3	1:A:380:THR:HG22	1.80	0.62
1:C:340:LEU:HD12	1:C:340:LEU:C	2.22	0.59
1:C:366:ASP:HA	1:C:369:SER:HB2	1.86	0.58
1:B:179:LYS:HD2	1:B:184:HIS:HB2	1.85	0.57
1:C:270:ASN:HB2	1:C:303:LEU:O	2.03	0.57
1:A:28:ALA:HA	1:C:41:VAL:O	2.04	0.57
1:C:231:HIS:ND1	2:C:501:CIT:H21	2.20	0.56
1:C:336:ASN:O	1:C:336:ASN:ND2	2.36	0.55
1:B:7:LEU:HB3	1:B:15:ILE:HG23	1.88	0.55
1:A:250:LEU:HD13	1:A:399:TRP:CD2	2.42	0.55
1:B:227:LEU:HD13	1:B:321:ILE:HG13	1.89	0.55
1:B:288:SER:HA	1:B:291:ILE:HB	1.90	0.53
1:B:313:ASN:HA	1:B:362:TYR:CE1	2.45	0.52
1:B:75:GLU:HG3	1:B:320:ILE:HD11	1.91	0.51
1:C:390:GLY:O	1:C:393:PRO:HD2	2.10	0.51
1:B:306:PHE:HZ	1:B:347:GLU:HA	1.75	0.51
1:B:359:ARG:HB3	1:B:361:LEU:CD2	2.41	0.50
1:C:274:ILE:HB	1:C:380:THR:HG22	1.94	0.50
1:B:108:ILE:O	1:B:112:THR:HG23	2.12	0.50
1:B:9:LEU:HB3	1:B:10:PRO:HD2	1.95	0.49
1:C:321:ILE:HG23	1:C:368:TYR:HB2	1.96	0.48
1:A:271:GLN:O	1:A:274:ILE:HG22	2.13	0.48
1:A:422:ARG:HG3	1:C:72:TYR:CE1	2.49	0.48

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:269:ALA:O	1:B:273:VAL:HG23	2.13	0.48
1:B:7:LEU:HB3	1:B:15:ILE:CG2	2.43	0.48
1:B:211:TYR:HA	3:B:601:HOH:O	2.14	0.47
1:B:372:ILE:O	1:B:376:LEU:HG	2.15	0.47
1:C:355:TYR:CD1	1:C:355:TYR:C	2.89	0.46
1:C:148:GLU:O	1:C:151:SER:HB2	2.16	0.45
1:A:263:GLY:HA3	1:C:246:ALA:O	2.16	0.45
1:B:413:ARG:NH1	3:B:609:HOH:O	2.45	0.45
1:C:189:PRO:HB3	1:C:198:ASN:HD21	1.81	0.45
1:C:108:ILE:O	1:C:112:THR:HG23	2.15	0.45
1:A:321:ILE:HG23	1:A:372:ILE:HD12	1.98	0.45
1:B:295:LYS:HG2	1:B:355:TYR:CD1	2.51	0.45
1:C:353:ASP:C	1:C:353:ASP:OD1	2.55	0.45
1:B:289:LYS:HD2	1:B:290:PHE:CE2	2.52	0.45
1:B:207:PRO:HD2	1:B:208:THR:HG23	1.98	0.44
1:C:229:ALA:HA	1:C:390:GLY:O	2.16	0.44
1:C:66:ILE:HD13	1:C:68:ARG:NH2	2.32	0.44
1:A:120:LYS:O	1:A:123:ASP:HB2	2.18	0.44
1:B:332:LEU:HA	3:B:660:HOH:O	2.17	0.44
1:B:189:PRO:HB3	1:B:198:ASN:HD21	1.82	0.44
1:A:250:LEU:HD13	1:A:399:TRP:CE3	2.53	0.44
2:C:501:CIT:O6	2:C:501:CIT:C5	2.66	0.44
1:C:75:GLU:HG3	1:C:320:ILE:HD11	2.00	0.44
1:C:311:TYR:O	1:C:362:TYR:OH	2.32	0.44
1:A:232:GLU:HG3	1:C:414:GLN:NE2	2.33	0.43
1:B:347:GLU:HG3	1:B:363:PRO:HG3	2.00	0.43
1:B:316:PRO:HA	1:B:319:LYS:HD3	2.00	0.43
1:C:350:ALA:HA	1:C:356:PHE:CD2	2.54	0.43
1:A:119:ARG:HD2	1:A:119:ARG:C	2.39	0.42
1:B:112:THR:HA	1:B:168:ALA:HB1	2.01	0.42
1:B:183:GLY:HA2	3:B:679:HOH:O	2.20	0.42
1:C:349:ALA:O	1:C:353:ASP:HB3	2.20	0.42
1:B:229:ALA:HA	1:B:390:GLY:O	2.19	0.42
1:C:270:ASN:OD1	1:C:273:VAL:HG23	2.20	0.42
1:C:173:ILE:O	1:C:177:THR:HG23	2.20	0.41
1:C:223:LYS:O	1:C:227:LEU:HG	2.21	0.41
1:A:280:ILE:HG23	1:A:290:PHE:CE1	2.56	0.41
1:C:268:GLY:HA2	1:C:366:ASP:OD2	2.21	0.41
1:C:308:HIS:CE1	2:C:501:CIT:H41	2.56	0.41
1:C:354:GLN:O	1:C:357:ILE:N	2.54	0.41
1:B:392:LEU:N	1:B:393:PRO:HD2	2.36	0.41

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:35:ARG:O	1:B:39:GLY:N	2.47	0.40
1:C:73:PRO:HG2	1:C:76:GLN:NE2	2.36	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	426/434 (98%)	411 (96%)	14 (3%)	1 (0%)	47 55
1	B	418/434 (96%)	400 (96%)	16 (4%)	2 (0%)	29 31
1	C	424/434 (98%)	391 (92%)	29 (7%)	4 (1%)	17 16
All	All	1268/1302 (97%)	1202 (95%)	59 (5%)	7 (1%)	25 26

All (7) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	287	THR
1	C	45	SER
1	C	301	PHE
1	C	269	ALA
1	A	297	LYS
1	B	303	LEU
1	C	289	LYS

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	360/364 (99%)	355 (99%)	5 (1%)	67	80
1	B	358/364 (98%)	343 (96%)	15 (4%)	30	38
1	C	358/364 (98%)	348 (97%)	10 (3%)	43	56
All	All	1076/1092 (98%)	1046 (97%)	30 (3%)	43	56

All (30) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	14	SER
1	A	140	CYS
1	A	232	GLU
1	A	387	PHE
1	A	407	GLU
1	B	4	THR
1	B	29	PHE
1	B	48	LYS
1	B	210	LYS
1	B	219	SER
1	B	231	HIS
1	B	232	GLU
1	B	265	LEU
1	B	274	ILE
1	B	282	LYS
1	B	302	ARG
1	B	330	GLN
1	B	353	ASP
1	B	411	ARG
1	B	431	LYS
1	C	203	MET
1	C	231	HIS
1	C	232	GLU
1	C	289	LYS
1	C	296	ASP
1	C	326	ASP
1	C	335	GLN
1	C	336	ASN
1	C	340	LEU
1	C	411	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such

sidechains are listed below:

Mol	Chain	Res	Type
1	B	354	GLN
1	C	154	GLN
1	C	270	ASN
1	C	271	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	CIT	C	501	-	12,12,12	1.12	0	17,17,17	0.95	0
2	CIT	B	501	1	12,12,12	0.96	0	17,17,17	1.61	6 (35%)
2	CIT	A	501	-	12,12,12	0.95	0	17,17,17	1.82	3 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	CIT	C	501	-	-	9/16/16/16	-
2	CIT	B	501	1	-	11/16/16/16	-
2	CIT	A	501	-	-	0/16/16/16	-

There are no bond length outliers.

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	CIT	O5-C6-C3	-5.02	115.15	122.25
2	B	501	CIT	O5-C6-C3	-3.10	117.86	122.25
2	A	501	CIT	O6-C6-C3	2.91	118.11	113.05
2	B	501	CIT	C3-C2-C1	2.37	119.56	113.81
2	A	501	CIT	C4-C3-C6	-2.29	105.19	110.11
2	B	501	CIT	O3-C5-C4	-2.14	116.68	122.94
2	B	501	CIT	O1-C1-C2	-2.12	116.76	122.94
2	B	501	CIT	C3-C4-C5	2.09	118.87	113.81
2	B	501	CIT	O2-C1-C2	2.04	120.91	114.35

There are no chirality outliers.

All (20) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	B	501	CIT	C2-C3-C4-C5
2	B	501	CIT	O7-C3-C4-C5
2	B	501	CIT	C6-C3-C4-C5
2	C	501	CIT	C1-C2-C3-O7
2	C	501	CIT	C1-C2-C3-C4
2	C	501	CIT	C1-C2-C3-C6
2	C	501	CIT	C2-C3-C6-O5
2	C	501	CIT	C2-C3-C6-O6
2	C	501	CIT	O7-C3-C6-O5
2	C	501	CIT	O7-C3-C6-O6
2	B	501	CIT	C2-C3-C6-O5
2	B	501	CIT	O7-C3-C6-O5
2	B	501	CIT	O7-C3-C6-O6
2	B	501	CIT	C2-C3-C6-O6
2	B	501	CIT	O1-C1-C2-C3
2	B	501	CIT	O2-C1-C2-C3
2	B	501	CIT	C4-C3-C6-O6
2	C	501	CIT	O7-C3-C4-C5
2	C	501	CIT	C6-C3-C4-C5
2	B	501	CIT	C4-C3-C6-O5


There are no ring outliers.

1 monomer is involved in 4 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	501	CIT	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	428/434 (98%)	0.02	26 (6%) 21 20	26, 42, 82, 108	0
1	B	424/434 (97%)	0.32	42 (9%) 7 6	25, 40, 96, 124	0
1	C	428/434 (98%)	0.44	39 (9%) 9 8	28, 49, 89, 113	0
All	All	1280/1302 (98%)	0.26	107 (8%) 11 9	25, 44, 89, 124	0

All (107) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	292	ALA	7.8
1	B	277	LEU	7.3
1	C	296	ASP	6.5
1	B	280	ILE	6.5
1	C	294	ALA	6.4
1	B	335	GLN	6.3
1	A	300	GLY	6.2
1	B	292	ALA	5.7
1	B	359	ARG	5.5
1	C	290	PHE	5.4
1	B	346	LEU	5.4
1	B	343	ALA	5.3
1	C	431	LYS	5.3
1	A	299	SER	5.2
1	B	287	THR	5.2
1	B	290	PHE	5.1
1	C	293	GLN	5.0
1	C	355	TYR	4.8
1	A	301	PHE	4.7
1	B	431	LYS	4.7
1	B	288	SER	4.7
1	B	291	ILE	4.5
1	C	354	GLN	4.5

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
1	B	281	GLU	4.5
1	B	358	ASP	4.5
1	B	283	ASP	4.5
1	B	301	PHE	4.5
1	A	302	ARG	4.4
1	C	357	ILE	4.4
1	B	339	LEU	4.2
1	C	356	PHE	4.1
1	C	333	GLY	4.0
1	A	297	LYS	3.8
1	B	333	GLY	3.8
1	C	346	LEU	3.8
1	B	13	GLN	3.7
1	B	336	ASN	3.7
1	A	282	LYS	3.6
1	B	357	ILE	3.6
1	B	302	ARG	3.6
1	C	358	ASP	3.6
1	B	276	MET	3.6
1	B	356	PHE	3.6
1	B	295	LYS	3.5
1	B	354	GLN	3.5
1	C	291	ILE	3.4
1	C	352	THR	3.4
1	C	359	ARG	3.4
1	C	283	ASP	3.3
1	A	285	GLY	3.3
1	A	298	ASN	3.3
1	C	153	ASN	3.2
1	C	279	ALA	3.2
1	C	300	GLY	3.1
1	B	282	LYS	3.1
1	B	279	ALA	3.1
1	A	213	ILE	3.1
1	A	296	ASP	3.1
1	A	153	ASN	3.0
1	C	98	GLU	3.0
1	B	1	MET	2.9
1	A	11	GLY	2.9
1	C	301	PHE	2.9
1	B	355	TYR	2.8
1	C	350	ALA	2.8

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
1	C	255	SER	2.8
1	A	428	THR	2.8
1	C	1	MET	2.8
1	B	330	GLN	2.7
1	C	295	LYS	2.7
1	C	287	THR	2.7
1	B	332	LEU	2.6
1	C	348	GLN	2.6
1	B	365	VAL	2.6
1	A	37	ALA	2.6
1	C	254	VAL	2.5
1	C	286	ASP	2.5
1	B	293	GLN	2.5
1	A	283	ASP	2.4
1	A	256	ALA	2.4
1	C	336	ASN	2.4
1	B	9	LEU	2.4
1	B	353	ASP	2.4
1	C	353	ASP	2.4
1	C	152	PRO	2.4
1	A	279	ALA	2.4
1	B	303	LEU	2.3
1	B	318	ALA	2.3
1	C	313	ASN	2.3
1	A	284	GLY	2.3
1	C	2	ALA	2.3
1	A	430	ARG	2.2
1	A	13	GLN	2.2
1	B	360	LYS	2.2
1	C	210	LYS	2.2
1	C	239	THR	2.2
1	A	9	LEU	2.2
1	C	351	LEU	2.2
1	B	320	ILE	2.2
1	A	365	VAL	2.1
1	B	316	PRO	2.1
1	A	425	VAL	2.1
1	A	243	VAL	2.1
1	B	152	PRO	2.0
1	C	94	HIS	2.0
1	A	420	THR	2.0
1	A	286	ASP	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	CIT	C	501	13/13	0.85	0.19	60,74,78,82	0
2	CIT	B	501	13/13	0.86	0.25	47,57,89,97	0
2	CIT	A	501	13/13	0.93	0.16	31,38,53,62	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




**Electron density around CIT C 501:**


$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around CIT B 501:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [i](#)

There are no such residues in this entry.