



# Full wwPDB NMR Structure Validation Report ⓘ

May 28, 2020 – 07:24 pm BST

PDB ID : 1E91  
Title : Structure of the complex of the Mad1-Sin3B interaction domains  
Authors : Spronk, C.A.E.M.; Tessari, M.; Kaan, A.M.; Jansen, J.F.A.; Vermeulen, M.;  
Stunnenberg, H.G.; Vuister, G.W.  
Deposited on : 2000-10-04

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)  
NmrClust : Kelley et al. (1996)  
MolProbity : 4.02b-467  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
RCI : v\_1n\_11\_5\_13\_A (Berjanski et al., 2005)  
PANAV : Wang et al. (2010)  
ShiftChecker : 2.11  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11



## 2 Ensemble composition and analysis i

This entry contains 20 models. Model 19 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:10-A:37, A:57-A:77, B:2-B:13 (61)	0.21	19

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 2 clusters and 1 single-model cluster was found.

Cluster number	Models
1	1, 2, 3, 4, 5, 6, 9, 10, 12, 13, 14, 15, 18, 19, 20
2	7, 8, 11, 17
Single-model clusters	16

### 3 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 1615 atoms, of which 795 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called PAIRED AMPHIPATHIC HELIX PROTEIN SIN3B.

Mol	Chain	Residues	Atoms					Trace	
			Total	C	H	N	O		S
1	A	85	1405	454	691	122	137	1	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	83	ALA	GLY	conflict	UNP Q62141

- Molecule 2 is a protein called MAD PROTEIN (MAX DIMERIZER).


Mol	Chain	Residues	Atoms					Trace	
			Total	C	H	N	O		S
2	B	13	210	67	104	15	23	1	0

## 4 Residue-property plots


### 4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: PAIRED AMPHIPATHIC HELIX PROTEIN SIN3B



- Molecule 2: MAD PROTEIN (MAX DIMERIZER)




### 4.2 Scores per residue for each member of the ensemble


Colouring as in section 4.1 above.

#### 4.2.1 Score per residue for model 1

- Molecule 1: PAIRED AMPHIPATHIC HELIX PROTEIN SIN3B




- Molecule 2: MAD PROTEIN (MAX DIMERIZER)




### 4.2.2 Score per residue for model 2

- Molecule 1: PAIRED AMPHIPATHIC HELIX PROTEIN SIN3B




- Molecule 2: MAD PROTEIN (MAX DIMERIZER)




### 4.2.3 Score per residue for model 3

- Molecule 1: PAIRED AMPHIPATHIC HELIX PROTEIN SIN3B




- Molecule 2: MAD PROTEIN (MAX DIMERIZER)




### 4.2.4 Score per residue for model 4

- Molecule 1: PAIRED AMPHIPATHIC HELIX PROTEIN SIN3B




- Molecule 2: MAD PROTEIN (MAX DIMERIZER)




### 4.2.5 Score per residue for model 5

- Molecule 1: PAIRED AMPHIPATHIC HELIX PROTEIN SIN3B




- Molecule 2: MAD PROTEIN (MAX DIMERIZER)




### 4.2.6 Score per residue for model 6

- Molecule 1: PAIRED AMPHIPATHIC HELIX PROTEIN SIN3B




- Molecule 2: MAD PROTEIN (MAX DIMERIZER)




### 4.2.7 Score per residue for model 7

- Molecule 1: PAIRED AMPHIPATHIC HELIX PROTEIN SIN3B




- Molecule 2: MAD PROTEIN (MAX DIMERIZER)




### 4.2.8 Score per residue for model 8

- Molecule 1: PAIRED AMPHIPATHIC HELIX PROTEIN SIN3B




- Molecule 2: MAD PROTEIN (MAX DIMERIZER)




### 4.2.9 Score per residue for model 9

- Molecule 1: PAIRED AMPHIPATHIC HELIX PROTEIN SIN3B




- Molecule 2: MAD PROTEIN (MAX DIMERIZER)




### 4.2.10 Score per residue for model 10

- Molecule 1: PAIRED AMPHIPATHIC HELIX PROTEIN SIN3B




- Molecule 2: MAD PROTEIN (MAX DIMERIZER)






#### 4.2.11 Score per residue for model 11

- Molecule 1: PAIRED AMPHIPATHIC HELIX PROTEIN SIN3B




- Molecule 2: MAD PROTEIN (MAX DIMERIZER)




#### 4.2.12 Score per residue for model 12

- Molecule 1: PAIRED AMPHIPATHIC HELIX PROTEIN SIN3B




- Molecule 2: MAD PROTEIN (MAX DIMERIZER)




#### 4.2.13 Score per residue for model 13

- Molecule 1: PAIRED AMPHIPATHIC HELIX PROTEIN SIN3B




- Molecule 2: MAD PROTEIN (MAX DIMERIZER)




#### 4.2.14 Score per residue for model 14

- Molecule 1: PAIRED AMPHIPATHIC HELIX PROTEIN SIN3B




- Molecule 2: MAD PROTEIN (MAX DIMERIZER)




#### 4.2.15 Score per residue for model 15

- Molecule 1: PAIRED AMPHIPATHIC HELIX PROTEIN SIN3B




- Molecule 2: MAD PROTEIN (MAX DIMERIZER)




#### 4.2.16 Score per residue for model 16

- Molecule 1: PAIRED AMPHIPATHIC HELIX PROTEIN SIN3B




- Molecule 2: MAD PROTEIN (MAX DIMERIZER)




#### 4.2.17 Score per residue for model 17

- Molecule 1: PAIRED AMPHIPATHIC HELIX PROTEIN SIN3B




- Molecule 2: MAD PROTEIN (MAX DIMERIZER)




#### 4.2.18 Score per residue for model 18

- Molecule 1: PAIRED AMPHIPATHIC HELIX PROTEIN SIN3B




- Molecule 2: MAD PROTEIN (MAX DIMERIZER)




#### 4.2.19 Score per residue for model 19 (medoid)

- Molecule 1: PAIRED AMPHIPATHIC HELIX PROTEIN SIN3B




- Molecule 2: MAD PROTEIN (MAX DIMERIZER)



#### 4.2.20 Score per residue for model 20

- Molecule 1: PAIRED AMPHIPATHIC HELIX PROTEIN SIN3B

Chain A:  58% 42%



- Molecule 2: MAD PROTEIN (MAX DIMERIZER)

Chain B:  92% 8%



## 5 Refinement protocol and experimental data overview

The models were refined using the following method: *SIMULATED ANNEALING, ITERATIVE NOE-ASSIGNMENT*.

Of the 100 calculated structures, 20 were deposited, based on the following criterion: *30 STRUCTURES WITH NO RESTRAINT VIOLATIONS, 20 LOWEST ENERGY STRUCTURES FINALLY SELECTED*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CNS	refinement	1.0
ARIA	structure solution	ARIA

No chemical shift data was provided. No validations of the models with respect to experimental NMR restraints is performed at this time.

## 6 Model quality [i](#)

### 6.1 Standard geometry [i](#)

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
All	All	10160	10000	0	-

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is -.

There are no clashes.

### 6.3 Torsion angles [i](#)

#### 6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	0	-	-	-	-
2	B	0	-	-	-	-
All	All	0	-	-	-	-

There are no Ramachandran outliers.

### 6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	0	-	-	-
2	B	0	-	-	-
All	All	0	-	-	-

There are no protein residues with a non-rotameric sidechain to report.

### 6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 6.7 Other polymers [i](#)

There are no such molecules in this entry.

### 6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 7 Chemical shift validation

No chemical shift data were provided