



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 15, 2024 – 09:39 AM EDT

PDB ID : 2EBF
Title : Crystal structures reveal a thiol-protease like catalytic triad in the C-terminal region of Pasteurella multocida toxin
Authors : Kitadokoro, K.; Horiguchi, Y.; Kamitani, S.
Deposited on : 2007-02-08
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 1.20.1
EDS : 2.37.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	X	746	 10% (Poor fit), 81% (0 outliers), 13% (1 outlier), 5% (2 outliers), 5% (3+ outliers)
2	A	2	 50% (0 outliers), 50% (1 outlier)
2	B	2	 50% (0 outliers), 50% (1 outlier)

2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 6450 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Dermonecrotic toxin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	X	711	5643	3603	939	1074	27	0	0	0

There are 29 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
X	540	MET	-	expression tag	UNP P17452
X	541	GLY	-	expression tag	UNP P17452
X	542	HIS	-	expression tag	UNP P17452
X	543	HIS	-	expression tag	UNP P17452
X	544	HIS	-	expression tag	UNP P17452
X	545	HIS	-	expression tag	UNP P17452
X	546	HIS	-	expression tag	UNP P17452
X	547	HIS	-	expression tag	UNP P17452
X	548	ASP	-	expression tag	UNP P17452
X	549	TYR	-	expression tag	UNP P17452
X	550	ASP	-	expression tag	UNP P17452
X	551	ILE	-	expression tag	UNP P17452
X	552	PRO	-	expression tag	UNP P17452
X	553	THR	-	expression tag	UNP P17452
X	554	THR	-	expression tag	UNP P17452
X	555	GLU	-	expression tag	UNP P17452
X	556	ASN	-	expression tag	UNP P17452
X	557	LEU	-	expression tag	UNP P17452
X	558	TYR	-	expression tag	UNP P17452
X	559	PHE	-	expression tag	UNP P17452
X	560	GLN	-	expression tag	UNP P17452
X	561	GLY	-	expression tag	UNP P17452
X	562	ALA	-	expression tag	UNP P17452
X	563	HIS	-	expression tag	UNP P17452
X	564	MET	-	expression tag	UNP P17452
X	565	GLY	-	expression tag	UNP P17452
X	566	ILE	-	expression tag	UNP P17452

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
X	567	GLN	-	expression tag	UNP P17452
X	568	ARG	-	expression tag	UNP P17452

- Molecule 2 is an oligosaccharide called alpha-D-glucopyranose-(1-1)-alpha-D-glucopyranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
2	A	2	Total	C	O	0	0	0
			23	12	11			
2	B	2	Total	C	O	0	0	0
			23	12	11			

- Molecule 3 is PHOSPHONATE (three-letter code: 2PO) (formula: HO₃P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	X	1	Total	O	P	0	0
			4	3	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	X	757	Total	O	0	0
			757	757		

3 Residue-property plots i

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Dermonecrotic toxin



- Molecule 2: alpha-D-glucopyranose-(1-1)-alpha-D-glucopyranose



- Molecule 2: alpha-D-glucopyranose-(1-1)-alpha-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	110.97Å 150.41Å 77.14Å 90.00° 105.46° 90.00°	Depositor
Resolution (Å)	20.00 – 1.90 19.99 – 1.90	Depositor EDS
% Data completeness (in resolution range)	97.5 (20.00-1.90) 97.5 (19.99-1.90)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.08 (at 1.90Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.205 , 0.233 0.205 , 0.234	Depositor DCC
R_{free} test set	4662 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	30.2	Xtrriage
Anisotropy	0.046	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 47.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	6450	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.62% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 2PO, GLC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	X	0.44	0/5769	0.59	0/7811

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	X	5643	0	5573	74	0
2	A	23	0	21	0	0
2	B	23	0	21	0	0
3	X	4	0	0	0	0
4	X	757	0	0	11	0
All	All	6450	0	5615	74	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (74) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:X:850:THR:HB	4:X:2141:HOH:O	1.55	1.02
1:X:589:GLY:HA3	1:X:1284:ALA:HB3	1.52	0.91
1:X:850:THR:HG23	1:X:853:PHE:H	1.35	0.90
1:X:1027:ILE:HD11	1:X:1062:TYR:HB3	1.56	0.88
1:X:850:THR:HG21	4:X:1497:HOH:O	1.75	0.84
1:X:1223:HIS:HD2	1:X:1250:ARG:HH12	1.28	0.79
1:X:589:GLY:HA3	1:X:1284:ALA:CB	2.16	0.75
1:X:759:TYR:HA	1:X:762:LEU:HD23	1.68	0.75
1:X:784:SER:HA	4:X:2137:HOH:O	1.87	0.74
1:X:1103:PRO:HB2	1:X:1104:GLN:HG2	1.73	0.70
1:X:809:MET:SD	4:X:2062:HOH:O	2.51	0.69
1:X:916:ASN:ND2	1:X:943:ASP:OD2	2.25	0.68
1:X:843:GLY:H	1:X:846:GLN:HE21	1.43	0.65
1:X:825:GLU:HG3	1:X:826:LEU:H	1.62	0.64
1:X:1001:HIS:HD2	4:X:1425:HOH:O	1.80	0.63
1:X:575:LEU:HD21	4:X:1808:HOH:O	2.01	0.61
1:X:784:SER:HB3	1:X:789:MET:O	2.02	0.60
1:X:591:TRP:H	1:X:663:GLN:HE22	1.48	0.60
1:X:749:ILE:HD11	1:X:800:LEU:HB2	1.83	0.60
1:X:1104:GLN:O	1:X:1257:PHE:CZ	2.55	0.60
1:X:804:HIS:HB2	1:X:809:MET:CE	2.32	0.59
1:X:577:GLY:HA2	4:X:2095:HOH:O	2.03	0.59
1:X:589:GLY:CA	1:X:1284:ALA:HB3	2.30	0.58
1:X:861:ARG:HD3	1:X:1003:ASP:OD1	2.04	0.57
1:X:1060:VAL:HG21	1:X:1067:ALA:HB2	1.86	0.57
1:X:1113:PHE:O	1:X:1252:ARG:HA	2.05	0.57
1:X:749:ILE:HD11	1:X:800:LEU:CB	2.35	0.57
1:X:591:TRP:H	1:X:663:GLN:NE2	2.02	0.57
1:X:804:HIS:HB2	1:X:809:MET:HE3	1.87	0.56
1:X:575:LEU:HD22	1:X:1112:GLN:H	1.71	0.56
1:X:814:SER:HB2	1:X:815:PRO:HD2	1.88	0.55
1:X:1223:HIS:CD2	1:X:1250:ARG:HH12	2.17	0.55
1:X:575:LEU:HB2	1:X:1111:LYS:HB2	1.90	0.54
1:X:1104:GLN:HB2	1:X:1259:GLU:HG2	1.90	0.54
1:X:1162:ALA:HA	1:X:1225:GLN:HE21	1.72	0.53
1:X:1104:GLN:HA	4:X:2147:HOH:O	2.09	0.52
1:X:1010:ASN:C	1:X:1010:ASN:HD22	2.13	0.52
1:X:970:LEU:HD11	1:X:986:LEU:HD12	1.92	0.51
1:X:1285:LEU:H	1:X:1285:LEU:HD13	1.74	0.51
1:X:814:SER:H	1:X:818:ASN:HD22	1.58	0.50
1:X:637:LEU:HD13	1:X:664:LEU:HD11	1.93	0.50
1:X:807:GLU:HG3	1:X:1006:VAL:HG23	1.92	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:X:850:THR:CG2	1:X:853:PHE:H	2.17	0.49
1:X:631:GLN:HE22	1:X:692:ARG:HA	1.78	0.49
1:X:752:PHE:O	1:X:755:ILE:HG12	2.13	0.49
1:X:1060:VAL:HG22	1:X:1065:ILE:HG13	1.95	0.48
1:X:593:PRO:HG2	1:X:619:GLN:NE2	2.28	0.48
1:X:1223:HIS:HD2	1:X:1250:ARG:NH1	2.05	0.48
1:X:1218:ALA:O	1:X:1237:ILE:HA	2.15	0.47
1:X:915:GLN:HB2	1:X:1069:VAL:HG12	1.97	0.47
1:X:1069:VAL:HG13	1:X:1079:LEU:HG	1.96	0.46
1:X:1060:VAL:HG13	1:X:1065:ILE:O	2.16	0.46
1:X:738:ILE:HD12	1:X:796:ILE:HD12	1.98	0.45
1:X:843:GLY:H	1:X:846:GLN:NE2	2.13	0.45
1:X:748:ASP:O	1:X:751:VAL:HG23	2.18	0.44
1:X:613:ARG:NE	4:X:1832:HOH:O	2.41	0.44
1:X:759:TYR:CE1	1:X:834:MET:HB3	2.52	0.44
1:X:915:GLN:HG2	1:X:916:ASN:N	2.32	0.44
1:X:1011:ASN:ND2	1:X:1013:TYR:H	2.16	0.43
1:X:1050:PRO:HA	1:X:1087:VAL:O	2.19	0.43
1:X:583:PHE:O	1:X:585:ILE:HD12	2.19	0.43
1:X:1245:LEU:C	1:X:1245:LEU:HD23	2.39	0.43
1:X:1071:ASP:HB3	1:X:1073:ASP:H	1.84	0.42
1:X:749:ILE:HG22	1:X:774:SER:HB2	2.01	0.42
1:X:941:ILE:HD12	1:X:996:PHE:CE2	2.54	0.42
1:X:813:ILE:HA	1:X:818:ASN:HD21	1.85	0.42
1:X:941:ILE:HD12	1:X:996:PHE:CZ	2.55	0.42
1:X:804:HIS:HB2	1:X:809:MET:HE2	2.02	0.41
1:X:745:VAL:HG23	4:X:2141:HOH:O	2.19	0.41
1:X:639:GLU:OE2	1:X:851:ARG:NH2	2.44	0.41
1:X:699:MET:HA	1:X:702:MET:HE3	2.02	0.41
1:X:1245:LEU:HD22	4:X:1904:HOH:O	2.19	0.41
1:X:602:VAL:HB	1:X:605:LYS:HB3	2.03	0.40
1:X:1004:ASN:HD22	1:X:1004:ASN:HA	1.67	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	X	709/746 (95%)	686 (97%)	20 (3%)	3 (0%)	34 24

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	X	916	ASN
1	X	980	GLY
1	X	1071	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	X	616/646 (95%)	589 (96%)	27 (4%)	28 19

All (27) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	X	596	LEU
1	X	597	LYS
1	X	618	LEU
1	X	627	LEU
1	X	631	GLN
1	X	637	LEU
1	X	668	LEU
1	X	738	ILE
1	X	851	ARG
1	X	852	ASN
1	X	916	ASN
1	X	919	LYS
1	X	941	ILE
1	X	955	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	X	985	LEU
1	X	986	LEU
1	X	1004	ASN
1	X	1010	ASN
1	X	1033	GLU
1	X	1079	LEU
1	X	1115	LEU
1	X	1133	VAL
1	X	1138	ARG
1	X	1139	GLU
1	X	1181	LEU
1	X	1283	ARG
1	X	1285	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (24) such sidechains are listed below:

Mol	Chain	Res	Type
1	X	619	GLN
1	X	631	GLN
1	X	663	GLN
1	X	798	ASN
1	X	818	ASN
1	X	839	GLN
1	X	846	GLN
1	X	867	ASN
1	X	907	ASN
1	X	916	ASN
1	X	926	ASN
1	X	934	ASN
1	X	951	GLN
1	X	972	ASN
1	X	1001	HIS
1	X	1004	ASN
1	X	1010	ASN
1	X	1011	ASN
1	X	1021	ASN
1	X	1131	ASN
1	X	1203	GLN
1	X	1223	HIS
1	X	1225	GLN
1	X	1266	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GLC	A	1	2	11,11,12	0.35	0	15,15,17	1.22	1 (6%)
2	GLC	A	2	2	12,12,12	0.60	0	17,17,17	0.68	0
2	GLC	B	1	2	11,11,12	0.45	0	15,15,17	0.91	1 (6%)
2	GLC	B	2	2	12,12,12	0.52	0	17,17,17	0.60	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	A	1	2	-	0/2/19/22	0/1/1/1
2	GLC	A	2	2	-	0/2/22/22	0/1/1/1
2	GLC	B	1	2	-	0/2/19/22	0/1/1/1
2	GLC	B	2	2	-	0/2/22/22	0/1/1/1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1	GLC	C1-O5-C5	4.16	117.77	112.19

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1	GLC	C1-O5-C5	3.00	116.21	112.19


There are no chirality outliers.


There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	2PO	X	1327	-	0,3,3	-	-	0,3,3	-	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	X	711/746 (95%)	0.57	76 (10%) 6 6	24, 33, 51, 64	0

All (76) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	X	979	ASN	9.0
1	X	1231	LEU	9.0
1	X	576	GLY	7.9
1	X	575	LEU	6.7
1	X	1285	LEU	6.6
1	X	1283	ARG	6.2
1	X	1284	ALA	5.5
1	X	577	GLY	5.4
1	X	958	GLN	5.1
1	X	1033	GLU	5.1
1	X	603	ILE	5.0
1	X	788	GLY	4.9
1	X	762	LEU	4.9
1	X	786	SER	4.8
1	X	769	ASN	4.8
1	X	771	GLY	4.8
1	X	975	LYS	4.6
1	X	780	LEU	4.6
1	X	756	GLN	4.5
1	X	783	GLU	4.4
1	X	1281	LEU	4.3
1	X	740	PHE	4.3
1	X	827	TYR	4.1
1	X	760	TRP	3.8
1	X	782	ARG	3.8
1	X	829	PHE	3.8
1	X	1072	SER	3.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	X	951	GLN	3.6
1	X	874	SER	3.5
1	X	787	LYS	3.4
1	X	815	PRO	3.4
1	X	739	ASP	3.4
1	X	641	ILE	3.2
1	X	959	GLY	3.2
1	X	961	ASN	3.2
1	X	1135	GLU	3.0
1	X	1103	PRO	2.9
1	X	1010	ASN	2.8
1	X	1073	ASP	2.8
1	X	1117	ALA	2.8
1	X	647	SER	2.8
1	X	722	VAL	2.8
1	X	968	ALA	2.7
1	X	736	GLU	2.7
1	X	824	TYR	2.7
1	X	1104	GLN	2.6
1	X	831	PHE	2.6
1	X	1282	THR	2.6
1	X	800	LEU	2.6
1	X	768	TRP	2.5
1	X	1031	ASP	2.5
1	X	755	ILE	2.5
1	X	1032	ARG	2.4
1	X	970	LEU	2.4
1	X	1136	ASP	2.4
1	X	652	GLU	2.4
1	X	850	THR	2.4
1	X	993	ASN	2.4
1	X	960	SER	2.3
1	X	1034	GLY	2.2
1	X	606	PRO	2.2
1	X	637	LEU	2.2
1	X	660	LEU	2.2
1	X	657	PHE	2.2
1	X	745	VAL	2.2
1	X	1207	THR	2.2
1	X	675	ALA	2.1
1	X	962	GLU	2.1
1	X	841	ASP	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	X	1017	TYR	2.1
1	X	1219	VAL	2.1
1	X	757	THR	2.1
1	X	826	LEU	2.1
1	X	955	GLN	2.0
1	X	605	LYS	2.0
1	X	1236	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	GLC	A	1	11/12	0.85	0.27	46,47,48,48	0
2	GLC	A	2	12/12	0.86	0.27	40,45,47,47	0
2	GLC	B	2	12/12	0.91	0.22	44,45,45,46	0
2	GLC	B	1	11/12	0.94	0.22	41,44,45,45	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around Chain A:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	2PO	X	1327	4/4	0.63	0.24	50,51,51,52	4

6.5 Other polymers [i](#)

There are no such residues in this entry.