

wwPDB X-ray Structure Validation Summary Report (i)

Mar 23, 2024 – 10:49 PM EDT

PDB ID : 1ECZ

Title: PROTEASE INHIBITOR ECOTIN

Authors: Shin, D.H.; Suh, S.W.

Deposited on : 1996-08-06

Resolution : 2.68 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.36.1 buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac : 5.8.0158

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.36.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$

The reported resolution of this entry is 2.68 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive	Similar resolution
1.133113	$(\# \mathrm{Entries})$	$(\# ext{Entries}, ext{ resolution range}(ext{Å}))$
Clashscore	141614	4210 (2.70-2.66)
Ramachandran outliers	138981	4141 (2.70-2.66)
Sidechain outliers	138945	4141 (2.70-2.66)
RSRZ outliers	127900	3780 (2.70-2.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain				
1	A	142	49%	46%	-		
1	В	142	51%	44%	•		

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	BOG	A	1001	X	-	-	-
2	BOG	A	1009	X	-	-	-





Continued from previous page...

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	BOG	A	1011	X	-	-	-
2	BOG	A	1015	X	-	-	-
2	BOG	A	1019	X	-	-	-
2	BOG	A	1021	X	-	-	-
2	BOG	A	1025	X	-	-	-
2	BOG	A	1029	X	-	-	-
2	BOG	A	1033	X	-	-	-
2	BOG	A	1041	X	-	-	-
2	BOG	В	1005	X	-	-	-
2	BOG	В	1007	X	-	=	-
2	BOG	В	1013	X	-	-	-
2	BOG	В	1017	X	-	=	-
2	BOG	В	1023	X	-	-	-
2	BOG	В	1027	X	-	-	-
2	BOG	В	1031	X	-	-	-
2	BOG	В	1035	X	-	X	-
2	BOG	В	1037	X	-	-	-
2	BOG	В	1039	X	-	-	-



2 Entry composition (i)


There are 3 unique types of molecules in this entry. The entry contains 2767 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called ECOTIN.

\mathbf{Mol}	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	A	142	Total			О	S	0	0	0
1	71	1 12	1130	721	187	216	6		O	U
1	D	142	Total	С	N	Ο	S	0	0	0
1	Ъ	142	1130	721	187	216	6	0	U	U

• Molecule 2 is octyl beta-D-glucopyranoside (three-letter code: BOG) (formula: C₁₄H₂₈O₆).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 20 14 6	0	0
2	A	1	Total C O 20 14 6	0	0
2	A	1	Total C O 20 14 6	0	0
2	A	1	Total C O 20 14 6	0	0

Continued on next page...



 $Continued\ from\ previous\ page...$

Mol	Chain	$oxed{ ext{Residues} }$	Atoms	ZeroOcc	AltConf
2	A	1	Total C O	0	0
			20 14 6		
2	A	1	Total C O 20 14 6	0	0
			Total C O		
2	A	1	20 14 6	0	0
			Total C O		
2	A	1	20 14 6	0	0
2	A	1	Total C O	0	0
2	A	1	20 14 6	0	U
2	A	1	Total C O	0	0
	11	1	20 14 6	0	Ů,
2	A	1	Total C O	0	0
			12 6 6 Total C O		
2	В	1	Total C O 20 14 6	0	0
			Total C O		
2	В	1	20 14 6	0	0
	D	1	Total C O	0	0
2	В	1	20 14 6	0	0
2	В	1	Total C O	0	0
	Ъ	1	20 14 6	U	U
2	В	1	Total C O	0	0
		_	20 14 6	Ů	
2	В	1	Total C O	0	0
			20 14 6 Total C O		
2	В	1	20 14 6	0	0
			Total C O		
2	В	1	20 14 6	0	0
0	D	1	Total C O	0	0
2	В	1	20 14 6	0	0
2	В	1	Total C O	0	0
	ע	1	20 14 6		


• Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	49	Total O 49 49	0	0
3	В	46	Total O 46 46	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	39.22Å 84.86Å 98.74Å	Donositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	6.00 - 2.68	Depositor
Resolution (A)	30.69 - 2.69	EDS
% Data completeness	78.1 (6.00-2.68)	Depositor
(in resolution range)	80.4 (30.69-2.69)	EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	2.27 (at 2.68Å)	Xtriage
Refinement program	X-PLOR 3.1	Depositor
D D.	0.195 , 0.309	Depositor
R, R_{free}	0.229 , (Not available)	DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	45.3	Xtriage
Anisotropy	0.299	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.28 , 148.2	EDS
L-test for twinning ²	$ < L > = 0.46, < L^2> = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.88	EDS
Total number of atoms	2767	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The analyses of the Patterson function reveals a significant off-origin peak that is 36.90 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 4.6568e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: BOG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	A	0.35	0/1152	0.63	0/1558	
1	В	0.35	0/1152	0.63	1/1558 (0.1%)	
All	All	0.35	0/2304	0.63	1/3116 (0.0%)	

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^{o})$	$\operatorname{Ideal}(^{o})$
1	В	5	GLN	N-CA-C	5.50	125.86	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1130	0	1144	72	0
1	В	1130	0	1146	77	0
2	A	212	0	291	38	0
2	В	200	0	280	30	0
3	A	49	0	0	3	0
3	В	46	0	0	3	0
All	All	2767	0	2861	175	0



The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 32.

The worst 5 of 175 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$egin{aligned} & ext{Interatomic} \ & ext{distance} \ & ext{(Å)} \end{aligned}$	$\begin{array}{c} \text{Clash} \\ \text{overlap (Å)} \end{array}$
2:B:1035:BOG:C3	2:B:1035:BOG:H1'2	1.70	1.19
2:A:1015:BOG:H2'1	2:A:1015:BOG:H5	1.20	1.18
2:A:1021:BOG:H1'1	2:A:1021:BOG:H5	1.09	1.08
1:B:112:LYS:HD3	2:B:1013:BOG:H8'3	1.40	1.02
2:B:1035:BOG:H3	2:B:1035:BOG:C1'	1.89	1.02

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentile	es
1	A	140/142 (99%)	113 (81%)	23 (16%)	4 (3%)	4 9	
1	В	140/142 (99%)	114 (81%)	21 (15%)	5 (4%)	3 6	
All	All	280/284 (99%)	227 (81%)	44 (16%)	9 (3%)	4 8	

5 of 9 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	5	GLN
1	В	5	GLN
1	A	3	SER
1	В	3	SER
1	В	96	PHE



5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Perce	ntiles
1	A	125/125 (100%)	122 (98%)	3 (2%)	49	75
1	В	125/125 (100%)	120 (96%)	5 (4%)	31	57
All	All	250/250 (100%)	242 (97%)	8 (3%)	39	65

5 of 8 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	В	117	VAL
1	В	72	TYR
1	В	52	LEU
1	В	38	VAL
1	В	70	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 6 such sidechains are listed below:

Mol	Chain	Res	Type
1	В	30	GLN
1	В	44	GLN
1	В	53	HIS
1	A	44	GLN
1	A	30	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.



5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

21 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Т	Clasia	Das	T :1-	Во	ond leng	ths	В	ond ang	les
Mol	Type	Chain	Res	Link	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	BOG	В	1017	-	20,20,20	0.47	0	25,25,25	0.59	0
2	BOG	A	1029	-	20,20,20	0.41	0	25,25,25	0.54	0
2	BOG	A	1003	_	20,20,20	0.46	0	$25,\!25,\!25$	0.77	2 (8%)
2	BOG	В	1023	-	20,20,20	0.47	0	25,25,25	0.65	0
2	BOG	A	1033	-	20,20,20	0.49	0	25,25,25	0.66	0
2	BOG	В	1007	-	20,20,20	0.42	0	25,25,25	0.59	0
2	BOG	A	1009	-	20,20,20	0.43	0	25,25,25	0.54	0
2	BOG	A	1011	-	20,20,20	0.51	0	25,25,25	0.59	0
2	BOG	В	1039	_	20,20,20	0.49	0	25,25,25	0.77	0
2	BOG	A	1019	-	20,20,20	0.43	0	25,25,25	0.64	0
2	BOG	В	1037	_	20,20,20	0.45	0	$25,\!25,\!25$	0.60	0
2	BOG	A	1021	_	20,20,20	0.45	0	$25,\!25,\!25$	0.80	2 (8%)
2	BOG	В	1013	-	20,20,20	0.45	0	25,25,25	0.59	0
2	BOG	В	1027	-	20,20,20	0.48	0	25,25,25	0.60	0
2	BOG	В	1005	-	20,20,20	0.49	0	25,25,25	0.56	0
2	BOG	В	1031	_	20,20,20	0.49	0	25,25,25	0.60	0
2	BOG	В	1035	-	20,20,20	0.44	0	$25,\!25,\!25$	0.88	2 (8%)
2	BOG	A	1025	-	20,20,20	0.45	0	25,25,25	0.64	0
2	BOG	A	1041	-	12,12,20	0.31	0	17,17,25	0.56	0
2	BOG	A	1001	-	20,20,20	0.50	0	25,25,25	0.62	0
2	BOG	A	1015	-	20,20,20	0.47	0	25,25,25	0.78	1 (4%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BOG	В	1017	-	1/1/5/5	5/11/31/31	0/1/1/1
2	BOG	A	1029	-	1/1/5/5	5/11/31/31	0/1/1/1
2	BOG	A	1003	-	-	6/11/31/31	0/1/1/1
2	BOG	В	1023	-	1/1/5/5	6/11/31/31	0/1/1/1
2	BOG	A	1033	-	1/1/5/5	8/11/31/31	0/1/1/1
2	BOG	В	1007	-	1/1/5/5	8/11/31/31	0/1/1/1
2	BOG	A	1009	-	1/1/5/5	5/11/31/31	0/1/1/1
2	BOG	A	1011	-	1/1/5/5	6/11/31/31	0/1/1/1
2	BOG	В	1039	-	1/1/5/5	9/11/31/31	0/1/1/1
2	BOG	A	1019	-	1/1/5/5	6/11/31/31	0/1/1/1
2	BOG	В	1037	-	1/1/5/5	7/11/31/31	0/1/1/1
2	BOG	A	1021	-	1/1/5/5	4/11/31/31	0/1/1/1
2	BOG	В	1013	-	1/1/5/5	4/11/31/31	0/1/1/1
2	BOG	В	1027	-	1/1/5/5	8/11/31/31	0/1/1/1
2	BOG	В	1005	-	1/1/5/5	7/11/31/31	0/1/1/1
2	BOG	В	1031	-	1/1/5/5	4/11/31/31	0/1/1/1
2	BOG	В	1035	-	1/1/5/5	5/11/31/31	0/1/1/1
2	BOG	A	1025	-	1/1/5/5	8/11/31/31	0/1/1/1
2	BOG	A	1041	-	1/1/5/5	0/2/22/31	0/1/1/1
2	BOG	A	1001	-	1/1/5/5	8/11/31/31	0/1/1/1
2	BOG	A	1015	-	1/1/5/5	5/11/31/31	0/1/1/1

There are no bond length outliers.

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$Ideal(^{o})$
2	В	1035	BOG	C1'-O1-C1	2.97	118.76	113.84
2	A	1003	BOG	C1'-O1-C1	2.64	118.22	113.84
2	A	1021	BOG	C1'-O1-C1	2.63	118.20	113.84
2	В	1035	BOG	O1-C1-C2	2.55	112.28	108.30
2	A	1021	BOG	O1-C1-C2	2.17	111.69	108.30

5 of 20 chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	A	1001	BOG	C1
2	A	1009	BOG	C1
2	A	1011	BOG	C1

Continued on next page...



Continued from previous page...

Mol	Chain	Res	Type	Atom
2	A	1015	BOG	C1
2	A	1019	BOG	C1

5 of 124 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1003	BOG	C2-C1-O1-C1'
2	A	1009	BOG	O5-C1-O1-C1'
2	A	1011	BOG	C2-C1-O1-C1'
2	A	1011	BOG	O5-C1-O1-C1'
2	A	1011	BOG	C2'-C1'-O1-C1

There are no ring outliers.


18 monomers are involved in 68 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1029	BOG	3	0
2	В	1023	BOG	4	0
2	A	1033	BOG	3	0
2	В	1007	BOG	1	0
2	A	1009	BOG	1	0
2	A	1011	BOG	8	0
2	В	1039	BOG	4	0
2	A	1019	BOG	1	0
2	В	1037	BOG	2	0
2	A	1021	BOG	7	0
2	В	1013	BOG	3	0
2	В	1005	BOG	3	0
2	В	1031	BOG	2	0
2	В	1035	BOG	11	0
2	A	1025	BOG	3	0
2	A	1041	BOG	2	0
2	A	1001	BOG	2	0
2	A	1015	BOG	8	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring




in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







Torsions


Rings































































5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	# RSRZ > 2		$OWAB(A^2)$	Q < 0.9
1	A	142/142 (100%)	-0.21	3 (2%) 63	63	7, 35, 70, 81	0
1	В	142/142 (100%)	-0.13	2 (1%) 75	76	11, 36, 68, 90	0
All	All	284/284 (100%)	-0.17	5 (1%) 68	69	7, 35, 70, 90	0

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	В	3	SER	3.3
1	A	12	PRO	2.6
1	В	89	ASP	2.5
1	A	4	VAL	2.0
1	A	2	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.



Mol	Type	Chain	Res	Atoms	RSCC	RSR	${f B ext{-}factors}({f \AA}^2)$	Q<0.9
2	BOG	A	1021	20/20	0.64	0.40	17,71,89,97	0
2	BOG	В	1005	20/20	0.66	0.32	31,69,89,95	0
2	BOG	В	1027	20/20	0.70	0.29	30,47,65,73	0
2	BOG	A	1001	20/20	0.71	0.23	10,60,74,82	0
2	BOG	A	1033	20/20	0.73	0.36	28,68,81,83	0
2	BOG	В	1017	20/20	0.74	0.38	21,56,66,70	0
2	BOG	A	1009	20/20	0.76	0.25	16,51,69,74	0
2	BOG	В	1037	20/20	0.76	0.36	26,55,78,85	0
2	BOG	В	1013	20/20	0.77	0.28	35,50,87,91	0
2	BOG	В	1035	20/20	0.78	0.26	10,56,72,84	0
2	BOG	A	1029	20/20	0.79	0.35	2,78,97,99	0
2	BOG	A	1015	20/20	0.80	0.29	11,47,79,84	0
2	BOG	В	1031	20/20	0.80	0.23	9,65,83,86	0
2	BOG	В	1023	20/20	0.81	0.27	9,44,80,90	0
2	BOG	В	1039	20/20	0.81	0.28	18,42,69,70	0
2	BOG	A	1003	20/20	0.82	0.20	15,50,66,75	0
2	BOG	A	1019	20/20	0.83	0.25	2,41,59,65	0
2	BOG	A	1025	20/20	0.84	0.34	5,51,77,98	0
2	BOG	В	1007	20/20	0.85	0.24	25,55,63,64	0
2	BOG	A	1011	20/20	0.88	0.17	13,51,64,78	0
2	BOG	A	1041	12/20	0.89	0.23	13,58,66,72	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







Electron density around BOG B 1005:


 $2 {\rm mF}_o\text{-}{\rm DF}_c$ (at 0.7 rmsd) in gray ${\rm mF}_o\text{-}{\rm DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)














Electron density around BOG A 1001: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray ${ m mF}_o{ m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive) Electron density around BOG A 1033: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray mF_o -DF_c (at 3 rmsd) in purple (negative) and green (positive)










Electron density around BOG B 1037:


 $2 \text{mF}_o\text{-DF}_c$ (at 0.7 rmsd) in gray $\text{mF}_o\text{-DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)





















Electron density around BOG A 1029:

 $2 {\rm mF}_o\text{-}{\rm DF}_c$ (at 0.7 rmsd) in gray ${\rm mF}_o\text{-}{\rm DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around BOG A 1015:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)












Electron density around BOG B 1031: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray ${\rm mF}_o\text{-}{\rm DF}_c$ (at 3 rmsd) in purple (negative) and green (positive) Electron density around BOG B 1023: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray ${ m mF}_o{ m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)











Electron density around BOG A 1019: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray ${ m mF}_o{ m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive) Electron density around BOG A 1025: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray mF_o -DF_c (at 3 rmsd) in purple (negative) and green (positive)



Electron density around BOG A 1011: 2mF_o-DF_c (at 0.7 rmsd) in gray mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)



6.5 Other polymers (i)

There are no such residues in this entry.

